

DAFTAR PUSTAKA

- Addini, A. N. B. (2012). Sifat Fisikokimia Campuran Minyak Kelapa Sawit dan Minyak Wijen Selama Penggorengan Berulang Tempe Kedelai. Skripsi. Universitas Gadjah Mada.
- Aguilar-Hernandez, I., Afseth, N.K., Lopez-Luke, T., Contreras-Torres, F.F., Wold, J.P., Ornelas-Soto, N. (2017). Surface enhanced Raman spectroscopy of phenolic anti- oxidants: a systematic evaluation of ferulic acid, p-coumaric acid, caffeic acid and sinapic acid. *Vibrational Spectroscopy*, **89**, 113–122.
- Aladedunye, F., & Przybylski, R. (2013). Frying stability of high oleic sunflower oils as affected by composition of tocopherol isomers and linoleic acid content. *Food Chemistry*, **141**, 2373-2380.
- Ali, H.M., Abo-Shady, A., Sharaf Eldeen, H.A., Soror, H.A., Shousha, W.G., Abdel-Barry, O.A., Saleh, A.M. (2013). Structural features, kinetics and SAR study of radical scavenging and antioxidant activities of phenolic and anilinic compounds. *Chemistry Central Journal*, **7**, 53–53.
- Anonim, 1992. *Standard Nasional Indonesia: Mutu dan Cara Uji Minyak Kelapa*. Dewan Standardisasi Nasional, 1-3.
- Anonim, 2015. *Statistik Hortikultura Daerah Istimewa Yogyakarta 2015*. Yogyakarta: BPS Provinsi DIY
- Anonim, 2017. <http://tipspetani.blogspot.co.id/2014/09/jenis-markisa-yang-dibudidayakan.html>. Diakses pada tanggal 10 Juni 2017 jam 15.29 WIB
- Anwar, F., Hussain, A.I., Iqbal, S. & Bhangar, M.I. (2007). Enhancement of the oxidative stability of some vegetable oils by blending with Moringa oleifera oil. *Food Chemistry*, **103**, 1181–1191.
- AOAC, 2006. *Official Methods of Analysis of the Association of Official Analytical Chemist*. Washington: Association of Official Analytical Chemist.
- AOCS, 2009. *Official Methods of Analysis of the American Oil Chemists Society*. Washington: American Oil Chemists Society.

- Astuti, S.D., Andarwulan, N. & Hariyadi, P. (2010). Formulasi dan karakterisasi gel minyak sawit (Palm oil gel) kaya karotenoid sebagai ingredien pangan fungsional. Prosiding Seminar Nasional “Peran Keamanan Pangan Produk Unggulan Daerah dalam Menunjang Ketahanan Pangan dan Menekan Laju Inflasi”.
- Azad, A.K.M., Ali, M.A., Akter, M.S., Rahman, J. & Ahmed, M. (2014). Isolation and characterization of pectin extracted from lemon *pomace* during ripening, **2**, 30–35.
- Banerjee, A., Dasgupta, N. & De, B. (2005). In vitro study of antioxidant activity of *Syzygium cumini* fruit. *Food Chemistry*, **90**, 727–733.
- Bell, J.G., McEvoy, J., Tocher, D.R., Sargent, J.R. (2000). Depletion of α -tocopherol and astaxanthin in Atlantic salmon (*Salmo salar*) affects autoxidative defense and fatty acid metabolism. *Journal of Nutrition*, **130**, 1800-1808.
- Bezerra, M.A., Santelli, R.E., Oliveira, E.P., Villar, L.S. & Escalera, L.A. (2008). Response surface methodology (RSM) as a tool for optimization in analytical chemistry. *Talanta*, **76**, 965–977.
- Bhatnagar, A.S., Prasanth Kumar, P.K., Hemavathy, J. & Gopala Krishna, A.G. (2009). Fatty Acid Composition, Oxidative Stability, and Radical Scavenging Activity of Vegetable Oil Blends with Coconut Oil. *Journal of the American Oil Chemists' Society*, **86**, 991–999.
- Botella-Pav'ia, P. & Rodr'iguez-Concepci'on, M. (2006). Carotenoid biotechnology in plants for nutritionally improved foods. *Physiology Plant*, **126**, 369-381.
- Britton, G., Liaaen-Jensen, S., & Pfander, H. (1995). Carotenoids Volume 1. Switzerland: Birkhauser.
- Budiyanto, Silsia, D., Efendi, Z., & Janika, R. (2010). Perubahan kandungan β -karoten, asam lemak bebas dan bilangan peroksida minyak sawit merah selama pemanasan. *AGRITECH*, **30(2)**, 75-79.
- Cai, M., Wang, S. & Liang, H.H. (2012). Optimization of ultrasound-assisted ultrafiltration of *Radix astragalus* extracts with hollow fiber membrane using response surface methodology. *Separation and Purification Technology*, **100**, 74–81.
- Carocho, M., Ferreira, Isabel, C.F.R.. (2013). A review on antioxidants, prooxidants and related controversy: natural and synthetic compounds, screening and analysis methodologies. *Food and Chemical Toxicology*, **51**,

15-25.

- Chandran, J., Nayana, N., Roshini, N. & Nisha, P. (2017). Oxidative stability, thermal stability and acceptability of coconut oil flavored with essential oils from black pepper and ginger. *Journal of Food Science and Technology*, **54**, 144–152.
- Chandrashekar, P., Lokesh, B.R. & Krishna, A.G.G. (2010). Hypolipidemic effect of blends of coconut oil with soybean oil or sunflower oil in experimental rats. *Food Chemistry*, **123**, 728–733.
- Chu, Y.-H. and Kung, Y.-H. 1997. A study on oxidative stability of vegetable oil blends. *Shipin Kexue*, **24(4)**, 389-397.
- Chemat, F., Zill-E-Huma & Khan, M.K. (2011). Applications of ultrasound in food technology: Processing, preservation and extraction. *Ultrasonics Sonochemistry*, **18**, 813–835.
- Codex Standard. (1999). *Codex Standard for Edible Fats and Oils not Covered by Individual Standards* : Codex Stan 19-1981 (Rev.2-1999). London: FAO United Nations.
- Coppin, E. A., & Pike, O. A. (2001). Oil stability index correlated with sensory determination of oxidative stability in light-exposed soybean oil. *Journal of the American Oil Chemists' Society*, **78**, 13–18.
- Deng, J., Xu, Z., Xiang, C., Liu, J., Zhou, L., Li, T., Yang, Z. & Ding, C. (2017). Comparative evaluation of maceration and ultrasonic-assisted extraction of phenolic compounds from fresh olives. *Ultrasonics Sonochemistry*, **37**, 328–334.
- Dong, J., Liu, Y., Liang, Z. & Wang, W. (2010). Investigation on ultrasound-assisted extraction of salvianolic acid B from *Salvia miltiorrhiza* root. *Ultrasonics Sonochemistry*, **17**, 61–65.
- Foo, S.C., Yusoff, F.M., Ismail, M., Basri, M., Yau, S.K., Khong, N.M.H., Chan, K.W. & Ebrahimi, M. (2017). Antioxidant capacities of fucoxanthin-producing algae as influenced by their carotenoid and phenolic contents. *Journal of Biotechnology*, **241**, 175–183.
- Gharby, S., Harhar, H. & Charrouf, Z. (2017). Chemical characterization and oxidative stability of seeds and oil of sesame grown in Morocco. *Journal of the Saudi Society of Agricultural Sciences*, **16**, 105–111.
- Gorinstein, S., Poovarodom, S., Leontowicz, H., Leontowicz, M., Namiesnik, J., Vearasilp, S., Haruenkit, R., Ruamsuke, P., Katrich, E., Tashma, Z. (2011). Antioxidant properties and bioactive constituents of some rare

- exotic Thai fruits and comparison with conventional fruits: in vitro and in vivo studies. *Food Research International*, **44**, 2222-2232.
- Goula, A.M. (2013). Ultrasound-assisted extraction of pomegranate seed oil - Kinetic modeling. *Journal of Food Engineering*, **117**, 492-498.
- Goula, A.M., Ververi, M., Adamopoulou, A. & Kaderides, K. (2017). Green ultrasound-assisted extraction of carotenoids from pomegranate wastes using vegetable oils. *Ultrasonics Sonochemistry*, **34**, 821-830.
- Gulla, S. & Waghray, K. (2011). Effect of Storage on Physico-chemical Characteristics and Fatty Acid Composition of Selected Oil Blends, **3**, 35-46.
- Hammouda, I. Ben, Triki, M., Matthäus, B. & Bouaziz, M. (2018). A Comparative Study on Formation of Polar Components, Fatty Acids and Sterols during Frying of Refined Olive Pomace Oil Pure and Its Blend Coconut Oil. *Journal of Agricultural and Food Chemistry*, **66**, 3514-3523.
- Handajani, S., Erlyna W.R dan Suminah Anantanya. (2006). *The Queen of Oil, Potensi Agribisnis Komoditas Wijen*. Andi: Yogyakarta.
- Handajani, S., Manuhara, G.J. & Anandito, R.B.K. (2010). Pengaruh suhu ekstraksi terhadap karakteristik fisik, kimia dan sensoris minyak wijen, **30**, 116-122.
- Handayani, A.D., Sutrisno, Indraswati, N. & Ismadji, S. (2008). Extraction of astaxanthin from giant tiger (*Panaeus monodon*) shrimp waste using palm oil: Studies of extraction kinetics and thermodynamic. *Bioresource Technology*, **99**, 4414-4419.
- Hashempour-baltork, F., Torbati, M., Azadmard-damirchi, S. & Savage, G.P. (2016). Trends in Food Science & Technology Vegetable oil blending: A review of physicochemical, nutritional and health effects. *Trends in Food Science & Technology*, **57**, 52-58.
- Jasna, C., Pajin, B., Djilas, S., Šaponjac, V.T., Gordana, C., Lonc, I. & Vulic, J. (2016). Sour cherry pomace extract encapsulated in whey and soy proteins: Incorporation in cookies. *Food Research International*, **207**, 27-33.
- Janzantti, N.S., Macoris, M.S., Garruti, D.S. & Monteiro, M. (2012). Influence of the cultivation system in the aroma of the volatile compounds and total antioxidant activity of passion fruit. *LWT - Food Science and Technology*, **46**, 511-518.
- Kalam, M.A., Rashed, M.M., Imdadul, H.K. & Masjuki, H.H. (2016). Property

development of fatty acid methyl ester from waste coconut oil as engine fuel. *Industrial Crops and Products*, **87**, 333–339.

Ketaren, S. (2008). *Minyak dan Lemak Pangan*. Jakarta : UI Press.

Kivevele, T., Agarwal, A.K., Gupta, T. & Mbarawa, M. (2011). Oxidation Stability of Biodiesel Produced from Non-Edible Oils of African Origin. *SAE International Technical Paper*.

Kulkarni, S.G. & Vijayanand, P. (2010). Effect of extraction conditions on the quality characteristics of pectin from passion fruit peel (*Passiflora edulis* f. *flavicarpa* L.). *LWT - Food Science and Technology*, **43**, 1026–1031.

Kumar, P.K.P. & Krishna, A.G.G. (2015). Physicochemical characteristics of commercial coconut oils produced in India. *Grasas Aceites*, 66:11.

Lee, E. & Choe, E. (2012). Biocatalysis and Agricultural Biotechnology Changes in oxidation-derived off-flavor compounds of roasted sesame oil during accelerated storage in the dark. *Biocatalysis and Agricultural Biotechnology*, **1**, 89–93.

Lee, J., Lee, Y. & Choe, E. (2008). Effects of sesamol, sesamin, and sesamolin extracted from roasted sesame oil on the thermal oxidation of methyl linoleate. *LWT - Food Science and Technology*, **41**, 1871–1875.

Lenth, R. V. (2012). Response-surface methods in R , using rsm. *Journal of Statistical Software*, **32**, 1–17.

Li, H., Zhou, P., Yang, Q., Shen, Y., Deng, J., Li, L. & Zhao, D. (2011). Comparative studies on anxiolytic activities and flavonoid compositions of *Passiflora edulis* “edulis” and *Passiflora edulis* “flavicarpa.” *Journal of Ethnopharmacology*, **133**, 1085–1090.

Li, Y., Fabiano-Tixier, A.S., Tomao, V., Cravotto, G. & Chemat, F. (2013). Green ultrasound-assisted extraction of carotenoids based on the bio-refinery concept using sunflower oil as an alternative solvent. *Ultrasonics Sonochemistry*, **20**, 12–18.

Lópes-Vargas, J.H., Fernández-Lópes, J., Pérez-Álvarez, J.A., Viuda-Martos M. (2013) Chemical, physico-chemical, technological, antibacterial and antioxidant properties of dietary fiber powder obtained from yellow passion fruit (*Passiflora edulis* var. *flavicarpa*) co-products. *Food Research International*, **51**, 756–763.

Luque-García, J.L. & De Castro, M.D.L. (2003). Ultrasound: a powerful tool for leaching. *Trends Analytical Chemistry*, **22**, 41–47.

- Malacrida, C.R. & Jorge, N. (2012). Yellow passion fruit seed oil (*Passiflora edulis* f. *flavicarpa*): Physical and chemical characteristics. *Brazilian Archives of Biology and Technology*, **55**, 127–134.
- Mathur, A., Verma, S.K., Purohit, R., Gupta, V., Dua, V.K., Prasad, G. B.K.S., Mathur, D., Singh, S. K., & Singh, S. (2011). Evaluation of in vitro antimicrobial and antioxidant activities of peel and pulp of some citrus fruits. *IJPI'S Journal of Biotechnology and Biotherapeutics*, **1:2**, 1-17.
- McGrady, J. 1993. Fats and Oils. IFT Short Course Ingredients Technology. Chicago:IL.
- Meléndez-martínez, A.J., Escudero-gilete, M.L., Vicario, I.M. & Heredia, F.J. (2010). Study of the influence of carotenoid structure and individual carotenoids in the qualitative and quantitative attributes of orange juice colour, **43**, 1289–1296.
- Mercadante, A.Z., Britton, G. & Rodriguez-Amaya, D.B. (1998). Carotenoids from Yellow Passion Fruit (*Passiflora e dulis*). *Journal of Agricultural and Food Chemistry*, **46**, 4102–4106.
- Metrohm, A. G. (2013). Oxidative stability of oils and fats—Rancimat method. *Metrohm application bulletin*, **204/2 e**, 1–4.
- Munhoz, D.R., Moreira, F.K., Bresolin, J.D., Bernardo, M.P., Sousa, C.P. De & Mattoso, L.H.C. (2018). Sustainable Production and In vitro Biodegradability of Edible Films from Yellow Passion Fruit Co-products via Continuous Casting. *ACS Sustainable Chemistry & Engineering*, **6**, 9883–9892.
- Ningrum, A., Minh, N.N., & Schreiner, M. (2015). Carotenoids and Norisoprenoids as Carotenoid Degradation Products in Pandan Leaves (*Pandanus amaryllifolius* Roxb.). *International Journal of Food Properties*, **18**, 1905–1914.
- Nishino, A., Yasui, H. & Maoka, T. (2017). Reaction and Scavenging Mechanism of β -Carotene and Zeaxanthin with Reactive Oxygen Species, **2016**, 1–8.
- Nitbani, F.O., Jumina, Siswanta, D. & Solikhah, E.N. (2016). Isolation and Antibacterial Activity Test of Lauric Acid from Crude Coconut Oil (*Cocos nucifera* L.). *Procedia Chemistry*, **18**, 132–140.
- Nugraha, W.S. 2004. Kendali Adsorben Karbon Aktif dan Magnesium Silikat dalam Efisiensi Pemakaian Minyak Goreng di *Further Processing* PT.Chaeroen Pokhand Indonesia-Serang. Skripsi. Institut Pertanian Bogor.

- Oliver, J. & Palou, A. (2000). Chromatographic determination of carotenoids in foods. *Journals of Chromatography A*, **881**, 543-555.
- Oliveira, R.C. de, Davantel De Barros, S.T. & Gimenes, M.L. (2013). The extraction of passion fruit oil with green solvents. *Journal of Food Engineering*, **117**, 458–463.
- Oliveira, A.C. de, Valentim, I.B., Silva, C.A., Bechara, E.J.H., Barros, M.P. de, Mano, C.M. & Goulart, M.O.F. (2009). Total phenolic content and free radical scavenging activities of methanolic extract powders of tropical fruit residues. *Food Chemistry*, **115**, 469–475.
- Oliveira, D.A., Angonese, M., Gomes, C.F., & Ferreira, S.R.S. (2016). Valorization of passion fruit (*Passiflora edulis* sp.) by-products: Sustainable recovery and biological activities. *Journal of Supercritical Fluids*, **111**, 55–62.
- Ou, B., Huang, D., Hampsch-Woodill, M., Flanagan, J.A. & Deemer, E.K. (2002). Analysis of antioxidant activities of common vegetables employing oxygen radical absorbance capacity (ORAC) and ferric reducing antioxidant power (FRAP) assays: A comparative study. *Journal of Agricultural and Food Chemistry*, **50**, 3122–3128.
- Ovani, I. (2013). Pengembangan minuman emulsi minyak bekatul berflavor kaya Antioksidan untuk pencegahan penyakit tidak menular. Skripsi. Institut Pertanian Bogor.
- Owen, J. P. (2015). Fruit and Pomace Extracts: Biological Activity, Potential Applications and Beneficial Health Effects. New York: Nova Publishers.
- Pabari, R.M. & Ramtoola, Z. (2012). Application of face centred central composite design to optimise compression force and tablet diameter for the formulation of mechanically strong and fast disintegrating orodispersible tablets. *International Journal of Pharmaceutics*, **430**, 18–25
- Park, P. W. & Goins, R. E. (1994). In situ preparation of fatty acids methyl ester for analysis of fatty acids composition in food. *Journal of Food Science*, **59**, 1262-1266.
- Paula, a N. a, Aukar, D.E.A., Gertrudes, E., Lemos, D.E.M. & Oliveira, J.C. (2002). Genetic Variations Among Passion Fruit Species Using Rapid Markers 1. *Biologia*, **1**, 738–740.
- Pereira, A.D., Correa, R.X., Oliveira, A.C.C. (2015). Molecular genetic diversity and differentiation of populations of ‘somnus’ passion fruit trees

- (*Passiflora setacea* DC): Implications for conservation and pre-breeding. *Biochemical System and Ecosystem*, **59**, 12-21.
- Pertuzatti, P.B., Sganzerla, M., Jacques, A.C., Barcia, M.T. & Zambiasi, R.C. (2015). Carotenoids, tocopherols and ascorbic acid content in yellow passion fruit (*Passiflora edulis*) grown under different cultivation systems. *LWT - Food Science and Technology*, **64**, 259–263.
- Petrauskaite, V., De Greyt, W., Kellens, M. & Huyghebaert, A. (1998). Physical and chemical properties of trans- free fats produced by chemical interesterification of vegetable oil blends. *Journal of the American Oil Chemists' Society*, **75**, 489- 493.
- Pinheiro, E.R., Silva, I.M.D.A., Gonzaga, L. V., Amante, E.R., Teófilo, R.F., Ferreira, M.M.C. & Amboni, R.D.M.C. (2008). Optimization of extraction of high-ester pectin from passion fruit peel (*Passiflora edulis flavicarpa*) with citric acid by using response surface methodology. *Bioresource Technology*, **99**, 5561–5566.
- Porto-Figueira, P., Freitas, A., Cruz, C.J., Figueira, J. & Câmara, J.S. (2015). Profiling of passion fruit volatiles: An effective tool to discriminate between species and varieties. *Food Research International*, **77**, 408–418.
- Ramli, N.S., Ismail, P. & Rahmat, A. (2014). Influence of Conventional and Ultrasonic-Assisted Extraction on Phenolic Contents , Betacyanin Contents , and Antioxidant Capacity of Red Dragon Fruit (*Hylocereus polyrhizus*). *The Scientific World Journal*, **2014**, 1-7.
- Reena, MB. & Lokesh, B.R. (2007). Hypolipidemic Effect of Oils with Balanced Amounts of Fatty Acids Obtained by Blending and Interesterification of Coconut Oil with Rice Bran Oil or Sesame Oil. *Journal of Agricultural and Food Chemistry*, **55**, 10461–10469.
- Rodriguez-Amaya, D. B. (2001). A guide to carotenoid analysis in food. Washington, DC: ILSI Press.
- Rodriguez-amaya, D.B. (2010). Journal of Food Composition and Analysis Quantitative analysis , in vitro assessment of bioavailability and antioxidant activity of food carotenoids — A review. *Journal of Food Composition and Analysis*, **23**, 726–740.
- Sachindra, N.M. & Mahendrakar, N.S. (2005). Process optimization for extraction of carotenoids from shrimp waste with vegetable oils. *Bioresource Technology*, **96**, 1195–1200.
- Saini, R.K. & Keum, Y. (2018). Carotenoid extraction methods : A review of

recent developments. *Food Chemistry*, **240**, 90–103.

- Santos, D.T., Cavalcanti, R.N., Rostagno, M.A., Queiroga, C.L., Eberlin, M.N., Meireles, M.A.A. (2013). Extraction of Polyphenols and Anthocyanins from the Jambul (*Syzygium cumini*). *Fruit Food & Publication Health* **3**, 12-20.
- Serjouie, S., Tan, C. P., Mirhosseini, H., & Che Man, Y. B. (2010). Effect of vegetable- based oil blends on physicochemical properties of oils during deep fat-frying. *American Journal of Food Technology*, **5**, 310-323.
- Shimizu, S., Fujii, G., Takahashi, M., Nakanishi, R., Komiya, M., Shimura, M., Noma, N., Onuma, W., Terasaki, M., Yano, T., & Mutoh, M. (2014). Sesamol suppresses cyclooxygenase transcriptional activity in colon cancer cells and modifies intestinal polyp development in *Apc*^{Min/+} mice. *Journal of Clinical Biochemistry and Nutrition*, **54**, 95-101.
- Silva, J.K. da, Cazarin, C.B.B., Batista, Āngela Giovana & Maróstica, M. (2014). Effects of passion fruit (*Passiflora edulis*) byproduct intake in antioxidant status of Wistar rats tissues. *LWT - Food Science and Technology*, **59**, 1213–1219.
- Silva, J.K. da, Cazarin, C.B.B., Colomeu, T.C., Batista, Āngela Giovana, Meletti, L.M.M., Paschoal, J.A.R., Bogusz Júnior, S., Furlan, M.F., Reyes, F.G.R., Augusto, F., Maróstica Júnior, M.R. & Lima Zollner, R. de. (2013). Antioxidant activity of aqueous extract of passion fruit (*Passiflora edulis*) leaves: In vitro and in vivo study. *Food Research International*, **53**, 882–890.
- Silva, L.M.R. Da, Figueiredo, E.A.T. De, Ricardo, N.M.P.S., Vieira, I.G.P., Figueiredo, R.W. De, Brasil, I.M. & Gomes, C.L. (2014). Quantification of bioactive compounds in pulps and by-products of tropical fruits from Brazil. *Food Chemistry*, **143**, 398–404.
- Sun, M. & Temelli, F. (2006). Supercritical carbon dioxide extraction of carotenoids from carrot using canola oil as a continuous co-solvent. *Journal of Supercritical Fluids*, **37**, 397–408.
- Sunil, L., Reddy, P. V., Krishna, A. G. G., & Urooj, A. (2015). Retention of natural antioxidants of blends of groundnut and sunflower oils with minor oils during storage and frying. *Journal of Food Science and Technology*, **52**, 849-857.
- Taghvaei, M., Jafari, S.M., Mahoonak, A.S., Nikoo, A.M., Rahmadian, N., Hajitabar, J., Meshginfar, N. (2014). The effect of natural antioxidants extracted from plant and animal resources on the oxidative stability of soybean oil. *LWT-Food Science & Technology*, **56**, 124–130.

- Tan, C.P., Man, Y.B.C., Selamat, J. & Yusoff, A. (2002). Analytical, Nutritional and Clinical Methods Section Comparative studies of oxidative stability of edible oils by differential scanning calorimetry and oxidative stability index methods. *Food Chemistry*, **76**, 385–389.
- Tapiero, H., Townsend, D.M. & Tew, K.D. (2004). The role of carotenoids in the prevention of human pathologies, **58**, 100–110.
- Tiwari, M.R., Tiwari, K.K., & Toliwal, S.D. 2014. Studies on Thermal Stability of Palm-Sesame Oil Blends during Deep-Fat Frying. *Journal of Scientific and Industrial Research*, **73**, 153-156.
- Vicario, I.M., Heredia, F.J. & Mele, A.J. (2007). Food Chemistry Provitamin A carotenoids and ascorbic acid contents of the different types of orange juices marketed in Spain, **101**, 177–184.
- Viganó, J. & Martinez, J. (2015). Trends for the Application of Passion Fruit Industrial By-products: A Review on the Chemical Composition and Extraction Techniques of Phytochemicals. *Food and Public Health*, **5**, 164–173.
- Wahyuni, D.T. & Widjanarko, S.B. (2015). The Effect of Different Solvent and Extraction Time of Carotenoids Extract From Pumpkin with Ultrasonic Method. *Jurnal Pangan dan Agroindustri*, **3**, 390–401.
- Widiyarti, G., Hanafi, M., Soewarso, W.P. (2009). Study on synthesis of monolaurin as antibacterial agent against *Staphylococcus aureus*. *Indonesian Journal of Chemistry*, **1**, 99-106.
- Yates, M., Ramos, M., Martin-luengo, M.A., Maria, A., Serrano, M. & Zurdo, V. (2017). MultivalORIZATION of apple *pomace* towards materials and chemicals . *Waste to wealth*, **143**.
- Yen, F., Fan, K., He, J. & Gao, M. (2015). Ultrasonic-Assisted Solvent Extraction of Carotenoids From Rapeseed Meal : Optimization Using Response Surface Methodology. *Journal of Food Quality*, **38**, 377–386.
- Yen, G.C. & Shyu, S.L. (1989). Oxidative stability of sesame oil prepared from sesame seed with different roasting temperatures. *Food Chemistry*, **31**, 215–224.
- Yoo, K. & Moon, B. (2016). Comparative carotenoid compositions during maturation and their antioxidative capacities of three citrus varieties. *Food Chemistry*, **196**, 544–549.

Zribi, A., Jabeur, H., Matthäus, B. & Bouaziz, M. (2016). Quality control of refined oils mixed with palm oil during repeated deep-frying using FT-NIRS, GC, HPLC, and multivariate analysis. *European Journal of Lipid Science and Technology*, **118**, 512–523.