

DAFTAR PUSTAKA

- Auldridge M. E., Block A., Vogel J. T., Dabney-Smith C., Mila I., Bouzayen M., Magallanes-Lundback M., DellaPenna D., McCarty D. R. & Klee H. J. (2006). Characterization of three members of the *Arabidopsis* carotenoid cleavage dioxygenase family demonstrates the divergent roles of this multifunctional enzyme family. *The Plant Journal* 45, 982-993
- Auldridge M. E., McCarty D. R. & Klee H. J. (2006) Plant carotenoid cleavage oxygenases and their apocarotenoid products. *Current Opinion in Plant Biology* 9, 315-321.
- AOAC. (2006). Official Methods of Analysis. 11th edition. Washington DC: Association of Official Analytical Chemists, Inc.
- Ayala-Zavala, J. F., Vega-Vega, V., Rosas-Domínguez, C., Palafox-Carlos, H., Villa-Rodríguez, J. a., Siddiqui, M. W., González-Aguilar, G. A. (2011). Agro-industrial potential of exotic fruit byproducts as a source of food additives. *Food Research International*, 44(7), 1866–1874. doi:10.1016/j.foodres.2011.02.021
- Bachrudin, Z. 1999. *Petunjuk Laboratorium: Isolasi, Identifikasi, dan Pewarnaan Protein*. Yogyakarta: PAU Bioteknologi UGM.
- Baldermann, S., Naim, M. & Fleischmann, P. (2005). Enzymatic carotenoid degradation and aroma formation in nectarines (*Prunus persica*), **38**, 833–836.
- Baldermann, S. (2008) Carotenoid oxygenases from *Camellia sinensis*, *Osmanthus fragrans*, and *Prunus persica nucipersica*. *Cuvillier Verlag Göttingen*.
- Baldermann, S. et al. (2010) Functional characterization of a carotenoid cleavage dioxygenase 1 and its relation to the carotenoid accumulation and volatile emission during the floral development of *Osmanthus fragrans* Lour. *J. Exp. Bot.* 61, 2967–2977
- Banerjee, J., Singh, R., Vijayaraghavan, R., Macfarlane, D., Patti, A. F., dan Arora, A. (2016). Bioactives From Fruit Processing Wastes: Green Approaches to Valuable Chemicals. *Food Chemistry*, 5(6), 333-383
- Baumes R., Wirth J., Bureau S., Gunata Y. & Razungles A. (2002) Biogenesis of C13- norisoprenoid compounds: experiments supportive for an apocarotenoid pathway in grapevines. *Analytica Chimica Acta* 458, 3-14.

- Britton, G., Liaaen-Jensen, S., Pfander, H. (1995) Carotenoids Volume 1. *Birkhauser, Basel, Switzerland*.
- Britton, G., Liaaen-Jensen, S., Pfander, H. (2008) Carotenoids Volume 4 – Natural functions. *Birkhauser, Basel, Switzerland*.
- Bouvier, F., Suire, C., Mutterer, J., Camara, B. (2003) Oxidative remodeling of chromoplast carotenoids: identification of the carotenoid dioxygenase *CsCCD* and *CsZCD* genes involved in crocus secondary metabolite biogenesis. *Plant Cell* 15: 47–62.
- Chaplin, M., dan Bucke, C. (1990). Enzyme Technology. Cambridge: Cambridge University Press
- Chau, C.F. & Huang, Y.L. (2004). Characterization of passion fruit seed fibres - A potential fibre source. *Food Chemistry*, **85**, 189–194.
- Chernys, J.T., Zeevaart, J.A.D., 2000. Characterization of the 9-cis-epoxycarotenoid dioxygenase gene family and the regulation of abscisic acid biosynthesis in avocado. *Plant Physiol.* 124, 343–353.
- Dunn, m.J. 1989. Determination of total Protein Concentration . di dalam : Protein Purification Methods. Harris, E.L.V. dan S. Angal (eds.). IRL Press, Oxford, England.
- Ferrari, R.A. et al. (2014). Characterization of by-products of passion fruit industrialization - utilization of seeds. **Revista Brasileira de Fruticultura**, v.26, n.1, p.101-102.
- Fleischmann, P., Lutz-Roeder, A., Winterhalter, P., Watanabe, N., (2001). Carotenoid cleavage enzymes in animals and plants. In: Winterhalter, P., Rouseff, R. (Eds.), Carotenoid-Derived Aroma Compounds. ACS Symp. Ser. 802, Vol. 219. American Chemical Society, Washington, DC, pp. 76–88.
- Fleischmann, P., Studer, K., Winterhalter, P., 2002. Partial purification and kinetic characterization of a carotenoid cleavage enzyme from quince fruit (*Cydonia oblonga*). *J. Agric. Food Chem.* 50, 1677–1680.
- Gonzalez-Jorge, S. et al. (2013) Carotenoid Cleavage Dioxygenase4 is a negative regulator of b-carotene content in *Arabidopsis seeds*. *Plant Cell* 25, 4812–4826
- Guiné, R. P., Martín-Belloso, O., Mínguez-Mosquera, M. I., Paliyath, G., Pessoa, F. L., Le Quéré, J.-L., Sidhu, J. S., Sinha, N., Peggy Stanfield, R., & Hui, Y. (2010). Handbook of fruit and vegetable flavors: John Wiley and Sons.

- Hames, B. D., dan Rickwood, (2004). *A Practical Approach : Gel Electrophoresis Protein*. Huntington: Robert E Krieger Publishing Company.
- Huang, F.C., Horváth, G., Molnár, P., Turcsi, E., Deli, J., Schrader, J., Sandmann, G., Schmidt, H., Schwab, W. (2009) Substrate promiscuity of RdCCD1, a carotenoid cleavage oxygenase from *Rosa damascena*. *Phytochem.* 70 (4): 457-464.
- Ilg A., Beyer P. & Al-Babili S. (2009) Characterization of the rice carotenoid cleavage dioxygenase1 reveals a novel route for geranial biosynthesis. *FEBS Journal* 3, 736-747.
- Isoe S.; Katsumura S.; Hyeon, S. B.; Sakan, T. Biogenetic type synthesis of grasshopper ketone and loliolide and a possible biogenesis of allenic carotenoids. *Tetrahedron Lett.* **1971**, 1089-1092.
- Janzantti, N.S., Macoris, M.S., Garruti, D.S. & Monteiro, M. (2012). Influence of the cultivation system in the aroma of the volatile compounds and total antioxidant activity of passion fruit. *LWT - Food Science and Technology*, **46**, 511–518.
- Janzantti, N.S. & Monteiro, M. (2014). Changes in the aroma of organic passion fruit (*Passiflora edulis* Sims f. *flavicarpa* Deg.) during ripeness. *LWT - Food Science and Technology*, **59**, 612–620.
- Kato, M., Matsumoto, H., Ikoma, Y., Okuda, H., Yano, M. (2006) The role of carotenoid cleavage dioxygenases in the regulation of carotenoid profiles during maturation of citrus fruit. *J. Exp. Bot.* 57: 2153–2164.
- Leão, K.M.M., Sampaio, K.L., Pagani, A.A.C. & Silva, M.A.A.P. Da. (2014). Odor potency, aroma profile and volatiles composition of cold pressed oil from industrial passion fruit residues. *Industrial Crops and Products*, **58**, 280–286.
- Leffingwell, J. (2014). Carotenoids as Flavor & Fragrance Precursors. A Review. *Leffingwell Reports*, Vol. 2 (No. 3)
- Lehninger, A.L., 1992, *Principles of Biochemistry*: 1st Edition, Worth Pub, New York.
- Lewinshon, E., Sitrit, Y., Bar, E., Azulay, Y., Ibdah, M., Meir, A., Yosef, E., Zamir, D., Tadmor, Y. (2005). Not just color carotenoid degradation as a link between pigmentation and aroma in tomato and watermelon fruit. *Trends Food Science Technology*. 16, 407-415

- Liu Jia Ren , Hong-Wei Dong , Xiang-Rong Sun , Qi Wang , Wen-Guang Sun , John W. Parry , Qian Liu , Xiao-Hui Han , Chang-Hao Sun , Bing-Qing Chen & Bao-Feng Yang (2009) Effects of β -Ionone on Mammary Carcinogenesis and Antioxidant Status in Rats Treated With DMBA, *Nutrition and Cancer*, 62:1, 58-65
- Ma, G., Zhang, L., Matsuta, A., Matsutani, K., Yamawaki, K., Yahata, M., Kato, M. (2013). Enzymatic formation of β -citraurin from β -cryptoxanthin and zeaxanthin by carotenoid cleavage dioxygenase 4 in the flavedo of citrus fruit. *Plant Physiology*, 163(2), 682-695
- Mathieu S., Terrier N., Procureur J., Bigey F. & Günata Z. (2005) Carotenoid cleavage dioxygenase from *Vitis vinifera* L.: functional characterization and expression during grape berry development in relation to C13 norisoprenoid accumulation. *Journal of Experimental Botany* 56, 2721-2731
- McCarty, D.R., Klee, H.J. (2006) Characterization of three members of the *Arabidopsis* carotenoid cleavage dioxygenase family demonstrates the divergent roles of this multifunctional enzyme family. *Plant. J.* 45: 982–993.
- Mendes-Pinto M. M. (2009) Carotenoid breakdown products: the “norisoprenoids” in wine aroma. *Archives of Biochemistry and Biophysics* 483, 236-245.
- Mercadante, A.Z., Britton, G. & Rodriguez-Amaya, D.B. (1998). Carotenoids from Yellow Passion Fruit (*Passiflora e dulis*). *Journal of Agricultural and Food Chemistry*, **46**, 4102–4106.
- Moraes, T. De, Vanessa, C., Hermes, S., Oliveira, A. De, Flo, S. H. (2016). Evaluation of bioactive compounds, chemical and technological properties of fruits byproducts powder. *Journal of Food Science and Technology*, 20(12), 100-102
- Morais, L., Altina, E., Figueiredo, T. De, Maria, N., Silva, P., Gusmao, L., Gomes, C.L. (2014). Quantification of bioactive compounds in pulps and by-products of tropical fruits from Brazil. *Food Chemistry*, 143, 398-404.
- Nacke, C., Hüttmann, S., Etschmann, M.M.W. & Schrader, J. (2012). Enzymatic production and in situ separation of natural β -ionone from β -carotene. *Journal of Industrial Microbiology and Biotechnology*, **39**, 1771–1778.
- Ningrum, A. (2010). Carotenoids, related aroma compounds and partial purification of a carotenoid cleavage enzymes from Aonori

(*Enteromorpha sp.*). Master Thesis. Faculty of Agriculture, Shizuoka University, Shizuoka, Japan

- Ningrum, A., Minh, N. N., & Schreiner, M. (2015). Carotenoids and Norisoprenoids as Carotenoid Degradation Products in Pandan Leaves (*Pandanus amaryllifolius* Roxb.). *International Journal of Food Properties*, 18(9), 1905–1914. doi:10.1080/10942912.2014.971186
- Ningrum, A., & Schreiner, M. (2014). Carotenoid-Cleavage Activities of Crude Enzymes from Pandanus amryllifolius. *Chemistry & Biodiversity*, 11, 1871–1881.
- Ohmiya, A. (2009) Carotenoid cleavage dioxygenases and their apocarotenoid products in plants. *Plant Biotech.* 26: 351–358.
- Oliveira, D.A., Angonese, M., Gomes, C. & Ferreira, S.R.S. (2016). The Journal of Supercritical Fluids Valorization of passion fruit (*Passiflora edulis* sp.) by-products : Sustainable recovery and biological activities. *The Journal of Supercritical Fluids*, 111, 55–62.
- Ruiz-Sola, M.A. and Rodriguez-Concepcion, M. (2012) Carotenoid biosynthesis in Arabidopsis: a colorful pathway. *Arabidopsis Book* 10, e0158
- Rodriguez-Amaya, D. B. (2001). A guide to carotenoid analysis in foods. Washington, DC: ILSI Press64
- Pertuzatti, P.B., Sganzerla, M., Jacques, A.C., Barcia, M.T. & Zambiasi, R.C. (2015). Carotenoids, tocopherols and ascorbic acid content in yellow passion fruit (*Passiflora edulis*) grown under different cultivation systems. *LWT - Food Science and Technology*, 64, 259–263.
- Pontes, M., Marques, J.C. & Câmara, J.S. (2009). Headspace solid-phase microextraction-gas chromatography-quadrupole mass spectrometric methodology for the establishment of the volatile composition of *Passiflora* fruit species. *Microchemical Journal*, 93, 1–11.
- Qin, X.Q., Zeevaart, J.A.D., 1999. The 9-cis-epoxycarotenoid cleavage reaction is the key regulatory step of abscisic acid biosynthesis in water-stressed bean. *Proc. Natl. Acad. Sci. USA* 96, 15354– 15361.
- Rotili, M. C. C., Vorpagel, J. A., Braga, G. C., Kuhn, O. J., & Salibe, A. B. (2013). Antioxidant activity, chemical composition and conservation of yellow passion fruit packed with pvc film. *Revista Brasileira de Fruticultura*, 35(4), 942-952.

- Schwartz S. H., Qin X. Q. & Loewen M. C. (2004) The biochemical characterization of two carotenoid cleavage enzymes from *Arabidopsis* indicates that a carotenoid-derived compound inhibits lateral branching. *Journal of Biological Chemistry* 279, 46940-46945.
- Schwartz, S.H., Qin, X.Q., Zeevaart, J.A.D., 2001. Characterization of a novel carotenoid cleavage dioxygenase from plants. *J. Biol. Chem.* 276, 25208–25211b.
- Schwartz, S.H., Tan, B.C., Gage, D.A., Zeevaart, J.A.D., McCarty, D.R., 1997. Specific oxidative cleavage of carotenoids by VP14 of maize. *Science* 276, 1872–1874.
- Scopes, R.K., 1987, *Protein Purification, Principle and Practice*, 2nd ed, Springer Verlag, New York.
- Septembre-Malaterre, A., Stanislas, G., Douraguia, E. & Gonthier, M.P. (2016). Evaluation of nutritional and antioxidant properties of the tropical fruits banana, litchi, mango, papaya, passion fruit and pineapple cultivated in Réunion French Island. *Food Chemistry*, **212**, 225–233.
- Simkin, A.J., Underwood, B.A., Auldridge, M., Loucas, H.M., Shibuya, K., Schmelz, E., Clark, D.G., Klee, H.J. (2004) Circadian regulation of the PhCCD1 carotenoid cleavage dioxygenase controls emission of β -ionone, a fragrance volatile of petunia flowers *Plant. Physiol.* 136: 3504-3514.
- Silva, T.V. et al. (2014) Determination of the peel color scale and juice yield of yellow passion fruits in different harvest seasons. *Revista Brasileira de Fruticultura*, v.30, n.4, p.880-884.
- Sudarmadji, S., Haryono, B., Suhardi. 1996. *Analisa Bahan Makanan dan Pertanian*. Yogyakarta: Penerbit Liberty.
- Sudha, M, L., Dharmesh, S.M., Pynam, H., and Bhimangouder, S.V. (2016). Antioxidant and cyto/DNA protective properties of apple pomace enriched bakery product. *Jurnal of Food Science and Technology*, 53(4), 1909-1918.
- Talcott, S. T., Percival, S. S., Pittet-Moore, J., & Celoria, C. (2003). Phytochemical composition and antioxidant stability of fortified yellow passion fruit (*Passiflora edulis*). *Journal of Agricultural and Food Chemistry*, 51, 935e941.
- Vogel, J.T., Tan, B.C., McCarty, D.R., Klee, H.J. (2008) The carotenoid cleavage dioxygenase 1 enzyme has broad substrate specificity, cleaving multiple carotenoids at two different bond positions. *J. Bio. Chem.* 283 (17): 11364-11373.

- Walter, M.H. and Strack, D. (2011) Carotenoids and their cleavage products: biosynthesis and functions. *Nat. Prod. Rep.* 28, 663–692
- Winterhalter P, Rouseff R (2002) Carotenoid-derived aroma compounds: an introduction In: Winterhalter P, Rouseff R (eds) ACS Symposium Series American Chemical Society. Washington, USA, pp 1–17
- Wijaya, S.K.S., & Rohman, L. 2005. *Fraksinasi dan Karakterisasi protein Utama Biji Kedelai*. Jember: Fakultas MIPA Universitas Jember.
- Wilson K. Walker 2010. Protein and enzyme techniques *In Practical Biochemistry*, (ed. Wilson K and Walker JM), Cambridge University Press. p.161-226.
- Yahyaa, M., Berim, A., Isaacson, T., Marzouk, S., Bar, E., Lewinshon (2015) Isolation and functional characterization of carotenoid cleavage dioxygenase-1 from *Laurus nobilis* L. (bay laurel) fruits. *J. Agric. Food Chem.* 63, 8275–82823.
- Yu SG, Anderson PJ, and Elson CE. (2012). Efficacy of β -ionone in the chemoprevention of rat mammary carcinogenesis. *J Agric Food Chem* **43**, 2144–2147,