

DAFTAR PUSTAKA

- Ackman, R.G. (1989). Nutritional composition of fats in seafoods. *Prog. Food Nutr. Sci.* 13: 161-289.
- Ahmed, Z., Donkor, O., Street, W.A., Vasiljevic, T. (2013). Proteolytic activities in fillets of selected underutilized Australian fish species. *Food Chem.* 140: 238–244.
- Alvarez, C., Couso, I., Tejada, I. (1999). Thermal gel degradation (“modori”) in sardine surimi gels. *J. Food Sci.* 64: 663-637.
- Ando, H., Adachi, M., Umeda, K., Matsuura, A., Nonaka, M., Uchio, R., Tanaka, H., Motoki, M. (1989). Purification and Characteristics of a Novel Transglutaminase Derived from Microorganisms. *Agric. Biol. Chem.* 53: 2613–2617.
- Ando, M., Tsukamasa, Y., Makinodan, Y. (1998). Identification of 170 k component which appears in the setting process of surimi gel. *Fish. Sci.* 64, 497–498.
- An, H., Weerasinghe, V., Seymour, T.A., Morrissey, M.T. (1994). Cathepsin degradation of Pacific whiting surimi proteins. *J. Food Sci.* 59:1013–1017.
- An, H., Peters, M.Y., Seymours, T.A. (1996). Roles of indigenous enzymes on surimi gelation. *Trends in Food Sci. Technol.* 7: 321-327.
- Anonim. (1993). Waters ACCQ Tag Chemistry Package, Manual Number WAT 052874. Waters Millipore Corporation. USA: Maple street Milford, MA 01757.
- Anonim. (2015). Peraturan Menteri Kelautan dan Perikanan Indonesia No. 17/PERMEN-KP/2015. Kementerian Kelautan dan Perikanan. Available at: http://infohukum.kkp.go.id/index.php/hukum/c/20?type_id=1&produk_id=7. (14 Januari 2016).
- Anonim. (2017a). Laporan kinerja Kementerian Kelautan dan Perikanan 2016. Kementerian Kelautan dan Perikanan. Jakarta.
- Anonim. (2017b). Analisis industri surimi Indonesia. Kementerian Kelautan dan Perikanan. Available at: <http://kkp.go.id/artikel/1103-analisis-industri-surimi-indonesia> (14 Agustus 2018).

- Anonim. (2018a). Statistik Perikanan Budidaya. Ditjen Perikanan Budidaya. Kementerian Kelautan dan Perikanan. Jakarta.
- Anonim. (2018b). Frozen minced fish paste (surimi) quality standard. Zhonggang (Fujian) Aquatic Food Co., Ltd. China. Available at: <http://zgseafood.foodmate.com/> (19 Agustus 2018).
- Anonim. (2018c). Spesifikasi surimi. PT. Indo Seafood, Indonesia. Available at: <https://www.indosurimi.com/surimi.html> (19 Agustus 2018).
- [AOAC] Analysis of the Association of Official Analytical Chemists. (1995). *Official Methods of Analysis of the Association of Official Analytical Chemists*. Edisi ke-16. Washington, D.C.
- Apriyantono, A., Fardiaz, D., Puspitasari, N.L., Sedarnawati, Budiyo, S. (1989). Analisis Pangan. Institut Pertanian Bogor: Pusat Antar Universitas Pangan dan Gizi. pp. 30-32.
- Araki, H., Seki, N. (1993). Comparison of reactivity of transglutaminase to various fish actomyosins. *Nippon Suisan Gakkaishi* 59:711-716.
- Ashie, I.N.A., Lanier, T.C. (2000). Transglutaminase in seafood processing. In N. F. Haard, & B. K. Simpson (Eds.), *Seafood enzymes: utilization and influence on postharvest seafood quality* (pp. 147–166). New York: Marcel Dekker.
- Arfat, Y.A., Benjakul, S. (2012). Gelling characteristics of surimi from yellow stripe trevally (*Selaroides leptolepis*). *Int. Aquatic Res.* 4: 1-13.
- Arunde, E., Sinjal, H.J., Monijung, R.D. (2016). Pengaruh penggunaan substrat yang berbeda terhadap daya tetas telur dan sintasan hidup larva ikan lele sangkuriang (*Clarias sp.*). *Budidaya Perairan* 4: 7-15.
- Bachtiar, I., Agustini, T.W., Dwi Anggo, A. (2014). Efektifitas pencucian dan suhu setting (25, 40, 50°C) pada gel kamaboko ikan lele dumbo (*clarias gariepinus*). *Jurnal Pengolahan dan Bioteknologi Hasil Perikanan* 3: 45-50.
- Belitz, H. D., Grosch, W., Schieberle, P. (2004). Fish, wales, crustaceans, mollusks. In: *Food Chemistry*. Berlin, Heidelberg: Springer, pp. 619–642.
- Benjakul, S., Seymour, T.A., Morrissey, M.T., Haejung, A.N. (1996). Proteinase in pacific whiting surimi wash water: identification and characterization. *J. Food Sci.* 61 (6): 1165-1170.
- Benjakul, S., Visessanguan, W., Ishizaki, S., Tanaka, M. (2001). Differences in Gelation Characteristics of Natural Actomyosin from Two Species of Bigeye

- Snapper, *Priacanthus tayenus* and *Priacanthus macracanthus*. *J. Food Sci.* 66: 1311-1318.
- Benjakul, S., Visessanguan, W., Leelapongwattana, K. (2002). Characteristics of muscle from two species of Bigeye snapper, *prlacanthus tayenus* and *Prlacanthus macracanthus*. *J. Food Biochem.* 26: 307-326.
- Benjakul, S., Chantarasuwan, C., Visessanguan, W. (2003). Effect of medium temperature setting on gelling characteristics of surimi from some tropical fish. *Food Chem.* 82: 567-574.
- Benjakul, S., Visessanguan, W. (2003). Transglutaminase-mediated setting in bigeye snapper Surimi. *Food Res.Int.* 36: 253–266.
- Benjakul, S., Visessanguan, W., Chantarasuwan, C. (2004). Effect of high-temperature setting on gelling characteristic of surimi from some tropical fish. *Int.J. Food Sci.Technol.* 39(6): 671-680.
- Benjakul, S., Visessanguan, W., Thongkaew, C., Tanaka, M. (2005). Effect of frozen storage on chemical and gel-forming properties of fish commonly used for surimi production in Thailand. *Food Hydrocol.* 19: 197–207.
- Bertak, J.A., Kahardian, C. (1995). Surimi-based imitation crab characteristic affected by heating method and end point temperature. *J. Food Sci.* 60 (2): 292-296.
- Binsi, P.K., Shamasundar, B.A. (2012). Purification and characterisation of transglutaminase from four fish species: Effect of added transglutaminase on the viscoelastic behaviour of fish mince. *Food Chem.* 132: 1922–1929.
- Blanco, M., Sotelo, C.G., Chapela, M.J., P´erez-Mart´ın, R.I. (2006). Towards sustainable and efficient use of fishery resources: present and future trends. *Trends Food Sci. Technol.* 18(1):29–36.
- Botta, J.R., (1995). *Evaluation of seafood freshness quality*. New York: VCH Publishers. p. 93.
- Boyer, C., Joandel, S., Roussilhes, V., Culioli, J., Ouali, A. (1996). Heat-induced gelation of myofibrillar proteins and myosin from fast- and slow-twitch rabbit muscles. *J. Food Sci.* 61:138–43.
- Camou, J.P, Sebranek, J.G., Olson, D.G. (1989). Effect of heating rate and protein concentration on gel strength and water loss of muscle protein gels. *J. Food Sci.* 54: 850–4.

- Cao, M. J., Hara, K., Osatomi, K., Tachibana, K., Izumi, T., Ishihara, T. (1999). Myofibril-bound serine proteinase (MBP) and its degradation of myofibrillar proteins. *J. Food Sci.* 64: 644–647.
- Cao, M.J., Osatomi, K., Hara, K., Ishihara, T. (2000). Identification of myofibril-bound serine proteinase (MBSP) in the skeletal muscle of lizard fish *Saurida wanieso* which specifically cleaves the arginine site. *Comp. Biochem. Physiol.* Part B 125:255–264.
- Cao, M.J., Jiang, X.J., Zhong, H.C., Zhang, Z.J., Su, W.J. (2006). Degradation of myofibrillar proteins by a myofibril-bound serine proteinase in the skeletal muscle of crucian carp (*Carasius auratus*). *Food Chem.* 94:7–13.
- Carvajal, P.A., Lanier, T.C., MacDonald, G.A. (2005). Stabilization of proteins in surimi. In: Park, J.W., editor. *Surimi and surimi seafood*. 2nd ed. Boca Raton: Taylor & Francis Group. p 163–225.
- Chang-Lee, M.V., Lampila, L.E., Crawford, D.L. (1990). Yield and Composition of Surimi from Pacific Whiting (*Merluccius productus*) and the Effect of Various Protein Additives on Gel Strength. *J. Food Sci.* 55: 83-86.
- Chaijan, M., Benjakul, S., Visessanguan, W., Faustman, C. (2004). Characteristics and gel properties of muscles from sardine (*Sardinella gibbosa*) and mackerel (*Rastrelliger kanagurta*) caught in Thailand. *Food Res. Int.* 37: 1021–1030.
- Chaijan, M., Benjakul, S., Visessanguan, W., Lee, S., Faustman, C. (2007) The effect of freezing and aldehydes on the interaction between fish myoglobin and myofibrillar proteins. *J Agric. Food Chem.* 55:4562–4568
- Chaijan, M., W. Panpipat and S. Benjakul. (2010). Physicochemical properties and gel-forming ability of surimi from three species of mackerel caught in Southern Thailand. *Food Chem.* 121: 85–92.
- Chaijan, M., Klomklao, S., Benjakul, S. (2013). Characterisation of muscles from Frigate mackerel (*Auxis thazard*) and catfish (*Clarias macrocephalus*). *Food Chem.* 139: 414–419.
- Chan, J.K., Gill, T.A., Paulson, A.T. (1992). Cross-linking of myosin heavy chains from cod, herring and silver hake during thermal setting. *J. Food Sci.* 57: 906–912.
- Chan, J.K., Gill, T.A., Paulson, A.T. (1993). Thermal aggregation of myosin subfragments from cod and herring. *J. Food Sci.* 58:1057-1061.
- Chen, J.S., Mehta, K. (1999). Tissue transglutaminase: an enzyme with a split personality. *Int. J. Biochem. Cell Biol.* 31: 817-836.

- Chen, H. H. (2002). Decoloration and gel-forming ability of horse mackerel mince by air-flotation washing. *J. Food Sci.* 67: 2970–2975.
- Chanarat, S., Benjakul, S., H-Kittikun, A. (2012). Comparative study on protein cross-linking and gel enhancing effect of microbial transglutaminase on surimi from different fish. *J. Sci. Food Agric.* 92(4): 844-852.
- Choi, Y.J., Lanier, T.C., Lee, H.G., Cho, Y.J. (1999). Purification and characterization of alkaline proteinase from Atlantic menhaden muscle. *J. Food Sci.* 64:768–771.
- Chomnawang, C., Nantachai, K., Yongsawatdigul, J., Thawornchinsombut, S., Tungkawachara, S. (2007). Chemical and biochemical changes of hybrid catfish fillet stored at 4 °C and its gel properties. *Food Chem.* 103: 420–427.
- CODEX COMMITTEE ON FISH AND FISHERY PRODUCTS. (2015). Appendices for Optional Final Product Requirements for Commodities in the *Code of Practice for Fish and Fishery Products* (CAC/RCP 52-2003). Alesund, Norway.
- Cortez-Vega, W. R., Fonseca, G.G., Prentice, C. (2012). Comparisons of the properties of whitemouth croaker (*Micropogonias furnieri*) surimi and mechanically deboned chicken meat surimi-like material. *Food Nutr. Sci.* 3: 1480–1483.
- Das, N., Khuntia, B.K., Raychaudhuri, U., Dora, K.C., Ganguly, S. (2015). Effect of water washing on the functional properties of fish meat. *Int. J. Med. Microbiol Med. Diseases* 1: 8-12.
- Debusca, A., Tahergorabi, R., Beamer, S.K., Partington, S., Jaczynski, J. (2013). Interactions of dietary fibre and omega-3 rich oil with protein in surimi gels developed with salt substitute. *Food Chem.* 141: 201-208.
- Dey, S.S., Dora, K.C. (2011). Suitability of Chitosan as Cryoprotectant on Croaker Fish (*Johnius gangeticus*) Surimi during Frozen Storage. *J. Food Sci. Technol.* 48(6): 699-705.
- Ding, Y., Liu, R., Rong, J., Xiong, S. (2014). Heat-induced denaturation and aggregation of actomyosin and myosin from yellowcheek carp during setting. *Food Chem.* 149: 237–243.
- ElShehawey, S.M., Gab-Alla, A.A., Mutwally, H.M.A. (2016). Amino Acids Pattern and Fatty Acids Composition of the Most Important Fish Species of Saudi Arabia. *Int. J. Food Sci. Nutr. Eng.* 6: 32-41.

- Folin, O., Ciocalteu, V. (1929). On tyrosine and thryptophane determinations in proteins. *J. Biol. Chem.* 73: 627.
- Folk, J.E., Cole, P.W. (1966). Mechanism of action of guinea pig liver transglutaminase. Purification and properties of the enzyme: Identification of a functional cysteine essential for activity. *J. Biol. Chem.* 241: 5518–5525.
- Folk, J.E. (1980). Transglutaminases. *Ann. Rev. Biochem.* 49: 517–531.
- Folk, J.E. (1983). Mechanism and basis for specificity of transglutaminase-catalyzed e-(c-glutamyl) lysine bond formation. *Adv. Enzymol.* 54: 1–56.
- Foegeding, E.A., Lanier, T.C., Hultin, H.O. (1996). Characteristics of edible muscle tissues. In O.R. Fennema (Ed.), *Food Chemistry. Third edition* (pp. 879-942). New York, USA: Marcel Dekker, Inc.
- Funatsu, Y., Kato, N., Arai, K. (1993). Gel Forming Ability and Cross-linking Ability of Myosin Heavy Chain of Salt-ground Meat from Sardine Surimi Acidified by Lactic Acid. *Nippon Suisan Gakkaishi* 59: 1093-1098.
- Ganesh, A., Dileep, A.O., Shamasundar, B.A., Singh, U. (2006). Gel-forming ability of common carp fish (*cyprinus Carpio*) meat: effect of freezing and frozen storage. *J. Food Biochem.* 30: 342–361.
- Gill, T.A., Conway, J.T. (1989). Thermal aggregation of cod (*Gadus morhua*) muscle proteins using l-ethyl-3-3-dimethylaminopropyl carbodiimide as a zero length crosslinker. *Agric. Biol. Chem.* 53:2553–62.
- Guenneugues, P., Morrissey, M.T. (2005). Surimi resources. In: Park JW, editor. *Surimi and surimi seafood*. 2nd ed. Boca Raton, Fla.: Taylor & Francis Group. p 3–32.
- Guenneugues, P., Lanelli, J. (2014). Surimi Resources and Market. In: Park JW, editor. *Surimi and surimi seafood*. 3th ed. Boca Raton, Fla.: Taylor & Francis Group. p 46–48.
- Haard, N. F., Simpson, B. K., Pan, B. S. (1994). Sarcoplasmic proteins and other nitrogenous compounds. In Z. E. Sikorski, B. S. Pan, and F. Shahidi (Eds.), *Seafood proteins*, New York, USA: Chapman and Hall. pp. 13–39.
- Hall GM, Ahmad NH. (1992). Surimi and fish minced products. *Dalam*: Hall GM (eds). *Fish Processing Technology*. New York: Blackie Academic and Professional.

- Harn-Gam, L., Yee-Leow, C., Baie, S. (2005). Amino acid composition of snakehead fish (*channa striatus*) of various sizes obtained at different times of the year. *Malaysian J. Pharmac. Sci.* 3: 19-30.
- Hashimoto, K., Watabe, S., Kono, M., Shiro, K. (1979). Muscle protein composition of sardine and mackerel. *Bull. Japanese Society Sci. Fish.* 45: 1435-1441.
- Hanan, T., Shaklai, N. (1995). Peroxidative interaction of myoglobin and myoglobin and myosin. *European J. Biochem.* 233: 930-936.
- Haejung An, Peters, M.Y., Seymour, T.A. 1996. Roles of indigenous enzymes in surimi gelation. *Trends Food Sci. Technol.* 7: 321-327.
- Hassan, M.A., Balange, A.K., Senapati, S.R., Martin Xavier, K.A. (2017). Effect of Different Washing Cycles on the Quality of *Pangasius hypophthalmus* Surimi. *Fish. Tech.* 54: 51-59.
- Hemung, B. O., Yongsawatdigul, J. (2005). Ca^{2+} Affects physicochemical and conformational changes of threadfin bream myosin and actin in a setting model. *J. Food Sci.* 70: 455-460.
- Hemung, B. O., Chin, K.B. (2013). Effects of fish sarcoplasmic proteins on the properties of myofibrillar protein gels mediated by microbial transglutaminase. *LWT-Food Sci. Technol.* 53: 184-190.
- Hermansson, A.M. (1979). Aggregation and denaturation involved in gel formation. In: Pour-El A. editor. *Functionality and protein structure*. Washington, D.C.: American Chemical Society. p 81-103.
- Hossain, M.I., Kamal, M.M., Shikha, F.H., Shahidul Haque, M.D. (2004). Effect of Washing and Salt Concentration on the Gel Forming Ability of Two Tropical Fish Species. *Int. J. Agri. Biol.* 6: 762-766.
- Hosseini-Shekarabi, S.P., Hosseini, S.E., Soltani, M., Kamali, A., Valinassab, T. (2014). A Comparative Study on Physicochemical and Sensory Characteristics of Minced Fish and Surimi from Black Mouth Croaker (*Ambloplites niloticus*). *J. Agr. Sci. Tech.* 16: 1289-1300.
- Hosseini-Shekarabi, S.P., Hosseini, S.E., Soltani, M., Kamali, A., Valinassab, T. (2015). Effect of heat treatment on the properties of surimi gel from black mouth croaker (*Ambloplites niloticus*). *Int. Food Res. J.* 22(1): 363-371.
- Hudson B.J.F. 1992. *Biochemistry of Food Proteins*. London: Elsevier Applied Sci.

- Hultin, H.O., Kelleher, S.D. (2000) Surimi processing from dark muscle fish. In: Park JW (ed) Surimi and surimi seafood. Marcel Dekker, New York, pp 59–77
- Hultin, H.O., Kristinsson, H.G., Lanier, T.C., Park, J.W. (2005). Process for recovery of functional proteins by pH shifts. In: Park JW, editor. *Surimi and surimi seafood*. 2nd ed. Boca Raton, Fla.: Taylor & Francis Group. p 107–39.
- Hu, Y., Morioka, K., Itoh, Y. (2007). Hydrolysis of surimi paste from walleye pollock (*Theragra chalcogramma*) by cysteine proteinase cathepsin L and effect of the proteinase inhibitor (E-64) on gelation. *Food Chem.* 104:702–708.
- Hu, Y., Morioka, K., Itoh, Y. (2008). Non-binding property of cathepsin L to myosin. *Food Chem.* 106:741–744.
- Hu, Y., Ji, R., Jiang, H., Zhang, J., Chen, J., Ye, X. (2012). Participation of cathepsin L in “modori” phenomenon in carp (*Cyprinus carpio*) surimi gel. *Food Chem.* 134:2014–2020.
- Ibhadon, S., Abdulsalami, M.S., Emere, M.C., Yilwa, V. (2015). Comparative Study of Proximate, Fatty and Amino Acids Composition of Wild and Farm-Raised African Catfish *Clarias gariepinus* in Kaduna, Nigeria. *Pakistan J. Nutr.* 14 (1): 56-61.
- Jafarpour, A., Gorczyca, E.M. (2009). Characteristics of sarcoplasmic proteins and their interaction with surimi and kamaboko gel. *J. Food Sci.* 74: N16–N22.
- James, C., Purnell, G., James, S.J. (2015). A Review of Novel and Innovative Food Freezing Technologies. *Food Bioprocess Technol.* 8:1616–1634.
- Jiang, S.T., Lee, B.L., Tsao, C.Y., Lee, J.J. (1997). Mackerel cathepsin B and L effects on thermal degradation of surimi. *J. Food Sci.* 62:310–315.
- Jiang, S.T. (2000). Enzymes and their effects on seafood texture. In: *Seafood Enzymes: Utilization and Influence on Postharvest Seafood Quality* (edited by N.F. Haard & B.K. Simpson). Pp. 411–450. New York: Marcel Dekker, Inc.
- Jia, D., Huang, Q., Xiong, S. (2015). Chemical interactions and gel properties of black carp actomyosin affected by MTGase and their relationships. *Food Chem.* Accepted Manuscript.
- Jianrong, W., Ikuo, K., Mikio, S., Nobuo, S. (1994). Effect of calcium ion concentration on the gelling properties and transglutaminase activity of walleye pollock surimi paste. *Nippon Suisan Gakkaishi* 60: 107-113.

- Julavittayanakul, O., Benjakul, S., Visessanguan, W. (2005). Effects of phosphate compounds on gel-forming ability of surimi from bigeye snapper (*Priacanthus tayenus*). *J. Food Hydrocol.* 20: 1153-1163.
- Kaba, N. (2006). The Determination of Technology and Storage Period of Surimi Production from Anchovy (*Engraulis encrasicolus* L., 1758). *Turk. J. Fisher. Aqua. Sci.* 6: 29-35.
- Karthikeyan, M., Shamasundar, B.A., Mathew, S., Kumar, P.R., Prakash, V. (2004). Physicochemical and functional properties of protein from pelagic fatty fish (*Sardinella longiceps*) as function of water washing. *Int. J. Food Prop.* 7 (3): 353-365.
- Karthikeyan, M., Dileep, A.O., Shamasundar, B.A. (2006). Effect of water washing on the functional and rheological properties of proteins from threadfin bream (*Nemipterus japonicus*) meat. *Int. J. Food Sci. Technol.* 41: 1002–1010.
- Kamal, M., Hossain, M.I., Sakib, M.N., Shikha, F.H., Neazuddin, M., Bapary, M.A.J., Islam, M.N. (2005). Effect of salt concentration and cryoprotectants on gel-forming ability of surimi prepared from queen fish (*Chorinemus lysan*) during frozen storage. *Pakistan J. Biologic. Sci.* 8:793-797.
- Kamath, G.G., Lanier, T.C., Foegeding, E.A., Hamann, D.D. (1992). Nondisulfide covalent cross-linking of myosin heavy chain in “setting” of alaska pollock and atlantic croaker surimi. *J. Food Biochem.* 16: 151-172.
- Kang, I.S., Wang, J.J., Shih, J.C.H., Lanier, T.C. (2004). Extracellular production of a functional soy cystatin by *Bacillus subtilis*. *J. Agric. Food Chem.* 52:5052–5056.
- Kang, E. J., Hunt, A.L., Park, J.W. (2008). Effects of salinity on physicochemical properties of alaska pollock surimi after repeated freeze-thaw cycles. *J. Food Sci.* 73: 347–355.
- Keillor, J.W., Clouthier, C.M., Apperley, K.Y.P., Akbar, A., Mulani, A. (2014). Acyl transfer mechanisms of tissue transglutaminase. *Bioorg. Chem.* 57: 186–197.
- Keillor, J.W., Apperley, K.Y.P., Akbar, A. (2015). Inhibitors of tissue transglutaminase. *Trends in Pharmacol. Sci.* 36: 32-40.
- Kinoshita, M., Toyohara, H., Shimizu, Y. (1990). Purification and properties of a novel latent proteinase showing myosin heavy chain-degrading activity from threadfin-bream muscle. *J. Biochem.* 107:587–591.

- Kishi, H., Nozawa, H., Seki, N. (1991). Reactivity of muscle transglutaminase on carp myofibrils and myosin B. *Nippon Suisan Gakkaishi*, 57: 1203–1210.
- Kimura, I., Sugimoto, M., Toyoda, K., Seki, N., Arai, K., Fujita, T. (1991). A study on the cross-linking reaction of myosin in kamaboko “suwari” gels. *Nippon Suisan Gakkaishi*. 57: 1389-1396.
- Kim, S.K. (2013). Marine Proteins and Peptides Biological Activities and Applications. USA: A John Wiley & Sons, Ltd., Publication. Pp. 5-8.
- Ko, W.C., Hwang, M.S. (1995). Contribution of milkfish sarcoplasmic protein to the thermal gelation of myofibrillar protein. *Fish. Sci.* 61:75–78.
- Klesk, K., Yonsawatdigul, J., Park, J.W., Viratchakul, S., Virulhakul, P. (2000). Gel Forming Ability of Tropical Tilapia Surimi as Compared with Alaska Pollock and Pacific Whiting Surimi. *J. Aquat. Food Prod. Technol.* 9: 91-104.
- Kristinsson, H.G., Hultin, H.O. (2003). Effect of low and high pH treatment on the functional properties of codmuscle proteins. *J. Agric. Food Chem.* 51:5103–10.
- Kumazawa, Y., Numazawa, T., Seguro, K., Motoki, M. (1995). Suppression of surimi gel setting by transglutaminase inhibitors. *J. Food Sci.* 60: 715-717.
- Kumazawa, Y., Nakanishi, K., Yasueda, H., Motoki, M. (1996). Purification and Characterization of Transglutaminase from Walleye Pollack Liver. *Fish. Sci.* 62: 959-964.
- Kumazawa, Y., Sano, Y., Seguro, K., Yasueda, H., Nio, N. (1997). Purification and characterization of transglutaminase from Japanese oyster (*Crassostrea gigas*). *J. Agric. Food Chem.* 45: 604–610.
- Ladrat, C., Verrez-Bagnis, V., Noel, J., Fleurence, J. (2003). *in vitro* proteolysis of myofibrillar and sarcoplasmic proteins of white muscle of sea bass (*Dicentrarchus labrax* L.): Effects of cathepsins B, D and L. *Food Chem.* 81:517–25.
- Lan, Y.H., Novakofski, J., McCusker, R.H., Brewer, M.S., Carr, T.R., McKeith, F.K. (1995). Thermal Gelation of Pork, Beef, Fish, Chicken and Turkey Muscles as Affected by Heating Rate and pH. *J. Food Sci.* 60: 936-940.
- Laemmli, U. K. (1970). Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* 227: 680-685.
- Lanier, T. C. (2000). Surimi gelation chemistry. In: Park, J. W. (Eds). Surimi and surimi seafood, p. 237-265. Marcel Dekker: New York.

- Lanier, T.C. (1992). Measurement of Surimi Composition and Functional Properties. In T. C. Lanier & C. M. Lee (Eds.), *Surimi technology* (pp. 123–154). New York: Marcel Dekker.
- Lanier, T.C., Lee, C.M. (1992). *Surimi Technology*. Marcel Dekker, New York. pp 167-179.
- Lanier, T.C., Yongsawatdigul, J., Carvajal-Rondanelli, P. (2014). Surimi Gelation Chemistry. In: Park JW, editor. *Surimi and surimi seafood*. 2nd ed. Boca Raton, Fla.: Taylor & Francis Group. p 101–39.
- Lee, C.M. (1984). Surimi process technology. *Food Technol.* 38:69–80.
- Lee, C.M. (1986). Surimi manufacturing and fabrication of surimi-based products. *Food Tech.* 40(3):115–124.
- Lee, J.J., Chen, H.C., Jiang, S.T. (1993). Purification and characterization of proteinases identified as cathepsin L and L-like (58kDa) proteinase from mackerel (*Scomber australasicus*). *Biosci. Biotech. Biochem.* 57(9): 1470–1476.
- Lee, H.G., Lanier, T.C., Hamann, D.D., Knopp, J.A. (1997). Transglutaminase Effects on Low Temperature Gelation of Fish Protein Sols. *J. Food Sci.* 62: 20–24.
- Lee, S.M., Chiang, C.F., Pan, B.S. (1998). Occurrence of transglutaminase in grey mullet (*Mugil cephalus*) muscle and it's effect on minced fish product. *J. Food Biochem.* 22: 475-487.
- Lee, N., Park, J.W. (1998). Calcium compounds to improve gel functionality of Pacific whiting and Alaska pollock surimi. *J. Food Sci.* 63: 969–974.
- Lee, H.G., Lanier, T.C., Hamann, D.D., Knopp, J.A. (2006). Transglutaminase effects on low temperature gelation of fish protein sols. *J. Food Sci.* 62: 20–24.
- Lertwittayanon, K., Benjakul, S., Maqsood, S., Encarnacion, A.B. (2013). Effect of different salts on dewatering and properties of yellowtail barracuda surimi. *Int. Aquatic Res.* 5:10.
- Li, S., Zhang, N., Liu, H., Ma, C. (2004). Preliminary study of the relationship between autolysis of silver carp (*Hypophthalmichthys molitrix*) myofibrillar proteins and indigenous cathepsin B, L and H. *J. Agric. China Agric. Univ.* 9:71–75.

- Lin, T.M., Park, J.W. (1996). Extraction of proteins from pacific whiting mince at various washing conditions. *J. Food Sci.* 61: 432-438.
- Lowry, O.H., Rosebrough, N.J., Farr, A.L., Randal, R.J. (1951). Protein measurement with the folin phenol reagent. *J. Biol. Chem.*, 193: 265–275.
- Luo, Y.K., Kuwahara, R., Kaneniwa, M., Murata, Y., Yokoyama, M. (2001). Comparison of Gel Properties of Surimi from Alaska Pollock and Three Freshwater Fish Species: Effects of Thermal Processing and Protein Concentration. *J. Food Sci.* 66: 548-554.
- Luo, Y., Kuwahara, R., Kaneniwa, M., Murata, Y., Yokoyama, M. (2004). Effect of soy protein isolate on gel properties of alaska pollock and common carp surimi at different setting conditions. *J. Sci. Food Agric.* 84: 663–671.
- Luo, Y., Shen, H., pan, d., Bu, G. (2008). Gel properties of surimi from silver carp (*Hypophthalmichthys molitrix*) as affected by heat treatment and soy protein isolate. *Food Hydrocol.* 22; 1513–1519.
- Makinodan, Y., Toyohara, H., Ikeda, S. (1984). Comparison of Muscle Proteinase Activity Among Fish Species. *Comp. Biochem. Physiol.* 798: 129-134.
- Makinodan, Y., Toyohara, H., Niwa, E. (1985). Implication of Muscle Alkaline Proteinase in the Textural Degradation of Fish Meat Gel. *Food Sci.* 50: 1351-1355.
- Mahawanich, T., Lekhavichitr, J., Duangmal, K. (2010). Gel properties of red tilapia surimi: effects of setting condition, fish freshness and frozen storage. *Int. J. Food Sci. Technol.* 45: 1777–1786.
- Mahyuddin, K. 2008. *Panduan Lengkap Agribisnis Lele*. Jakarta: Penebar Swadaya.
- Martin-Sanchez, A.M., Navarro, C., Perez-Alvarez, J.A., Kuri, V. (2009). Alternatives for Efficient and Sustainable Production of Surimi: A Review. *Comprehens. Rev. Food Sci. Food Saf.* 8: 359-374.
- Maruyama, N, Nozawa, H, Kimura, I, Satake, M., Seki N. (1995). Transglutaminase-induced polymerization of a mixture of different fish myosins. *Fish. Sci.* 61:495–500.
- Mazrouh, M.M. (2015). Effects of freezing storage on the biochemical composition in muscles of *Saurida undosquamis* (Richardson, 1848) comparing with imported frozen. *Int. J. Fish. Aquatic Studies* 3(2): 295-299.

- Montejano, J.G., Hamann, D.D., Lanier, T.C. (1984). Thermally induced gelation of selected comminuted muscle systems: rheological changes during processing, final strengths, and microstructure. *J. Food Sci.* 49:1496–505.
- Morioka, K., Shimizu, Y. (1990). Contribution of sarcoplasmic proteins to gel forming of fish meat. *Nippon Suisan Gakkaishi* 56:929–33.
- Morioka, K., Shimizu, Y. (1992). Heat-coagulation property of fish sarcoplasmic proteins. *Nippon Suisan Gakkaishi* 58(8):1529–33.
- Morioka, K., Shimizu, Y. (1993). Relationship between the heat-gelling property and composition of fish sarcoplasmic proteins. *Nippon Suisan Gakkaishi* 59(9): 1631 p.
- Morioka, K., Nishimura, T., Obatake, A., Shimizu, Y. (1997). Relationship between the myofibrillar protein gel strengthening effect and the composition of sarcoplasmic proteins from Pacific mackerel. *Fish. Sci.* 63(1):111–114.
- Morrissey, M.T., Wu, J.W., Lin, D., An, H. (1993). Proteinase inhibitor effects on torsion measurements and autolysis of Pacific whiting surimi. *J. Food Sci.* 58:1050-1054.
- Morrissey, M.T., Hartley, P.S., An, H. (1995). Proteolytic activity in Pacific Whiting and effects of surimi processing. *J. Aquat. Food Prod. Technol.* 4:5-18.
- Morrissey, M. T., Tan, S. (2000). World resources for surimi. In J. W. Park (Ed.), *Surimi and surimi seafood* (pp. 1–21). New York, USA: Marcel Dekker.
- Morales, O.G., Ramirez, J.A., Vivanco, D.I., Vazquez, M. (2001). Surimi of fish species from the Gulf of Mexico: evaluation of the setting phenomenon. *Food Chem.* 75: 43-48.
- Muhammad, W.N., Andriyanto, S. (2013). Manajemen budidaya ikan lele dumbo (*Clarias gariepinus*) di Kampung Lele, Kabupaten Boyolali, Jawa Tengah. *Media Akuakultur* 8: 63-71.
- Murthy, L.N., Panda, S.K., Shamasundar, B.A. (2011). Physico-chemical and functional properties of proteins of tilapia (*Oreochromis mossambicus*). *J. Food Proc. Eng.* 34: 83-107.
- Nakagawa, T., Nagayama, F., Ozaki, H., Watabe, S., Hashimoto, K. (1989). Effect of glycolytic enzymes on the gel-forming ability of fish muscle. *Nippon Suisan Gakkaishi* 55: 1045–1050.

- Nakahara, C., Nozawa, H., Seki, N. (1999). A comparison of cross-linking of fish myofibrillar proteins by indigenous and microbial transglutaminases. *Fish. Sci.* 65: 138–144.
- Nakamura, S., Ogawa, M., Saito, M., Nakai, S. (1998). Application of polymannosylated cystatin to surimi from roe-herring to prevent gel weakening. *FEBS Lett* 427:252–54.
- Ni, S., Nozawa, H., Seki, N. (2001). Effect of pH on the gelation of walleye pollack surimi and carp actomyosin pastes. *Fish. Sci.* 67: 920–927.
- Niwa, E. (1992). Chemistry of surimi gelation. In: Lanier, T.C., Lee, C.M., editors. *Surimi technology*. New York: Marcel Dekker, Inc. P 389–438.
- Nolsoe, H., Undeland, I. (2009). The acid and alkaline solubilization process for the isolation of muscle proteins: state of the art. *Food Bioproc. Technol.* 2(1):1–27.
- Nolsoe, H., Marmon, S.K., Undeland, I. (2011). Application of Filtration to Recover Solubilized Proteins During pH-Shift Processing of Blue Whiting (*Micromesistius poutassou*); Effects on Protein Yield and Qualities of Protein Isolates. *Open Food Sci. J.* 5:1-9.
- Nowsad, A.A., Katoh, E., Kanoh, S., Niwa, E. (1995). Effect of sarcoplasmic proteins on the setting of transglutaminase-free paste. *Fish. Sci.* 61(6):1039–40.
- Nowsad, A.A., Katoh, E., Kanoh, A., Niwa, E. (1996). Contribution of transglutaminase on the setting fish pastes at various temperatures. *Fish. Sci.* 62: 94–97.
- Nozawa, H., Mamagoshi, S., Seki, N. (1997). Partial purification and characterization of six transglutaminase from ordinary muscles of various fishes and marine invertebrates. *Comp. Biochem. Physiol.* 118B: 313–317.
- Ochiai, Y., Ochiai, L., Hashimoto, K., Watabe, S. (2001). Quantitative estimation of dark muscle content in the mackerel meat paste and its products using antisera against myosin light chains. *J. Food Sci.* 66: 1301–1305.
- Offer, G. (1987). Myosin filaments. In: *Fibrous Protein Structure*. JM Squire, PJ Vibert, eds. New York: Academic Press. pp 307–56.
- Ofstad, R., Grahl-Madsen, E., Gundersen, B., Lauritzen, K., Solberg, T., Solberg, C. (1993). Stability of cod (*Gadus morhua* L.) surimi processed with CaCl₂ and MgCl₂ added to the wash water. *Int. J. Food Sci. Tech.* 28(5):419–27.

- Ogata, H., Aranishi, F., Hara, K., Osatomi, K., Ishihara, T. (1998). Proteolytic degradation of myofibrillar components by carp cathepsin L. *J. Sci. Food Agric.* 76: 499-504.
- Ohkubo, M., Osatomi, K., Hara, K., Ishihara, T., Aranishi, F. (2005). A novel type of myofibril-bound serine protease from white croaker (*Argyrosomus argentatus*). *Comp Biochem Physiol. Part B* 141:231–236.
- Orak, H.H., Kayisoglu, S. (2008). Quality changes in whole, gutted and filleted three fish species (*Gadus euxinus*, *Mugil cephalus*, *Engraulis encrasicolus*) at frozen storage period (–26°C). *Acta Sci. Pol., Technol. Aliment.* 7(3): 15-28.
- Osatomi, K., Sasai, H., Cao, M., Hara, K., Ishihara, T. (1997). Purification and characterization of myofibril-bound serine proteinase from carp *Cyprinus carpio* ordinary muscle. *Comp Biochem Physiol. Part B* 116:183–190.
- Osibona, A.O., Kusemiju, K.I., Akande, G.R. (2009). Fatty acid composition and amino acid profile of two freshwater species, african catfish (*clarias gariepinus*) and tilapia (*tilapia zillii*). *African J. Food Agric. Nutr. Develop.* 9: 608-621.
- Paker, I. Matak, K.E. (2015). Impact of sarcoplasmic proteins on texture and color of silver carp and alaska pollock protein gels. *LWT-Food Sci. Technol.* 63: 985–991.
- Panpipat, W., Chaijan, M., Benjakul, S. (2010). Gel properties of croaker-mackerel surimi blend. *Food Chem.* 122: 1122–1128.
- Park J.W., Morrissey, M.T. (2000). Manufacturing of surimi from light muscle fish. *Dalam: Park JW (eds). Surimi and Surimi Seafood*. New York: Marcel Dekker, Inc.
- Park, J.W., Lin, T.M. (2005). Surimi: manufacturing and evaluation. *In: Park JW, editor. Surimi and surimi seafood*. 2nd ed. Boca Raton, Fla.: Taylor & Francis Group. p 33–106.
- Park, J.W. (2005). Surimi seafood: Products, Market and Manufacturing. *In: Park JW, editor. Surimi and surimi seafood*. 2nd ed. Boca Raton, Fla.: Taylor & Francis Group. p 375–433.
- Park, D., Choi, S.S., Ha, K.S. (2010). Transglutaminase 2: a multi-functional protein in multiple subcellular compartments. *Amino Acids* 39, 619–631.
- Park, J.W., Nozaki, H., Suzuki, T., Beliveau, J. (2014) Historical Review of Surimi Technology and Market Developments. *In: Park JW, editor. Surimi and surimi seafood*. 3th ed. Boca Raton, Fla.: Taylor & Francis Group. p 4–6.

- Phu, N.V., Morioka, J., Itoh, Y. (2010). Gel-forming characteristics of surimi from white croaker under the inhibition of the polymerization and degradation of protein. *J. Biol. Sci.* 10: 432–439.
- Pipatsattayanuwong, S., Park, J.W., Morrissey, M.T. (1995). Functional properties and shelf life of fresh surimi from pacific whiting. *J. Food Sci.* 60(6): 1241-1244.
- Piyadhamviboon, P., Yongsawatdigul, J. (2009). Proteinase inhibitory activity of sarcoplasmic proteins from threadfin bream (*Nemipterus spp.*). *J. Sci. Food Agric.* 90:291–298.
- Prawira, A. (2008). Pengaruh penambahan tepung alginat (na-alginat) terhadap mutu kamaboko berbahan dasar surimi ikan gabus (*Channa striata*). *Skripsi. Fakultas Perikanan dan Ilmu Kelautan. Bogor: Institut Pertanian Bogor.*
- Raghavan, S., Kristinsson, H.G. (2007). Conformational and rheological changes in catfish myosin as affected by different acids during acid-induced unfolding and refolding. *J. Agric. Food Chem.* 55: 4144–4153.
- Ramirez, J.A., Barrera, M., Morales, O.G., Vazquez, M. (2002). Effect of xanthan and locust bean gums on the gelling properties of myofibrillar protein. *Food Hydrocol.* 16(1):11–6.
- Ramírez, J.A., Velazquez, G., Echevarría, G.L., Torres, J.A. (2007). Effect of adding insoluble solids from surimi wash water on the functional and mechanical properties of pacific whiting grade A surimi. *Bioresour. Technol.* 98: 2148-2153.
- Rawdkuen, S., Sai-Ut, S., Khamsorn, S., Chaijan, M., Benjakul, S. (2009). Biochemical and gelling properties of tilapia surimi and protein recovered using an acid-alkaline process. *Food Chem.* 112: 112-119.
- Reynolds, J., Park, J.W., Choi, Y.J. (2002). Physicochemical properties of pacific whiting surimi as affected by various freezing and storage conditions. *J. Food Sci.* 67(6): 2072-2078.
- Rusherlistyani, Sudaryati, D., Heriningsih, S. (2017). Budidaya lele dengan sistem kolam bioflok. LPPM UPN VY. Yogyakarta.
- Saanin H. 1984. *Taksonomi dan kunci identifikasi ikan*. Jakarta: Bina Cipta.
- Samejima, K., Ishioroshi, M., Yasui, T. (1982). Heat induced gelling properties of actomyosin: effect of tropomyosin and troponin. *Agric. Biol. Chem.* 46:535–40.

- Santana, P., Huda, N., Yang, T.A. (2012). Technology for production of surimi powder and potential of applications. *Int. Food Res. J.* 19(4): 1313-1323.
- Saeki, H., Iseya, Z., Sugiura, S., Seki, N. (1995). Gel forming characteristics of frozen surimi from chum salmon in the presence of protease inhibitors. *J. Food Sci.* 60: 917-921.
- Seki, N., Uno, H., Lee, N.H., Kimura, I., Toyoda, K., Fujita, T., Arai, K. (1990). Transglutaminase activity in Alaska pollack muscle and surimi and its reaction with myosin B. *Nippon Suisan Gakkaishi* 56: 125-132.
- Seymour, T.A., Morrissey, M.T., Peters, M.Y., An, H.J. (1994). Purification and characterization of Pacific whiting proteases. *J. Agric. Food Chem.* 42: 2421–7.
- Sikorski, Z. E. (1994). The myofibrillar proteins in seafoods. In Z. E. Sikorski, B. S. Pan, and F. Shahidi (Eds.), *Seafood proteins*. New York, USA: Chapman & Hall. pp. 40-57.
- Silva, J.L., Chamul, R.S. (2000). Composition of marine and freshwater finfish and shellfish species and their products. In: Martin, R.E., E.P. Carter, G.J. Flick and L.M. Davis, editor. *Marine and Freshwater Products Handbook*. Pennsylvania: Technomic Publishing Company Inc. pp 31–45.
- Shimizu, Y. (1975). Technology of kamaboko. *Science of Cookery* 8: 184.
- Shimizu, Y., Toyohara, H., Lanier, T.C. (1992). Surimi production from fatty and dark-fleshed fish species. In T. C. Lanier & C. M. Lee (Eds.), *Surimi technology* (pp. 181–207). New York: Marcel Dekker.
- Shi, J., Luo, Y., Shen, H., Li, Z. (2014). Gel properties of surimi from silver carp (*Hypophthalmichthys molitrix*): Effects of whey protein concentrate, CaCl_2 , and setting condition. *J. Aquat. Food Prod. Technol.* 23: 489–497.
- Sriket, C. (2014). Proteases in fish and shellfish: Role on muscle softening and prevention. *Int. Food Res. J.* 21: 433-445.
- Stone, A.P., Stanley, D.W. (1992). Mechanisms of fish muscle gelation. *Food Res. Int.* 25: 381-388.
- Stoknes, I., Rustad, R., Mohr, V. (1993). Comparative Studies of the Proteolytic Activity of Tissue Extracts from Cod (*Cadus morhua*) and Herring (*Clupea harengus*). *Comp. Biochem. Physiol.* 106B: 613-619.
- Stoknes, I., Rustad, T. (1995). Proteolytic Activity in Muscle from Atlantic Salmon (*Salmo salar*). *Food Sci.* 60: 71 1-71 4.

- Suzuki T. 1981. *Fish and Krill Protein in Processing Technology*. London: Applied Science Publishing. Ltd.
- Sun, X.D., Holley, R.A. (2011). Factors Influencing Gel Formation by Myofibrillar Proteins in Muscle Foods. *Comprehens. Rev. Food Sci. Food Saf.* 10: 33-51.
- Suvanich, V., Marshall, D.L. (1998). Influence of storage time and temperature on quality of cat fish (*Ictalurus punctatus*) frames. *J. Aquat. Food Prod. Technol.* 7: 61–76.
- Suvanich, V., Jahncke, M.L., Marshall, D.L. (2000). Changes in selected chemical quality characteristics of channel catfish frame mince during chill and frozen storage. *J. Food Sci.* 65:24–29.
- Suyanto, S.R. (2009). *Budidaya Ikan Lele*. Penebar Swadaya. Jakarta.
- Sych, J., Lacroix, C., Adamounou, L.T., Castaigne, F. (1990). Cryoprotective effects of lactitol, palatinit and polydextrose on cod surimi proteins during frozen storage. *J. Food Sci.* 55: 356–360.
- Szczesniak, A.S. (1986). Sensory texture evaluation methodology. *Reciprocal Meat Conference Proceedings* 39: 86-96.
- Takeda, H., Seki, N. (1996). Enzyme-catalyzed cross-linking and degradation of myosin heavy chain in walleye pollack surimi paste during setting. *Fish. Sci.* 62: 462–467.
- Taguchi, T., Ishizaka, H., Tanaka, M., Nagashima, Y., Amano, K. (1987). Protein-protein interaction of fish myosin fragments. *J. Food Sci.* 52: 1103.
- Thrash, B. (1983). The Japanese kamaboko industry. In “Pastry and Peanuts.” Alaska Fisheries Development Foundation, Anchorage, AK.
- Toyoda, K., Kimura, I., Fujita, T., Noguchi, S.F., Lee, C.M. (1992). The surimi manufacturing process. In: Lanier TC, Lee CM, editors. *Surimi technology*. New York: Marcel Dekker Inc. p 79–112.
- Toyohara, H., Kinoshita, M., Shimizu, Y. (1990). Proteolytic degradation of threadfin-bream meat gel. *J Food Sci.* 55:259–260.
- Tsukamasa, Y., Shimizu, Y. (1990). Setting property of sardine and Pacific mackerel meat. *Nippon Suisan Gakkaishi*, 56, 1105-1112.

- Tsukamasa, Y., Shimizu, Y. (1991). Factors affecting the transglutaminase-associated setting phenomenon in fish meat sol. *Nippon Suisan Gakkaishi* 57: 535-540.
- Tsukamasa, Y., Miyake, Y., Ando, M., Makinodan, Y. (2002). Total activity of transglutaminase at various temperatures in several fish meats. *Fish. Sci.* 68: 929-933.
- Vareltsiz, K., Zetou, F., Soultos, N., Tsiaras, I. (1989). Use of Hake (*Merluccius merluccius*) Surimi in a Frankfurter Formulation. *Int. J. Food Sci. Technol.* 45: 277-281.
- Vareltzis, K. (2000). Fish proteins from unexploited and underdeveloped sources. In: Doxastakis, G., Kiosseoglou, V., eds. *Novel Macromolecules in Food Systems*. Amsterdam: Elsevier, pp. 133-159.
- Venugopal, V. (2009). Seafood proteins: functional properties and protein supplements. In: *Marine Products for Healthcare: Functional and Bioactive Nutraceutical Compounds from the Ocean*. Boca Raton: CRC Press, pp. 51-102.
- Vilhelmsson O. (1997). The state of enzyme biotechnology in the fish processing industry. *Trends Food Sci. Technol.* 8(8): 266-70.
- Visessanguan, W., Menino, A.R., Kim, S.M., An, H. (2001). Cathepsin L: A predominant heat-activated proteinase in arrowtooth flounder muscle. *J. Agric. Food Chem.* 49:2633-2640.
- Wan, J., Kimura, I., Satake, M., Seki, N. (1994). Effect of calcium ion concentration on the gelling properties and transglutaminase activity of walleye pollack surimi past. *J. Fish Sci.* 60: 107-114.
- Wan, J., Kimura, I., Satake, M., Seki, N. (1995). Causes of inferior gel forming ability of salmon surimi paste. *Fish. Sci.* 61 : 711-715.
- Wang, X., Hirata, T., Fukuda, Y., Kinoshita, M., Sakaguchi, M. (2002). Acceptability comparison of kamaboko gels derived from silver carp surimi and from walleye pollack surimi between the Chinese and Japanese. *Fish. Sci.* 68: 165-169.
- Wasson DH. 1992. Fish muscle proteases and heat-induced myofibrillar degradation: a review. *J. Aquat. Food Prod. Technol.* 8: 23-41.
- Weng, W.Y., Hamaguchi, P.Y., Osako, K., Tanaka, M. (2007). Effect of indigenous acid proteinases on the properties of edible films prepared from Alaska pollack surimi. *Food Chem.* 105: 996-1002.

- Wijayanti, I., Santoso, J., Jacob, A.M. (2012). Pengaruh frekuensi pencucian terhadap karakteristik gel surimi ikan lele dumbo (*clarias gariepinus*). *J. Saintek Perikanan* 8: 32-37.
- Worratao, A., Yongsawatdigul, J. (2003). Cross-linking of actomyosin by crude tilapia (*oreochromis niloticus*) transglutaminase. *J. Food Biochem.* 27: 35-51.
- Worratao, A., Yongsawatdigul, J. (2005). Purification and characterization of transglutaminase from Tropical tilapia (*Oreochromis niloticus*). *Food Chem.* 93: 651-658.
- Xiong, Y.L., Brekke, C.J. (1991). Protein extractability and thermally induced gelation properties of myofibrils isolated from pre-rigor and post-rigor chicken muscles. *J. Food Sci.* 56:210-5.
- Xiong, Y.L. (1994). Myofibrillar protein from different muscle fiber types: implications of biochemical and functional properties in meat processing. *CRC Crit Rev Food Sci. Nutr.* 34:293-320.
- Yamashita, M., Konagaya, S. (1990). Participation of Cathepsin L into Extensive Softening of the Muscle of Chum Salmon Caught During Spawning Migration. *Nippon Suisan Gakkaishi* 56: 1271-1277.
- Yang, Z., Wang, W., Wang, H., Ye, Q. (2014). Effects of a highly resistant rice starch and pre-incubation temperatures on the physicochemical properties of surimi gel from grass carp (*Ctenopharynxodon Idellus*). *Food Chem.* 145: 212-219.
- Yasueda, H., Kumazawa, H., Motoki, M. (1994). Purification and characterization of a tissue-type transglutaminase from red sea bream (*Pagrus major*). *Biosci. Biotechnol. Biochem.* 58: 2041-2045.
- Yasui, T., Ishioroshi, M., Samejima K. (1980). Heat-induced gelation of myosin in the presence of actin. *J Food Biochem* 4:61-78.
- Yin, T., Park, J.W. (2015). Optimum processing conditions for slowly heated surimi seafood using protease-laden Pacific whiting surimi. *LWT- Food Sci. Technol.* 63: 490-496.
- Yongsawatdigul, J., Park, J.W., Virulhakul, P., Viratchakul, S. (2000). Proteolytic Degradation of Tropical Tilapia Surimi. *J. Food Sci.* 65: 129-133.
- Yongsawatdigul, J., Worratao, A., Park, J.W. (2002). Effect of Indigenous Transglutaminase on Threadfin Bream Surimi Gelation. *J. Food Sci.* 67: 3258-3263.

- Yongsawatdigul, J., Park, J.W. (2004). Gelation of threadfin bream surimi as affected by thermal denaturation, transglutaminase, and proteinase(s) activities. *Dev. Food Sci.* 42:343–356.
- Yongsawatdigul, J., Piyadhamviboon, P., Singchan, K. (2006). Gel-forming ability of small scale mud carp unwashed and washed mince as related to indigenous proteinases and transglutaminase activities. *Eur. Food Res. Technol.* 223(6):769–74.
- Yongsawatdigul, J., Piyadhamviboon, P. (2007). Gel-enhancing effect and protein cross-linking ability of tilapia sarcoplasmic proteins. *J. Sci. Food Agric.* 87:2810–2816.
- Zaitsev, V., Kizeveter, I., Lagunov, L., Makarova, T., Minder, L., Podsevalov. (1969). *Fish Curing and Processing*. Mir Publisher, Moscow.
- Zayas, J.F. (1997). *Functionality of proteins in food*. Springer-Verlag Berlin Heidelberg, New York.
- Zepeda, C.M.G., Kastner, C.L., Kropf, D.H., Hunt, M.C., Kenney, P.B., Schwenke, J.R., Schleusener, D.S. (1993). Utilization of surimi-like products from pork with sex-odor in restructured, precooked pork roast. *J. Food Sci.* 58: 53-58.
- Zhang, L., Q. Li, J. Shi, B. Zhu and Y. Luo. (2018). Changes in chemical interactions and gel properties of heat-induced surimi gels from silver carp (*Hypophthalmichthys molitrix*) fillets during setting and heating: Effects of different washing solutions. *Food Hydrocoll.* 75: 116–124.
- Zhou, A., Benjakul, S., Pan, K., Gong, J., Liu, X. (2006). Cryoprotective effects of trehalose and sodium lactate of tilapia (*Sarotherodon nilotica*) surimi during frozen storage. *J. Food Chem.* 96: 96-103.