

DAFTAR PUSTAKA

- Abbate, A. 2017. Tackling the Problem of Adverse Ventricular Remodeling After Myocardial Infarction in Morrow, D.A. *Myocardial Infarction A Companion to Braunwald's Heart Disease*. Missouri: Elsevier. p:449-461.
- Aniyathodiyil G, Bohra SS, Mottengar A, Govind SC. 2017. Speckle-Tracking echocardiography to assess global and regional left ventricular function in acute myocardial infarction. *J Indian Acad Echocardiogr Cardiovasc Imaging*. 1(3):177-84.
- Antoni, M. L., Mollema, S. A., Atary, J. Z., Borleffs, C. J. W., Boersma, E., Van De Veire, N. R. L., Holman, E. R., Van Der Wall, E. E., Schalij, M. J., Bax, J. J. 2010. Time course of global left ventricular strain after acute myocardial infarction. *European Heart Journal*. 31(16): 2006–2013.
- Bao, Y.S., Na, S.P., Zhang, P., Jia, X.B., Liu, R.C., Yu, C.Y., Mu, S.H., Xie, R.J. 2012. Characterization of Interleukin-33 and Soluble ST2 in Serum and Their Association with Disease Severity in Patients with Chronic Kidney Disease. *J Clin Immunol*. 32:587-594.
- Bahuleyan, C.G., Alummoottil, G.K., Abdullakutty, J., Lordson, A.J., Babu, S., Krishnakumar, V.V., Pillai, A.M., Abraham, G., Dilip, M.N. 2017. Prognostic Value of Soluble ST2 Biomarker in Heart Failure Patients with Reduced Ejection Fraction – A Multicenter Study. *Indian Heart Journal*. 1-6.
- Basu, S., Senior, R., van der Does, R., Bruckner, T., Lahiri, A. 1997. Beneficial Effects of Intravenous and Oral Carvedilol Treatment in Acute Myocardial Infarction. A placebo-controlled, randomized trial. *Circulation*. 96(1):183-191.
- Bergis, D., Kassis, V., Ranglack, A., Koeberle, V., Piiper, A., Kronenberger, B., Zeuzem, S., Waidmann, O., Radeke, H.H. 2013. High Serum Level of the Interleukin-33 Receptor Soluble ST2 as a Negative Prognostic Factor in Hepatocellular Carcinoma. *Transal Oncol*. 6(3):311-318.
- Cartensen, H.G., Larsen, L.H., Hassager, C., Kofoed, K.F., Dalsgaard, M., Kristensen, C.B., Jensen, J.S., Mogelvang, R. 2015. Tissue Velocities and Myocardial Deformation in Asymptomatic and Symptomatic Aortic Stenosis. *Journal of the American Society of Echocardiography*. 1(1):1-12.
- Chistiakov, D.A., Orekhov, A.N., Bobryshev, Y.V. 2016. The Role of Cardiac Fibroblast in Post-Myocardial Heart Tissue Repair. *Experimental and Molecular Pathology*. 101(1): 231-240.
- Choe, J.C., Cha, K.S., Yun, E.Y., Ahn, J., Kim, J.S., Choi, J.H., Park, Y.H., Lee, H.C., Kim, H.J., Chun, K.J., Hong, T.J., Ahn, Y., Jeong, M.H., Chae, S.C., Kim, Y.J. 2018. Reverse Left Ventricular Remodelling in ST-Elevation Myocardial Infarction Patients Undergoing Primary

- Percutaneous Coronary Intervention: Incidence, Predictors and Impact on Outcome. *Heart, Lung and Circulation*. 27:154-164.
- Claus, P., Omar, A.M.S., Pedrizzetti, G., Sengupta, P.P., Nagel, E. 2015. Tissue Tracking Technology for Assessing Cardiac Mechanics: Principles, Normal Values, and Clinical Application. *JACC: Cardiovascular Imaging*. 8(12): 1444-1460.
- Cohn, J.N., Ferrarri, R., Sharpe, N. 2000. Cardiac Remodeling – Concepts and Clinical Implications: A Concensus Paper From an International Forum on Cardiac Remodeling. *Journal of the American College of Cardiology*. 35(3): 569-582.
- Cui, G., Qi, H., Gundersen, M.D., Yang, H., Christiansen, I., Sorbye, S.W., Goll, R., Florholmen, J., 2015. Dynamics of the IL-33/ST2 Network in the Progression of Human Colorectal Adenoma to Sporadic Colorectal Cancer. *Cancer Immunol Immunother*. 64(2):181-190.
- D'Andrea, A., Cocchia, R., Caso, P., Riegler, L., Scarafilo, R., Salerno, G., Golia, E., Di Salvo, G., Calabro, P., Bigazzi, M.C., Liccardo, B., Esposito, N., Cuomo, S., Bossone, E., Russo, M.G., Calabro, R. 2011. Global Longitudinal Speckle-Tracking Strain is Predictive of Left Ventricular Remodeling After Coronary Angioplasty In Patients With Recent Non-ST Elevation Myocardial Infarction. *Int J Cardiol*. 153(2):185-191.
- D'Elia, N., D'hooge, J., Marwick, T.H. 2015. Association Between Myocardial Mechanics and Ischemic LV Remodeling. *JACC : Cardiovascular Imaging*. 8(12): 1430-1443.
- Dhillon, O. S., Narayan, H. K., Khan, S. Q., Kelly, D., Quinn, P. A., Squire, I. B., Davies, J. E., Ng, L. L. 2013. Pre-discharge risk stratification in unselected STEMI: Is there a role for ST2 or its natural ligand IL-33 when compared with contemporary risk markers?. *International Journal of Cardiology*. 167(5):2182–2188.
- Diao, K., Yang, Z., Ma, M., He, Y., Zhao, Q., Liu, X., Gao, Y., Xie, L.J., Guo, Y. 2017. The Diagnostic Value of Global Longitudinal Strain (GLS) on Myocardial Infarction Size by Echocardiography: A Systematic Review and Meta-analysis. *Scientific Reports*, 7(1): 1-8.
- Doughty, R.N., Whalley, G.A., Gamble, G., MacMahon, S., Sharpe, N. 1997. Left Ventricular Remodeling with Carvedilol in Patients with Congestive Heart Failure due to Ischemic Heart Disease. *J Am Coll Cardiol*. 29: 1060-1066.
- Erlebacher, J.A., Weiss, J.L., Weisfeldt, M.L., Bulkley, B.H. 1984. Early Dilation of the Infarcted Segment in Acute Transmural Myocardial Infarction: Role of Infarct Expansion in Acute Left Ventricular Enlargement. *J Am Coll Cardiol*. 4:201-208.
- Falk E, Nakano M, Bentzon JF, Finn AV, Vimani R. 2013. Update on acute coronary syndromes: The pathologists' view. *Eur Heart J*; 34:719-28
- Gerber, Y., Weston, S.A., Enrique-Sarano, M., Manemann, S.M., Chamberlain, A.M., Jiang, R., Roger, V.L. 2016. Atherosclerotic Burden and Heart Failure After Myocardial Infarction. *JAMA Cardiology*. 2(1): 156-162.

- Hartopo, A. B., Setianto, B. Y., Pradana, A. D., Murti, H., Hadwiono, M.R. 2016. Pengaruh Kadar Soluble ST2 Serum Terhadap Kejadian Kardiovaskular Mayor Selama Perawatan Infark Miokard Akut dengan Elevasi Segmen ST. Yogyakarta.
- Hutomo, P., Hartopo, A.B., Sukmasari, I., Puspitawati, I., Gharini, P.P.R., Setianto, B.Y. 2017. Prognostic Factor of Soluble ST2 Serum on 90 Days-Major Cardiovascular Events in ST-Elevation Acute Myocardial Infarction Patients with Reperfusion Therapy. *Acta Cardiologia Indonesiana*. 3(1):1-13.
- Hong, Y.S., Moon, S.J., Joo, Y.B., Jeon, C.H., Cho, M.L., Ju, H.J., Oh, H.J., Heo, Y.J., Park, S.H., Kim, H.Y., Min, J.K. 2011. Measurement of Interleukin-33 (IL-33) and IL-33 Receptors (sST2 and ST2L) in Patients with Rheumatoid Arthritis. *Journal of Korean Medical Science*. 26(9):1132-1139.
- Hur, M., Kim, H., Kim, H.J., Yang, H.S., Magrini, L., Marino, R., Cardelli, P., Di Somma, S. Soluble ST2 Has a Prognostic Role in Patients With Suspected Sepsis. *Annals of Laboratory Medicine*. 35(6):570-577.
- Joyce, E., Hoogslag, G.E., Leong, D.P., Donnaire, P., Katsanos, S., Boden, H., Schali, M.J., Marsan, N.A., Bax, J.J., Delgado, V. 2014. Association Between Left Ventricular Global Longitudinal Strain and Adverse Left Ventricular Dilatation After ST-Segment-Elevation Myocardial Infarction. *Circ Cardiovasc Imaging*. 7(1):74-81.
- Kakkar, R. dan Lee, R. T. 2008. The IL-33/ST2 pathway: therapeutic target and novel biomarker. *Nat Rev Drug Discov*. 7(10):827-840.
- Kober, L., Torp-Pedersen, C., Carlsen J.E., Bagger, H., Eliassen, P., Lyngborg, K., Videbaek, J., Cole, D.S., Auclert, L., Pauly, N.C. 1995. A Clinical Trial of The Angiotensin-Converting-Enzyme Inhibitor Trandolapril in Patients With Left Ventricular Dysfunction After Myocardial Infarction. Trandolapril Cardiac Evaluation (TRACE) Study Group. *N Engl J Med*. 333(25):1670-6.
- Kondo, Y., Yoshimoto, T., Yasuda, K., Futatsugi-Yumikura, S., Morimoto, M., Hayashi, N., Hoshino, T., Fujimoto, J., Nakanishi, K. 2008. Administration of IL-33 Induces Airway Hyperresponsiveness and Goblet Cell Hyperplasia in the Lungs in the Absence of Adaptive Immune System. *Int Immunol*. 20(6):791-800.
- Kuroiwa, K., Arai, T., Okazaki, H., Minota, S., Tominaga, S.I. 2001. Identification of Human ST2 Protein in the Sera of Patients with Autoimmune Diseases. *Biochemical and Biophysical Research Communications*. 284:1104-1108.
- Lang, R.M., Badano, L.P., Mor-Avi, V., Afilalo, J., Armstrong, A., Ernande, L., Flachskampf, F.A., Foster, E., Goldstein, S.A., Kuznetsoc, T., Lancellotti, P., Muraru, D., Picard, M.H., Rietzschel, E.R., Rudski, L., Spencer, K.T., Tsang, W., Voigt, J.U. 2015. Recommendations for Cardiac Chamber Quantification by Echocardiography in Adults: an Update from the American Society of Echocardiography and the

- European Association of Cardiovascular Imaging. *J Am Soc Echocardiogr.* 28(1):1-39.
- Lancellotti, P., Dulgheru, R., Magne, J., Henri, C., Servais, L., Bouznad, N., Ancion, A., Martinez, C., Davin, L., Le Goff, C., Nchimi, A., Pierrard, L., Oury, C. 2015. Elevated Plasma Soluble ST2 is Associated with Heart Failure Symptoms and Outcome in Aortic Stenosis. *PLoS ONE.* 10(9):e0138940.
- Lu, D.P., Zhou, X.Y., Yao, L.T., Liu C.G., Ma, W., Jin, F., Wu, Y.F. 2014. Serum Soluble ST2 is Associated with ER-positive Breast Cancer. *BMC Cancer.* 14:198-205.
- Marino, P., Zanolla, L., Zardini, P. 1988. Effect of Streptokinase on Left Ventricular Remodeling and Function After Myocardial Infarction. *J Am Coll Cardiol.* 11:35-41.
- Mele, D., Nardoza, M., & Chiodi, E. 2017. Early Speckle - tracking Echocardiography Predicts Left Ventricle Remodeling after Acute ST - Segment Elevation Myocardial Infarction. *J Cardiovasc Echogr.* 27(2): 93–98.
- Minana, G., Nunez, J., Bayes-Genis, A., Revuelta-Lopez, E., Rios-Navarro, C., Nunez, E., Chorro, F.J., Lopez-Lereu, M.P., Monmeneu, J.C., Lupon, J., Bodi, V. 2018. ST2 and Left Ventricular Remodeling After ST-Segment Elevation Myocardial Infarction: A Cardiac Magnetic Resonance Study. *International Journal of Cardiology.* 270(6):336-342.
- Mueller, T., Gegenhuber, A., Leitner, I., Poelz, W., Haltmayer, M., Dieplinger, B. 2016. Diagnostic and Prognostic Accuracy of Galectin-3 and Soluble ST2 for Acute Heart Failure. *Clinica Chimica Acta.* 463:158-164.
- O’Gara, P. T., Kushner, F. G., Ascheim, D. D., Casey, D. E., Chung, M. K., Lemos, J. A. De, Ettinger, S. M., Fang, J. C., Fesmire, F. M., Franklin, B. A., Granger, C. B., Krumholz, H. M., Linderbaum, J. A., Morrow, D. A., Newby, L. K., Ornato, J. P., Ou, N., Radford, M. J., Tamis-holland, J. E., Tommaso, C. L., Tracy, C. M., Woo, Y. J., Zhao, D. X., Jacobs, A. K., Halperin, J. L., Albert, N. M., Brindis, R. G., Creager, M. A., Demets, D., Guyton, R. A., Hochman, J. S., Kovacs, R. J., Kushner, F. G., Ohman, E. M., Stevenson, W. G., Yancy, C. W. 2013. 2013 ACCF / AHA Guideline for the Management of ST-Elevation Myocardial Infarction. *J Am Coll Cardiol.* 61(4):e78–e140.
- Oshikawa, K., Kuroiwa, K., Tago, K., Iwahana, H., Yanagisawa, K., Ohno, S., Tominaga, S.I., Sugiyama, Y. 2001. Elevated soluble ST2 protein levels in sera of patients with asthma with an acute exacerbation. *Am J Respir Crit Care Med.* 164:277–281.
- Oshikawa, K., Kuroiwa, K., Tokunaga, T., Hagihara, S.I., Tominaga S.I., Sugiyama, Y., 2001. Acute Eosinophilic Pneumonia with Increased Soluble ST2 in Serum and Bronchoalveolar Lavage Fluid. *Respiratory Medicine.* 95532-533.

- Oshikawa, K., Yanagisawa, K., Ohno, S., Tominaga, S.I., Sugiyama, Y. 2002. Expression of ST2 in Helper T Lymphocytes of Malignant Pleural Effusions. *American Journal of Respiratory and Critical Medicine*. 165(7):1005-1009.
- Park, Y.H., Kang, S.J., Song, J.K., Lee, E.Y., Song, J.M., Kang, D.H., Kim, Y.H., Lee, C.W., Hong, M.K., Kim, J.J., Park, S.W., Park, S.J. 2008. Prognostic Value of Longitudinal Strain After Primary Reperfusion Therapy in Patients With Anterior-Wall Acute Myocardial Infarction. *J Am Soc Echoardiogr*. 21(3):263-267.
- Pascual-Figal, D. A. dan Januzzi, J. L. 2015. The Biology of ST2: The International ST2 Consensus Panel. *The American journal of cardiology*. 115(7 Suppl):3B–7B.
- Pfeffer, M.A., Braunwald, E., Moye, L.A., Brown, E.J. Jr., Cuddy, T.E., Davis, B.R., Geltman, E.M., Goldman, S., Flaker, G.C., Klein, M., Lamas, G.A., Packer, M., Rouleau, J., Rouleau, J.L., Rutherford, J., Wertheimer, J.H., Hawkins, C.M. 1992. Effect of Captopril On Mortality And Morbidity In Patients With Left Ventricular Dysfunction After Myocardial Infarction. *N Engl J Med*. 327(10):669-677.
- Pfeffer, M.A., Lamas, G.A., Vaughan, D.E., Parisi, A.F., Braunwald, E. 1988. Effect of Captopril on Progressive Ventricular Dilatation After Anterior Myocardial Infarction. *N Engl J Med*. 319(2):80-86.
- Pfeffer, M.A., MacMurray, J.J.V., Velazquez E.J., Rouleau J.L., Kober, L., Maggioni, A.P., Solomon, S.D., Swedberg, K., Van de Werf, F., White, H., Leimberger, J.D., Henis, M., Edward, S., Zelenkofske, S., Sellers, M.A., Califf, R.M. 2003. Valsartan, Captopril, or Both in Myocardial Infarction Complicated By Heart Failure, Left Ventricular Dysfunction, or Both. *N Engl J Med*. 349(20):1893-906.
- Prastaro, M., Pirozzi, E., Gaibazzi, N., Paolilo, S., Santoro, C., Savarese, G., Losi, M.A., Esposito, G., Filardi, P.P., Trimarco, B., Galderisi, M. 2017. Expert Review on the Prognostic Role of Echocardiography After Acute Myocardial Infarction. *J Am Soc Echocardiogr*. 30:431-443.
- Richards, A. M., Di Somma, S., Mueller, T. 2015. ST2 in stable and unstable ischemic heart diseases. *The American journal of cardiology*. 115(7 Suppl):48B–58B.
- Sabatine, M. S., Morrow, D. A., Higgins, L. J., MacGillivray, C., Guo, W., Bode, C., Rifai, N., Cannon, C. P., Gerszten, R. E., Lee, R. T. 2008. Complementary Roles for Biomarkers of Biomechanical Strain ST2 and N-Terminal Prohormone B-Type Natriuretic Peptide in Patients with ST-Elevation Myocardial Infarction. *Circulation*. 117(15):1936-1944.
- Sanada, S., Hakuno, D., Higgins, L. J., Schreiter, E. R., Mckenzie, A. N. J., Lee, R. T. 2007. IL-33 and ST2 comprise a critical biomechanically induced and cardioprotective signaling system. *J Clin Invest*. 117(6):1538–1549.
- Sanchez-Mas, J., Lax, A., Asensio-Lopez, M.C., Pascual-Figal, D.A. 2014. Modulation of IL-33/ST2 System in Postinfarction Heart Failure: Correlation with Cardiac Remodelling Markers. *European of Clinical Investigation*. 44(7).

- Sawada, H., Naito, Y., Hirotsu, S., Akahori, H., Toshihiro, I., Okuhara, Y., Miki, K., Eguchi, A., Mitsuno, M., Miyamoto, Y., Ohyanagi, M., Tsujino, T., Masuyama, T. 2013. Expression of Interleukin-33 and ST2 in Nonrheumatic Aortic Valve Stenosis. *International Journal of Cardiology*. 168:529-530.
- Schwammenthal, E., Adler, Y., Amichai, K., Sagie, A., Behar, S., Hod, H., Feinberg, M.S. 2003. Prognostic Value of Global Myocardial Performance Indices in Acute Myocardial Infarction: Comparison to Measures of Systolic and Diastolic Left Ventricular Function. *Chest*. 124:1645-1651.
- Scirica BM, Morrow DA. 2015. ST-Elevation Myocardial Infarction: Pathology, Pathophysiology, and Clinical Features in Mann DL *Braunwald's Heart Disease*. 10th ed. Philadelphia: Elsevier Saunders. p:1088-93.
- Sharpe, N., Murphym J., Smith, H., Hannan, S. 1988. Treatment of Patients with Symptomless Left Ventricular Dysfunction After Myocardial Infarction. *Lancet*. 1(8580):255-259.
- Shih, H., Lee, B., Lee, R.J., Boyle, A.J. 2010. The Aging Heart and Post-Infarction Left Ventricular Remodeling. *J Am Coll Cardiol*. 57(1):9-17.
- Shimpo, M., Morrow, D. A., Weinberg, E. O., Sabatine, M. S., Murphy, S. A., Antman, E. M., Lee, R. T. 2004. Serum Levels of the *Interleukin-1* Receptor Family Member ST2 Predict Mortality and Clinical Outcome in Acute Myocardial Infarction. *Circulation*. 109:2186–2190.
- Sjøli, B., Ørn, S., Grenne, B., Vartdal, T., Smiseth, O. A., Edvardsen, T., Brunvand, H. 2009. Comparison of Left Ventricular Ejection Fraction and Left Ventricular Global Strain as Determinants of Infarct Size in Patients with Acute Myocardial Infarction. *Journal of the American Society of Echocardiography*, 22(11): 1232–1238
- St. John Sutton, M.G. dan Sharpe, N. 2000. Left Ventricular Remodeling After Myocardial Infarction: Pathophysiology and Therapy. *Circulation*. 101(1): 2981-2988.
- Steg, P.G., James, S.K., Badano, L.P., Borger, M.A., Blomstorm-Lundqvist, C., Mario, C. Di, Ducrocq, G., Gershlick, A.H., Atar, D., Dickstein, K., Fernandez-Aviles, F., Giannuzzi, P., Halvorsen, S., Huber, K., Juni, P., Kastrati, A., Knuuti, J., Lenzen, M.J., Mahaffey, K.W., Valgimigli, M., van Hof, A., Widimsky, P., Zahger, D. 2012. ESC Guidelines for the management of acute myocardial infarction in patients presenting with ST-segment elevation The Task Force on the management of ST-segment elevation acute myocardial infarction of the European Society of Cardiology. *European Heart Journal*. 33:2569–2619.
- Stundl, A., Lunstedt, N.S., Courtz, F., Freitag-Wolf, S., Frey, N., Holdenrieder, S., Zur, B., Grube, E., Nickenig, G., Werner, N., Frank, D., Sinning, J.M. 2017. Soluble ST2 for Risk Stratification and the Prediction of Mortality in Patients Undergoing Transcatheter Aortic Valve Implantation. *American Journal of Cardiology*. 120:986-993.

- Tajima, S., Bando M., Ohno, S., Sugiyama, Y., Oshikawa, K., Tominaga, S., Itoh, K., Takada, T., Suzuki, E., Gejyo, F. 2007. ST2 gene induced by type 2 helper T cell (Th2) and proinflammatory cytokine stimuli may modulate lung injury and fibrosis. *Exp Lung Res.* 33:81–97.
- Tajima, S., Oshikawa, K., Tominaga, S., Sugiyama, Y. 2003. The increase in serum soluble ST2 protein upon acute exacerbation of idiopathic pulmonary fibrosis. *Chest.* 124:1206–1214.
- Touchstone, D.A., Beller, G.A., Nygaard, T.W., Tedesco, C., Kaul, S. 1988. Effects of Successful Intravenous Reperfusion Therapy on Regional Myocardial Function and Geometry in Humans: a Topographic Assessment Using Two-Dimensional Echocardiography. *J Am Coll Cardiol.* 12:289-300.
- Verma, A., Meris, A., Skali, H., Ghali, J.K., Arnold, J.M., Bourgoun, M., Velaquez, E.J., McMurray, J.J., Kober, L., Pfeffer, M.A., Califf, R.M., Solomon, S.D. 2008. Prognostic Implications of Left Ventricular Mass and Geometry Following Myocardial Infarction: the VALIANT (VALsartan In Acute myocardial iNfarcTion) Echocardiographic Study. *JACC Cardiovasc Imaging.* 1(5):582-591.
- Wagner GS, Macfarlane P, Wellens H, Josephson M, Gorgels A, Mirvis DM, Pahlm O, Surawicz B, Kligfield P, Childers R, Gettes LS, Bailey JJ, Deal BJ, Gorgels A, Hancock EW, Kors JA, Mason JW, Okin P, Rautaharju PM, van Herpen G. 2009. AHA/ACCF/HRS recommendations for the standardization and interpretation of the electrocardiogram: Part VI: Acute ischemia/infarction. *J Am Coll Cardiol*; 53:1003-11
- Weir, R.A.P., Miller, A.M., Murphy G.E.J., Clements, S., Steedman, T., Connell, J.M.C., McInnes, I.B., Dargie, H.J., McMurray, J.J.V. 2010. Serum Soluble ST2 A Potential Novel Mediator in Left Ventricular and Infarct Remodeling After Myocardial Infarction. *Journal of the American College of Cardiology.* 55(3): 243-250.
- Weinberg, E. O., Shimpo, M., Keulenaer, G. W. De, Macgillivray, C., Tominaga, S., Solomon, S. D., Rouleau, J., Lee, R. T. 2002. Expression and Regulation of ST2, an Interleukin-1 Receptor Family Member , in Cardiomyocytes and Myocardial Infarction. *Circulation.* 106:2961–2967.
- Yang, H.S., Mur, M., Kim, H., Magrini, L., Marino, R., Di Somma, S. 2016. Soluble Suppression of Tumorigenicity 2 and Echocardiography in Sepsis. *Annals of Laboratory Medicine.* 36(6):590-594.
- Yingchoncharoen, T., Agarwal, S., Popovic, Z.B., Marwick, T.H. 2013. Normal Ranges of Left Ventricular Strain: a Meta-analysis. *J Am Soc Echocardiogr.* 26:185-191.
- Zaliaduonyte-Peksiene, D., Simonyte, S., Lesauskaite, V., Vaskelyte, J., Gustiene, O., Mizariene, V., Jurkevicius, R., Jariene, G., Tamosiunas, A., Zaliunas, R. 2014. Left Ventricular Remodelling After Acute Myocardial Infarction: Impact of Clinical, Echocardiographic

Parameters and Polymorphism of Angiotensinogen Gene. *Journal of the Renin-Angiotensin-Aldosterone System*. 15(3):286-293.

Zaliaduonyte-Peksiene, D., Vaskelyte, J.J., Mizariene, V., Jurkevicius, R., Zaliunas, R. 2012. Does Longitudinal Strain Predict Left Ventricular Remodeling After Myocardial Infarction?. *Echocardiography*. 29(4):419-427.