

REFERENCE

- Andiarsa, D. and Meliyanie, G. (2018). Lymphatic Filariasis Elimination Program in Indonesia. *Journal of Health Epidemiology and Communicable Diseases*, 3(2), pp.63-70.
- Bal, R., Erdogan, S., Theophilidis, G., Baydas, G. and Naziroglu, M. (2010). Assessing the effects of the neonicotinoid insecticide imidacloprid in the cholinergic synapses of the stellate cells of the mouse cochlear nucleus using whole-cell patch-clamp recording. *NeuroToxicology*, 31(1), pp.113-120.
- Belkin JN. 1977. *Quinquefasciatus* or *Fatigans* for the tropical (southern) house mosquito (Diptera: Culicidae). Proceedings of the Entomological Society of Washington.
- Cang, T., Dai, D., Yang, G., Yu, Y., Lv, L., Cai, L., Wang, Q. and Wang, Y. (2017). Combined toxicity of imidacloprid and three insecticides to the earthworm, *Eisenia fetida* (Annelida, Oligochaeta). *Environmental Science and Pollution Research*, 24(9), pp.8722-8730.
- Bates M. 1949. *The Natural History of Mosquitoes*. Macmillan Company. New York, NY. 379 pp.
- Cdc.gov. (2018). *CDC - Lymphatic Filariasis - Biology - Life Cycle of Wuchereria bancrofti*. [online] Available at: https://www.cdc.gov/parasites/lymphaticfilariasis/biology_w_bancrofti.html [Accessed 3 Apr. 2018].
- Christophers, S. (2009). *Aedes aegypti* (L.) the yellow fever mosquito. Cambridge: Cambridge University Press, p.412.
- Corbel, V., Duchon, S., Zaim, M. and Hougard, J. (2004). Dinotefuran: A Potential Neonicotinoid Insecticide Against Resistant Mosquitoes. *Journal of Medical Entomology*, 41(4), pp.712-717.
- Daniel, I. E., and Ekam, U. S. (2016). Preliminary Phytochemical Screening and Toxic Effect of *Melanthera scandens* Leaf Extracts Using Brine Shrimp (*Artemia salina*) Test. *Journal of Complementary and Alternative Medical Research*, 1(3), pp.1-7.
- Darsie Jr RF, Morris CD. 2000. Keys to the adult females and fourth-instar larvae of the mosquitoes of Florida (Diptera: Culicidae). *Technical Bulletin of the Florida Mosquito Control Association* 1: 148-155.
- Darsie Jr RF, Ward RA. 2005. *Identification and Geographical Distribution of the Mosquitoes of North America, North of Mexico*. University of Florida Press. Gainesville, FL. 300 pp.

- Gervais, J. A.; Luukinen, B.; Buhl, K.; Stone, D. 2010. Imidacloprid General Fact Sheet; National Pesticide Information Center, Oregon State University Extension Services. <http://npic.orst.edu/factsheets/imidagen.html>.
- Harbach, R. (2014). Culicidae Classification | Mosquito Taxonomic Inventory. [online] Mosquito-taxonomic-inventory.info. Available at: <http://mosquito-taxonomic-inventory.info/simpletaxonomy/term/6165> [Accessed 5 Oct. 2018].
- Lima CA, Almeida WR, Hurd H, Albuquerque CM. 2003. Reproductive aspects of the mosquito *Culex quinquefasciatus* (Diptera: Culicidae) infected with *Wuchereria bancrofti* (Spirurida: Onchocercidae). *Memórias do Instituto Oswaldo Cruz*. 98: 217-222.
- Sholichah, Z. (2009). Ancaman dari nyamuk *Culex* sp yang terabaikan. *Balaba*, 5(1), pp.21-23.
- Sirivanakarn S, White GB. 1978. Neotype designation of *Culex quinquefasciatus* Say (Diptera: Culicidae). *Proceedings of the Entomological Society of Washington* 80: 360-372.
- Srinivasa Rao, M.*et al.* (2008). Larvicidal Efficacy of Neonicotinoid Classes of Compounds on *Culex quinquefasciatus*. *Journal of Entomology*, 5(1), pp.45-50.
- Tomizawa, M. and Casida, J. (2005). Neonicotinoid Insecticide Toxicology: Mechanisms of Selective Action. *Annual Review of Pharmacology and Toxicology*, 45(1), pp.247-268.
- Uragayala, S., Verma, V., Natarajan, E., Velamuri, P. and Kamaraju, R. (2015). Adulticidal & larvicidal efficacy of three neonicotinoids against insecticide susceptible & resistant mosquito strains. *Indian Journal of Medical Research*, 142(7), p.64.
- WHO recommended classification of pesticides by hazard and guidelines to classification. (2010). Geneva: World Health Organization.
- WHO First Meeting of The Vector Control Advisory Group. (2013). Geneva: World Health Organization.
- World Health Organization. (2005). Guidelines for laboratory and field testing of mosquito larvicides. Geneva: WHO/Cds/Whopes/Gcdpp/2005.13; pp. 1–39.
- World Health Organization. (2006). Guidelines for testing mosquito adulticides for indoor residual spraying and treatment of mosquito nets. Geneva: WHO/Cds/Whopes/Gcdpp/2006.3.
- World Health Organization. (2016). Lymphatic filariasis. [online] Available at: <http://www.who.int/news-room/fact-sheets/detail/lymphatic-filariasis> [Accessed 4 Oct. 2018].

- World Health Organization. (2016). Monitoring and managing insecticide resistance in Aedes mosquito populations [online] Available at: http://apps.who.int/iris/bitstream/handle/10665/204588/WHO_ZIKV_VC_16.1_eng.pdf?sequence=2 [Accessed 4 Oct. 2018].
- Yamamoto, Izuru. 1999. Nicotin to Nicotinoids: 1962 to 1997. In Yamamoto, Izuru. Casida, John. Nicotinoid Insecticides and the Nicotinic Acetylcholine receptor. Springer-Verlag pp. 3-27.
- Zhu, F., Lavine, L., O'Neal, S., Lavine, M., Foss, C. and Walsh, D. (2016). Insecticide Resistance and Management Strategies in Urban Ecosystems. *Insects*, 7(1), p.2.