UNIVERSITAS
GADJAH MADA
NEONICOTINOID INSECTICIDE'S EFFICACY TOWARDS CULEX QUINQUEFASCIATUS (DIPTERA: CULICIDAE)
RYAN KIKO SAPHIRO, Dr. Budi Mulyaningsih, Apt., MS.; Dr. Drh. Sitti Rahmah Umniyati, SU.
Universitas Gadjah Mada, 2018 | Diunduh dari http://etd.repository.ugm.ac.id/

REFERENCE

Andiarsa, D. and Meliyanie, G. (2018). Lymphatic Filariasis Elimination Program in Indonesia. *Journal of Health Epidemiology and Communicable Diseases*, 3(2), pp.63-70.

Bal, R., Erdogan, S., Theophilidis, G., Baydas, G. and Naziroglu, M. (2010). Assessing the effects of the neonicotinoid insecticide imidacloprid in the cholinergic synapses of the stellate cells of the mouse cochlear nucleus using whole-cell patch-clamp recording. NeuroToxicology, 31(1), pp.113-120.

Belkin JN. 1977. Quinquefasciatus or Fatigans for the tropical (southern) house mosquito (Diptera: Culicidae). Proceedings of the Entomological Society of Washington.

Cang, T., Dai, D., Yang, G., Yu, Y., Lv, L., Cai, L., Wang, Q. and Wang, Y. (2017). Combined toxicity of imidacloprid and three insecticides to the earthworm, Eisenia fetida (Annelida, Oligochaeta). Environmental Science and Pollution Research, 24(9), pp.8722-8730.

Bates M. 1949. The Natural History of Mosquitoes. Macmillian Company. New York, NY. 379 pp.

Cdc.gov. (2018). *CDC - Lymphatic Filariasis - Biology - Life Cycle of Wuchereria bancrofti*. [online] Available at: https://www.cdc.gov/parasites/lymphaticfilariasis/biology_w_bancrofti.html [Accessed 3 Apr. 2018].

Christophers, S. (2009). *Aedes aegypti* (L.) the yellow fever mosquito. Cambridge: Cambridge University Press, p.412.

Corbel, V., Duchon, S., Zaim, M. and Hougard, J. (2004). Dinotefuran: A Potential Neonicotinoid Insecticide Against Resistant Mosquitoes. *Journal of Medical Entomology*, 41(4), pp.712-717.

Daniel, I. E., and Ekam, U. S. (2016). Preliminary Phytochemical Screening and Toxic Effect of Melanthera scandens Leaf Extracts Using Brine Shrimp (Artemia salina) Test. Journal of Complementary and Alternative Medical Research, 1(3), pp.1-7.

Darsie Jr RF, Morris CD. 2000. Keys to the adult females and fourth-instar larvae of the mosquitoes of Florida (Diptera: Culicidae). Technical Bulletin of the Florida Mosquito Control Association 1: 148-155.

Darsie Jr RF, Ward RA. 2005. Identification and Geographical Distribution of the Mosquitoes of North America, North of Mexico. University of Florida Press. Gainesville, FL. 300 pp.

UNIVERSITAS
GADJAH MADA
NEONICOTINOID INSECTICIDE'S EFFICACY TOWARDS CULEX QUINQUEFASCIATUS (DIPTERA: CULICIDAE)
RYAN KIKO SAPHIRO, Dr. Budi Mulyaningsih, Apt., MS.; Dr. Drh. Sitti Rahmah Umniyati, SU.
Universitas Gadjah Mada, 2018 | Diunduh dari http://etd.repository.ugm.ac.id/

Gervais, J. A.; Luukinen, B.; Buhl, K.; Stone, D. 2010. Imidacloprid General Fact Sheet; National Pesticide Information Center, Oregon State University Extension Services. http://npic.orst.edu/factsheets/imidagen.html.

Harbach, R. (2014). Culicidae Classification | Mosquito Taxonomic Inventory. [online] Mosquito-taxonomic-inventory.info. Available at: http://mosquito-taxonomic-inventory.info/simpletaxonomy/term/6165 [Accessed 5 Oct. 2018].

Lima CA, Almeida WR, Hurd H, Albuquerque CM. 2003. Reproductive aspects of the mosquito *Culex quinquefasciatus* (Diptera: Culicidae) infected with Wuchereria bancrofti (Spirurida: Onchocercidae). Memórias do Instituto Oswaldo Cruz. 98: 217-222.

Sholichah, Z. (2009). Ancaman dari nyamuk *Culex* sp yang terabaikan. *Balaba*, 5(1), pp.21-23.

Sirivanakarn S, White GB. 1978. Neotype designation of *Culex quinquefasciatus* Say (Diptera: Culicidae). Proceedings of the Entomological Society of Washington 80: 360-372.

Srinivasa Rao, M.*et al*. (2008). Larvicidal Efficacy of Neonicotinoid Classes of Compounds on *Culex quinquefasciatus.* Journal of Entomology, 5(1), pp.45-50.

Tomizawa, M. and Casida, J. (2005). Neonicotinoid Insecticide Toxicology: Mechanisms of Selective Action. Annual Review of Pharmacology and Toxicology, 45(1), pp.247-268.

Uragayala, S., Verma, V., Natarajan, E., Velamuri, P. and Kamaraju, R. (2015). Adulticidal & larvicidal efficacy of three neonicotinoids against insecticide susceptible & resistant mosquito strains. *Indian Journal of Medical Research*, 142(7), p.64.

WHO recommended classification of pesticides by hazard and guidelines to classification. (2010). Geneva: World Health Organization.

WHO First Meeting of The Vector Control Advisory Group. (2013). Geneva: World Health Organization.

World Health Organization. (2005). Guidelines for laboratory and field testing of mosquito larvicides. Geneva: WHO/Cds/Whopes/Gcdpp/2005.13; pp. 1–39.

World Health Organization. (2006). Guidelines for testing mosquito adultcides for indoor residual spraying and treatment of mosquito nets. Geneva: WHO/Cds/Whopes/Gcdpp/2006.3.

World Health Organization. (2016). Lymphatic filariasis. [online] Available at: http://www.who.int/news-room/fact-sheets/detail/lymphatic-filariasis [Accessed 4 Oct. 2018].

UNIVERSITAS
GADJAH MADA
NEONICOTINOID INSECTICIDE'S EFFICACY TOWARDS CULEX QUINQUEFASCIATUS (DIPTERA: CULICIDAE)
RYAN KIKO SAPHIRO, Dr. Budi Mulyaningsih, Apt., MS.; Dr. Drh. Sitti Rahmah Umniyati, SU.
Universitas Gadjah Mada, 2018 | Diunduh dari http://etd.repository.ugm.ac.id/

World Health Organization. (2016). Monitoring and managing insecticide resistance in Aedes mosquito populations [online] Available at: http://apps.who.int/iris/bitstream/handle/10665/204588/WHO_ZIKV_VC_16.1_eng.pdf?sequence=2 [Accessed 4 Oct. 2018].

Yamamoto, Izuru. 1999. Nicotin to Nicotinoids: 1962 to 1997. In Yamamoto, Izuru. Casida, John. Nicotinoid Insecticides and the Nicotinic Acetylcholine receptor. Springer-Verlag pp. 3-27.

Zhu, F., Lavine, L., O'Neal, S., Lavine, M., Foss, C. and Walsh, D. (2016). Insecticide Resistance and Management Strategies in Urban Ecosystems. Insects, 7(1), p.2.