

REFERENCE

- Behera, P., Gupta, S., Nongkynrih, B., Kant, S., Mishra, A. and Sharan, P. (2017). Screening instruments for assessment of depression. *Indian Journal of Medical Specialities*, 8(1), pp.31-37.
- Bennett, D. (2001). Neurotrophic Factors: Important Regulators of Nociceptive Function. *The Neuroscientist*, 7(1), pp.13-17.
- Brennan-Olsen, S., Cook, S., Leech, M., Bowe, S., Kowal, P., Naidoo, N., et al. (2017). Prevalence of arthritis according to age, sex and socioeconomic status in six low and middle income countries: analysis of data from the World Health Organization study on global AGEing and adult health (SAGE) Wave 1. *BMC Musculoskeletal Disorders*, 18(1).
- Choy, E. (2012). Understanding the dynamics: pathways involved in the pathogenesis of rheumatoid arthritis. *Rheumatology*, 51(suppl 5), pp.v3-v11.
- Dean, J. and Keshavan, M. (2017). The neurobiology of depression: An integrated view. *Asian Journal of Psychiatry*, 27, pp.101-111.
- Forsgren, S. (2009). New data favouring that neurotrophins are of importance in arthritis. *Arthritis Research & Therapy*, 11(4), p.122.
- Forsgren, S., Grimsholm, O., Dalén, T. and Rantapää-Dahlqvist, S. (2011). Measurements in the Blood of BDNF for RA Patients and in Response to Anti-TNF Treatment Help Us to Clarify the Magnitude of Centrally Related Pain and to Explain the Relief of This Pain upon Treatment. *International Journal of Inflammation*, 2011, pp.1-7.
- Fournier, J., DeRubeis, R., Hollon, S., Dimidjian, S., Amsterdam, J., Shelton, R., et al. (2010). Antidepressant Drug Effects and Depression Severity. *JAMA*, 303(1), p.47.
- Giles, J. (2010). Are the Incidence and Prevalence of RA Changing in the United States?. [online] Johns Hopkins Arthritis Center. Available at: <https://www.hopkinsarthritis.org/arthritis-news/arthritis-news-are-the-incidence-and-prevalence-of-ra-changing-in-the-united-states/> [Accessed 17 Oct. 2018].

- Sadock, B., Sadock, V. and Ruiz, P. (2015). Kaplan & Sadock's synopsis of psychiatry. Philadelphia: Wolter Kluwer/Lippincott Williams & Wilkins.
- Klein, A., Williamson, R., Santini, M., Clemmensen, C., Ettrup, A., Rios, M., et al. (2010). Blood BDNF concentrations reflect brain-tissue BDNF levels across species. *The International Journal of Neuropsychopharmacology*, 14(03), pp.347-353.
- Margaretten, M., Julian, L., Katz, P. and Yelin, E. (2011). Depression in patients with rheumatoid arthritis: description, causes and mechanisms. *International Journal of Clinical Rheumatology*, 6(6), pp.617-623.
- Murphy, J. (2017). Rheumatoid Arthritis. [online] American College of Rheumatology. Available at: <https://www.rheumatology.org/I-Am-A/Patient-Caregiver/Diseases-Conditions/Rheumatoid-Arthritis> [Accessed 5 Sep. 2018].
- Ogłodek, E., Just, M., Szromek, A. and Araszkievicz, A. (2016). Melatonin and neurotrophins NT-3, BDNF, NGF in patients with varying levels of depression severity. *Pharmacological Reports*, 68(5), pp.945-951.
- Pedard, M., Demougeot, C., Prati, C. and Marie, C. (2018). Brain-derived neurotrophic factor in adjuvant-induced arthritis in rats. Relationship with inflammation and endothelial dysfunction. *Progress in Neuro-Psychopharmacology and Biological Psychiatry*, 82, pp.249-254.
- Phillips, C. (2017). Brain-Derived Neurotrophic Factor, Depression, and Physical Activity: Making the Neuroplastic Connection. *Neural Plasticity*, 2017, pp.1-17.
- Rathbun, AM. (2014). Depression in Rheumatoid Arthritis and an Estimation of the Bi-directional Association of Depression and Disease Burden: A Dissertation. University of Massachusetts Medical School. *GSBS Dissertations and Theses*. Paper 699. DOI:10.13028/M2RC8S. http://escholarship.umassmed.edu/gsbs_diss/699
- Riset Kesehatan Dasar(Riskesdas). (2013). Badan Penelitian dan Pengembangan Kesehatan Kementerian RI tahun 2013.

<http://www.depkes.go.id/resources/download/general/Hasil%20Ris-kesdas%202013.pdf>. [Accessed 17 April 2018]

- Simão, A., Mendonça, V., de Oliveira Almeida, T., Santos, S., Gomes, W., Coimbra, C., et al. (2014). Involvement of BDNF in knee osteoarthritis: the relationship with inflammation and clinical parameters. *Rheumatology International*.
- Symmons, D. (2002). Epidemiology of rheumatoid arthritis: determinants of onset, persistence and outcome. *Best Practice & Research Clinical Rheumatology*, 16(5), pp.707-722.
- Thompson, S., Bennett, D., Kerr, B., Bradbury, E. and McMahon, S. (1999). Brain-derived neurotrophic factor is an endogenous modulator of nociceptive responses in the spinal cord. *Proceedings of the National Academy of Sciences*, 96(14), pp.7714-7718.
- van Vollenhoven, R. (2009). Sex differences in rheumatoid arthritis: more than meets the eye. *BMC Medicine*, 7(1).
- Wirasto, R. (2017). Cognitive Visual Perception Therapy for Reducing Depression in Patients with Rheumatoid Arthritis. Ph.D. Universitas Gadjah Mada.
- Yu, H. and Chen, Z. (2010). The role of BDNF in depression on the basis of its location in the neural circuitry. *Acta Pharmacologica Sinica*, 32(1), pp.3-11.