UNIVERSITAS
GADJAH MADA
**Effect of Centella asiatica (L.) Urban Extract on the Expression of Heme Oxygenase-1 mRNA in Kidney
of Type II Diabetes Wistar Rat Model**
BERNADHET DAISY K, dr. Dwi Aris Agung N, M.Sc., Ph.D, 2. Dr. dr. Woro Rukmi P, M.Kes., Sp.PD
Universitas Gadjah Mada, 2018 | Diunduh dari http://etd.repository.ugm.ac.id/

REFERENCES

Abdou HM, Mohamed NA (2016). *Centella asiatica* Ameliorates Diabetic Complications and Oxidative Stress in Streptozotocin-Induced Diabetes Mellitus in Male Rats. *Int J Pharm Sci Rev Res*, 36(2):61–70.

Agarwal A, Nick HS (2000). Renal Response to Tissue Injury: Lessons from Heme Oxygenase-1 Gene Ablation and Expression. *J Am Soc Nephrol,* 11:965–973.

Alam J, Killeen E, Gong P, Naquin R, Hu B, Stewart D, Ingelfinger JR, Nath KA. (2003). Heme activates the heme oxygenase-1 gene in renal epithelial cells by stabilizing Nrf2. *Am J Physiol Renal Physiol*, 284(4):F743–F752.

Anon (2017). 2. Classification and Diagnosis of Diabetes. *Diabetes Care*, 40(Supplement 1):S11–S24.

Avogaro A, Albiero M, Menegazzo L, Kreutzenberg SD, Fadini GP (2011). Endothelial Dysfunction in Diabetes: The role of reparatory mechanisms. *Diabetes Care*, 34(Supplement 2):S285–S290.

Baranano DE, Rao M, Ferris CD, Snyder SH (2002). Biliverdin reductase: A major physiologic cytoprotectant. *Proceedings of the National Academy of Sciences*, 99(25):16093–16098.

Bhaskaran M, Reddy K, Radhakrishanan N, Franki N, Ding G, Singhal PC. (2003). Angiotensin II induces apoptosis in renal proximal tubular cells. *Am J Physiol Renal Physiol*, 284(5):F955–F965.

Bloomgarden ZT (2008). Diabetic Nephropathy. *Diabetes Care*, 31(4):823–827.

Butler AE, Janson J, Bonner-Weir S, Ritzer R, Rizza RA, Butler PC (2003). β-Cell Deficit and Increased β-Cell Apoptosis in Humans With Type 2 Diabetes. *Diabetes*, 52(1):102–110.

Cheng H, Harris R (2014). Renal Endothelial Dysfunction in Diabetic Nephropathy. *Cardiovasc Hematol Disord Drug Targets*, 14(1):22–33.

Farrell RE (2010). *RNA methodologies: a laboratory guide for isolation and characterization*, 4th ed. Amsterdam, Elsevier, Academic.

Fleige S, Pfaffl MW (2006). RNA integrity and the effect on the real-time qRT-PCR performance. *Mol Aspects Med*, 27(2–3):126–139.

Gozzelino R, Jeney V, Soares MP (2010). Mechanisms of Cell Protection by Heme Oxygenase-1. *Annu Rev Pharmacol Toxicol*, 50(1):323–354.

UNIVERSITAS GADJAH MADA
Effect of Centella asiatica (L.) Urban Extract on the Expression of Heme Oxygenase-1 mRNA in Kidney of Type II Diabetes Wistar Rat Model
BERNADHET DAISY K, dr. Dwi Aris Agung N, M.Sc., Ph.D, 2. Dr. dr. Woro Rukmi P, M.Kes., Sp.PD
Universitas Gadjah Mada, 2018 | Diunduh dari http://etd.repository.ugm.ac.id/

Griffin KA, Bidani AK (2009). Angiotensin II type 2 receptor in chronic kidney disease: the good side of angiotensin II? *Kidney Int* 75(10):1006–1008.

Hayashi K, Haneda M, Koya D, Maeda S, Isshiki K, Kikkawa R (2001). Enhancement of glomerular heme oxygenase-1 expression in diabetic rats. *Diabetes Res Clin Pract*, 52(2):85–96.

Kabir AU, Samad MB, D'Costa NM, Akhter F, Ahmed A, Hannan JMA (2014). Anti-hyperglycemic activity of Centella asiatica is partly mediated by carbohydrase inhibition and glucose-fiber binding. *BMC Complement Altern Med*, 14(31)1–14.

Kanwar YS, Sun L, Xie P, Liu F, Chen S. (2011). A Glimpse of Various Pathogenetic Mechanisms of Diabetic Nephropathy. *Annu Rev Pathol*, 6(1):395–423.

Kumari S, Kumar SH, Navis S (2015). Angiotensin Converting Enzyme (Ace) Inhibitors in the Treatment of Diabetic Nephropathy. *Int J Pharm Phytopharmacol*, 4(4):211–217.

L'Abbate A, Neglia D, Vecoli C, Novelli M, Ottaviano V, Baldi S, Barsacchi R, Paolicchi A, Masiello P, Drummond GS, McClung JA, Abraham NG (2007). Beneficial effect of heme oxygenase-1 expression on myocardial ischemia-reperfusion involves an increase in adiponectin in mildly diabetic rats. *Am J Physiol Heart Circ Physiol*, 293(6):H3532–H3541.

Lee SC, Han SH, Li JJ, Lee SH, Jung D, Kwak S, Kim SH, Kim DK, Yoo TH, Kim JH, Chang S, Han DS, Kang S (2009). Induction of heme oxygenase-1 protects against podocyte apoptosis under diabetic conditions. *Kidney Int*, 76(8):838–848.

Lee T-S, Chau L-Y (2002). Heme oxygenase-1 mediates the anti-inflammatory effect of interleukin-10 in mice. *Nat Med*, 8(3):240–246.

Lever JM, Boddu R, George JF, Agarwal A (2016). Heme Oxygenase-1 in Kidney Health and Disease. *Antioxid Redox Signal*, 25(3):165–183.

Ling APK, Chan HH, Koh RY, Wong YP (2017). Neuroprotective Rols of Asiaticoside on Hydrogen Peroxide-Induced Toxicity in SH-SY5Y Cells. *J Fundam Appl Sci*, 9(7S):636–649.

Liu J, He T, Lu Q, Shang J, Sun H, Zhang L (2010). Asiatic acid preserves beta cell mass and mitigates hyperglycemia in streptozocin-induced diabetic rats. *Diabetes Metab Res Rev*, 26(6):448–454.

Ndisang JF (2010). Role of Heme Oxygenase in Inflammation, Insulin-Signalling, Diabetes and Obesity. *Mediators Inflamm*, 2010:1–18.

UNIVERSITAS
GADJAH MADA
**Effect of Centella asiatica (L.) Urban Extract on the Expression of Heme Oxygenase-1 mRNA in Kidney
of Type II Diabetes Wistar Rat Model**
BERNADHET DAISY K, dr. Dwi Aris Agung N, M.Sc., Ph.D, 2. Dr. dr. Woro Rukmi P, M.Kes., Sp.PD
Universitas Gadjah Mada, 2018 | Diunduh dari http://etd.repository.ugm.ac.id/

Nordquist L, Friederich-Persson M, Fasching A, Liss P, Shoji K, Nangaku M, Hansell P, Palm P. (2015). Activation of Hypoxia-Inducible Factors Prevents Diabetic Nephropathy. *J Am Soc Nephrol*, 26(2):328–338.

Parving H-H, Hommel E, Jensen BR, Hansen HP. (2001). Long-term beneficial effect of ACE inhibition on diabetic nephropathy in normotensive type 1 diabetic patients. *Kidney Int*, 60(1):228–234.

Persson P, Palm F (2017). Hypoxia-inducible factor activation in diabetic kidney disease. *Curr Opin Nephrol Hy*, 26(5):345–350.

Pfeiffer AFH, Klein HH (2014). The Treatment of Type 2 Diabetes. *Dtschs Arzteblt Int*. 111(5):69–82

Pham CG, Bubici C, Zazzeroni F, Papa S, Jones J, Alvarez K, Jayawardena S, Smaele ED, Cong R, Beaumont C, Torti FM, Torti SV, Franzoso G (2004). Ferritin Heavy Chain Upregulation by NF-κB Inhibits TNFα-Induced Apoptosis by Suppressing Reactive Oxygen Species. *Cell.* 119:529–542.

Ponka P (1999). Cell Biology of Heme. *The American Journal of the Medical Sciences*, 318(4):241–256.

Ryter SW, Alam J, Choi AMK (2006). Heme Oxygenase-1/Carbon Monoxide: From Basic Science to Therapeutic Applications. *Physiol Rev*, 86(2):583–650.

Santos CF dos, Sakai VT, Machado MAAM, Schippers DN, Greene AS (2004). Reverse transcription and polymerase chain reaction: principles and applications in dentistry. *J Appl Oral Sci*, 12(1):1–11.

Suttner DM, Dennery PA (1999). Reversal of HO-1 related cytoprotection with increased expression is due to reactive iron. *FASEB J*, 13(13):1800–1809.

Tang SCW, Chan GCW, Lai KN (2016). Recent advances in managing and understanding diabetic nephropathy. *F1000Res*, 5:1044.

Toth-Manikowski S, Atta MG (2015). Diabetic Kidney Disease: Pathophysiology and Therapeutic Targets. *J Diabetes Res*, 2015:1–16.

Tracz MJ, Alam J, Nath KA (2007). Physiology and Pathophysiology of Heme: Implications for Kidney Disease. *J Am Soc Nephrol*, 18(2):414–420.

Wan J, Gong X, Jiang R, Zhang Z, Zhang L. (2013). Antipyretic and Anti-inflammatory Effects of Asiaticoside in Lipopolysaccharide-treated Rat through Up-regulation of Heme Oxygenase-1. *Phytother Res*, 27(8):1136–1142.

UNIVERSITAS
GADJAH MADA
**Effect of Centella asiatica (L.) Urban Extract on the Expression of Heme Oxygenase-1 mRNA in Kidney
of Type II Diabetes Wistar Rat Model**
BERNADHET DAISY K, dr. Dwi Aris Agung N, M.Sc., Ph.D, 2. Dr. dr. Woro Rukmi P, M.Kes., Sp.PD
Universitas Gadjah Mada, 2018 | Diunduh dari http://etd.repository.ugm.ac.id/

Wegiel B, Otterbein LE (2012). Go Green: The Anti-Inflammatory Effects of Biliverdin Reductase. *Front Pharmacol*, 3(47):1–8.

Weyer C, Funahashi T, Tanaka S, Hotta K, Matsuzawa Y, Pratley RE, Tataranni PA (2001). Hypoadiponectinemia in Obesity and Type 2 Diabetes: Close Association with Insulin Resistance and Hyperinsulinemia. *J Clin Endocrinol Metab*, 86(5):1930–1935.

Wilks A (2002). Heme Oxygenase: Evolution, Structure, and Mechanism. *Antioxid Redox Signal*, 4(4):603–614.

Wolf G, Butzmann U, Wenzel UO (2003). The Renin-Angiotensin System and Progression of Renal Disease: From Hemodynamics to Cell Biology. *Nephron Physiol*, 93(1):p3–p13.

Yang L, Quan S, Nasjletti A, Laniado-Schartzman M, Abraham NG (2004). Heme Oxygenase-1 Gene Expression Modulates Angiotensin II–Induced Increase in Blood Pressure. *Hypertension*, 43(6):1221–1226.

Yang Z-Z, Zou A-P (2001). Transcriptional regulation of heme oxygenases by HIF-1α in renal medullary interstitial cells. *Am J Physiol Renal Physiol*, 281(5):F900–F908.

Yim HE, Kim JH, Yoo KH, Bae IS, Hong YS, Lee JW. (2011). Spironolactone and Enalapril Differentially Up-Regulate the Expression of VEGF and Heme Oxygenase-1 in the Neonatal Rat Kidney: *Pediatr Res*, 69(5 Part 1):378–383.

Yoshida T, Biro P, Cohen T, Müller RM, Shibahara S, (1988). Human heme oxygenase cDNA and induction of its mRNA by hemin. *Eur J Biochem*, 171(3):457–461.