

REFERENCES

- Alemzadeh, R. and Kichler, J. (2014). Gender differences in the association of insulin resistance and high-sensitivity c-reactive protein in obese adolescents. *Journal of Diabetes & Metabolic Disorders*, 13(1), p.35.
- Balkau, B., Deanfield, J., Després, J., Bassand, J., Fox, K., Smith, S., Barter, P., Tan, C., Van Gaal, L., Wittchen, H., Massien, C. and Haffner, S. (2007). International Day for the Evaluation of Abdominal Obesity (IDEA). *Circulation*, 116(17), pp.1942-1951.
- Bibiloni, M., Pons, A. and Tur, J. (2013). Prevalence of Overweight and Obesity in Adolescents: A Systematic Review. *ISRN Obesity*, 2013, pp.1-14.
- Boden, G. (2009). Endoplasmic Reticulum Stress: Another Link Between Obesity and Insulin Resistance/Inflammation?. *Diabetes*, 58(3), pp.518-519.
- Cdc.gov. (2016). Defining Childhood Obesity | Overweight & Obesity | CDC. [online] Available at: <https://www.cdc.gov/obesity/childhood/defining.html> [Accessed 9 Aug. 2016]
- Cempaka, V. and Sidiartha, I. G. (2018) "Waist circumference and insulin levels in obese children", *Paediatrica Indonesiana*, 57(4), pp. 194-7. doi: 10.14238/pi57.4.2017.194-7.
- Cheng, Y., Tsao, Y., Tzeng, I., Chuang, H., Li, W., Tung, T. and Chen, J. (2017). Body mass index and waist circumference are better predictors of insulin resistance than total body fat percentage in middle-aged and elderly Taiwanese. *Medicine*, 96(39), p.e8126.
- Departemen Kesehatan Republik Indonesia. Laporan Hasil Riset Kesehatan Dasar Indonesia (Riskesdas). 2013
- Dietz, W. and Bellizzi, M. (1999). Introduction: the use of body mass index to assess obesity in children. *The American Journal of Clinical Nutrition*, 70(1), pp.123S-125S.
- Esteghamati, A., Ashraf, H., Khalilzadeh, O., Zandieh, A., Nakhjavani, M., Rashidi, A., Haghazali, M. and Asgari, F. (2010). Optimal cut-off of homeostasis model assessment of insulin resistance (HOMA-IR) for

the diagnosis of metabolic syndrome: third national surveillance of risk factors of non-communicable diseases in Iran (SuRFNCD-2007). *Nutrition & Metabolism*, 7(1), p.26.

Geer, E. and Shen, W. (2009). Gender differences in insulin resistance, body composition, and energy balance. *Gender Medicine*, 6, pp.60-75.

Geneva: World Health Organization; 2008. World Health Organization (WHO). Waist Circumference and Waist-Hip Ratio. Report of WHO Expert Consultation.

Hardy, O., Czech, M. and Corvera, S. (2012). Curr Opin Endocrinol Diabetes. 19(2): 81–87. doi:10.1097/MED.0b013e3283514e13

Karpe, F., Dickmann, J. and Frayn, K. (2011). Fatty Acids, Obesity, and Insulin Resistance: Time for a Reevaluation. *Diabetes*, 60(10), pp.2441-2449.

Keskin, M., Kurtoglu, S., Kendirci, M., Atabek, M. and Yazici, C. (2005). Homeostasis Model Assessment Is More Reliable Than the Fasting Glucose/Insulin Ratio and Quantitative Insulin Sensitivity Check Index for Assessing Insulin Resistance Among Obese Children and Adolescents. *PEDIATRICS*, 115(4), pp.e500-e503.

Lentferink, Y., Elst, M., Knibbe, C. and van der Vorst, M. (2017). Predictors of Insulin Resistance in Children versus Adolescents with Obesity. *Journal of Obesity*, 2017, pp.1-7.

Lim, S., Choi, D., Rhee, Y. and Kim, H. (2015). Association between Obesity Indices and Insulin Resistance among Healthy Korean Adolescents: The JS High School Study. *PLOS ONE*, 10(5), p.e0125238.

Ling, J., Mohamed, M., Jalaludin, M., Rampal, S., Zaharan, N. and Mohamed, Z. (2016). Determinants of High Fasting Insulin and Insulin Resistance Among Overweight/Obese Adolescents. *Scientific Reports*, 6(1).

Maffeis, C., Corciulo, N., Livieri, C., Rabbone, I., Trifirò, G., Falorni, A., Guerraggio, L., Peverelli, P., Cuccarolo, G., Bergamaschi, G., Di Pietro, M. and Grezzani, A. (2003). Waist circumference as a predictor of cardiovascular and metabolic risk factors in obese girls. *European Journal of Clinical Nutrition*, 57(4), pp.566-572.

- Mehdad, S., Hamrani, A., El Kari, K., El Hamdouchi, A., Barakat, A., El Mzibri, M., Mokhtar, N. and Aguenau, H. (2012). Body Mass Index, Waist Circumference, Body Fat, Fasting Blood Glucose in a Sample of Moroccan Adolescents Aged 11–17 Years. *Journal of Nutrition and Metabolism*, 2012, pp.1-7.
- Méthot, J., Houle, J. and Poirier, P. (2010). Obesity: how to define central adiposity?. *Expert Review of Cardiovascular Therapy*, 8(5), pp. 639-644.
- Morais, P., Sousa, A., Jardim, T., Nascente, F., Mendonça, K., Povia, T., Carneiro, C., Ferreira, V., Souza, W. and Jardim, P. (2016). Correlation of Insulin Resistance with Anthropometric Measures and Blood Pressure in Adolescents. *Arquivos Brasileiros de Cardiologia*.
- Nambiar, S., Truby, H., Davies, P. and Baxter, K. (2013). Use of the waist-height ratio to predict metabolic syndrome in obese children and adolescents. *Journal of Paediatrics and Child Health*, 49(4), pp.E281-E287.
- Parikh, R., Joshi, S., Menon, P. and Shah, N. (2007). Index of central obesity – A novel parameter. *Medical Hypotheses*, 68(6), pp.1272-1275.
- Pinhas-Hamiel, O. and Zeitler, P. (2017). The global spread of type 2 diabetes mellitus in children and adolescents.
- Romualdo, M., de Nóbrega, F. and Escrivão, M. (2014). Insulin resistance in obese children and adolescents. *Jornal de Pediatria*, 90(6), pp. 600-607.
- Ropelle, E., Pauli, J., Cintra, D., da Silva, A., De Souza, C., Guadagnini, D., Carvalho, B., Caricilli, A., Katashima, C., Carvalho-Filho, M., Hirabara, S., Curi, R., Velloso, L., Saad, M. and Carvalheira, J. (2012). Targeted Disruption of Inducible Nitric Oxide Synthase Protects Against Aging, S-Nitrosation, and Insulin Resistance in Muscle of Male Mice. *Diabetes*, 62(2), pp.466-470.
- Tabata, S., Yoshimitsu, S., Hamachi, T., Abe, H., Ohnaka, K. and Kono, S. (2009). Waist circumference and insulin resistance: a cross-sectional study of Japanese men. *BMC Endocrine Disorders*, 9(1).

Third Report of the National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III) Final Report. (2002). *Circulation*, 106(25), pp.3143-3143.

Wei, M., Gaskill, S., Haffner, S. and Stern, M. (1997). Waist Circumference as the Best Predictor of Noninsulin Dependent Diabetes Mellitus (NIDDM) Compared to Body Mass Index, Waist/hip Ratio and Other Anthropometric Measurements in Mexican Americans-A 7-Year Prospective Study. *Obesity Research*, 5(1), pp.16-23.

Weyer, C., Bogardus, C., Mott, D. and Pratley, R. (1999). The natural history of insulin secretory dysfunction and insulin resistance in the pathogenesis of type 2 diabetes mellitus. *Journal of Clinical Investigation*, 104(1), pp.43-49.

World Health Organization. (2017). Obesity and overweight. [online] Available at: <http://www.who.int/mediacentre/factsheets/fs311/en/> [Accessed 2 Dec. 2017].

World Health Organization. (2017). Diabetes. [online] Available at: <http://www.who.int/mediacentre/factsheets/fs312/en/> [Accessed 2 Dec. 2017].

World Health Organization. (2007). Growth reference 5-19 years: Height-for-age (5-19 years). [online] Available at: https://www.who.int/growthref/who2007_height_for_age/en/ [Accessed 2 Dec. 2018].

Zimmet, P., Alberti, K., Kaufman, F., Tajima, N., Silink, M., Arslanian, S., Wong, G., Bennett, P., Shaw, J. and Caprio, S. (2007). The metabolic syndrome in children and adolescents ? an IDF consensus report. *Pediatric Diabetes*, 8(5), pp.299-306.