

DAFTAR PUSTAKA

- Abdel-Kader, K., Unrul, M.L., Weisbord, S.D., 2009. Symptom Burden, Depression, and Quality of Life in Chronic and End-Stage Kidney Disease. *Clin J Am Soc Nephrol* 4: 1057– 64.
- Agarwal, D.K., Sharma, A., Gupta, A., 2000. Peritoneal equilibration test in Indian patients on continuous ambulatory peritoneal dialysis: Does it affect patient outcome?. *Adv Perit Dial* 16 : 148-151.
- Aitken, G.R., Roderick, P.J., Fraser, S., Mindell, J.S., O'Donoghue, D., Day, J. *et al.*, 2014. Change in prevalence of chronic kidney disease in England over time: comparison of nationally representative cross-sectional surveys from 2003 to 2010. *BMJ Open*: 339-49.
- Amir, N., 2005. *Depresi Aspek Neurobiologi Diagnosis dan Tatalaksana*. Fakultas Kedokteran Universitas Indonesia, Jakarta.
- Andrade, C. P., Sesso, R. C., 2012. Depression in Chronic Kidney Disease and Hemodialysis Patients. *Psychology* 2012. Vol.3, No.11 : 974-8.
- Anggraeni, K.N., Sarwono, B., Sunarmi., 2017. Hubungan dukungan Keluarga dengan Tingkat Depresi pada Pasien yang Menjalani terapi Hemodialisa di Unit Hemodialisa Rumah Sakit Tentara Dr. Soedjono Magelang. *Jurnal Keperawatan Soedirman (The Soedirman Journal of Nursing)*, Volume 12, No.2 Juli 2017: 109 – 17.
- APA. 2013. Diagnostic and statistical manual of mental disorder fifth edition (DSM-V). *American Psychiatric Association*. 155 – 87.
- Bautovich, A., Katz, I., Smith, M., Loo, C. K., Harvey S.B., 2014. Depression and chronic kidney disease: A review for clinicians. *Australian & New Zealand Journal of Psychiatry*: 1-12.
- Baykan, H., Yargic, I., 2012. Depression, Anxiety Disorders, Quality of Life and Stress Coping Strategies in Hemodialysis and Continuous Ambulatory Peritoneal Dialysis Patients. *Bulletin of Clinical Psychopharmacology*: 167-76.
- Beck, A.T., Steer, R.A., Brown, G.K., 1996. *Beck Depression Inventory, Manual 2nd edition*. San Antonio: The Psychological corporation.

- Brown, E.A., Davies, S.J., Rutherford, P., 2003. Survival of functionally anuric patients on automated peritoneal dialysis: the European APD Outcome Study. *J Am Soc Nephrol* 14: 294S-S7.
- Busch, M.A., Maske, U.E., Ryl L., Schlak, R., Hapke, U., 2013. Prevalence of depressive symptoms and diagnosed depression among adults in Germany. *Bundesgesundheitsbl*:733 – 9.
- Chan, M.K., Cheung, C.Y., Chan, Y.H., Chan, H.W., Chak, W.L., Chau, K.F., 2015. Prevalence and impact of anxiety and depression in Chinese peritoneal dialysis patients: a single center study. *doi: 10.1111/nep.12970*.
- Chaudhary, K., Sangha, H., Khanna, R., 2011. Peritoneal dialysis first: rationale. *Clin J Am Soc Nephrol* :447 - 56.
- Coresh, J., Selvin, E., Stevens, L.A., Manzi, J., Kusek, J.W., Eggers, P. *et al.*, 2007. Prevalence of chronic kidney disease in the United States. *JAMA* :2038–47.
- Dahlan, M. S., 2010. *Besar Sampel dan Cara Pengambilan Sampel Seri 2 Evidence Based Medicine*, Pp: 76. Salemba Medika, Jakarta.
- Davies, S.J., Mushahar, L., Yu, Z., Lambie, M., 2011. Determinants of peritoneal membrane function over time. *Semin Nephrol*. 31(2): 172-82.
- Fadem, S.Z., 2017. Charlson Comorbidity Scoring System Estimating Prognosis for Dialysis Patients. Dari: <http://touchcalc.com/calculators/cci-js>.
- Farinde, A., 2013. The Beck Depression Inventory. *The Pharma Innovation Journal Vol 2* :56-63.
- Forget, P., Khalifa, C., Defour, JP., Latinne, D., Van Pel, M., De Kock, M., 2017. What is normal value of the neutrophil-to-lymphocyte ratio?. *BMC Res Notes*: 1-4.
- Gansevoort, R.T., Correa-Rotter, R., Hemmelgarn, B.R., Jafar, T.H., Heerspink, H.J., Mann, J.F., *et al.*, 2013. Chronic kidney disease and cardiovascular risk: epidemiology, mechanisms, and prevention. *Lancet*: 339-49.
- Ginting, H., Naring, G., van der Veld, V.M., Srisayekti, W., Becker, E.S., 2013. Validating the Beck Depression Inventory-II in Indonesia's general population and coronary heart disease patients. *Int J Clin health Psychol*: 235-42.

- Güney, I., Biyik, M., Yeksan, M., Biyik, Z., Atalay, H., Solak, Y. *et al.*, 2015. Sleep Quality and Depression in Peritoneal Dialysis Patients. *Renal Failure*, 30: 1017–22.
- Harwood, L., Wilson, B., Locking-Cusolito, H., Sontrop, J., Spittal, J., 2009. Stressors and Coping in Individuals with Chronic Kidney Disease. *Nephrology Nursing Journal Vol. 36 No. 3* : 265-75.
- Ho-dac-Pannekeet, M.M., 1997. Analysis of ultrafiltration failure in peritoneal dialysis patients by means of standard peritoneal permeability analysis. *Perit Dial Int* 17(2):144-150.
- Hedayati, S.S., Bosworth, H.B., Briley, L.P., Sloane, R.J., Pieper, C.F., Kimmel, P.L. *et al.*, 2008. Death or hospitalization of patients on chronic hemodialysis is associated with a physician- based diagnosis of depression. *Kidney Int* 74: 930–36.
- Hedayati, S.S., Finkelstein, F.O., 2009. Epidemiology, diagnosis, and management of depression in patients with CKD. *Am J Kidney Dis*: 741-52.
- Hedayati, S.S., Minhajuddin, A.T., Afshar, M., Toto, R.D., Trivedi, M.H., Rush, A.J., 2010. Association between major depressive episodes in patients with chronic kidney disease and initiation of dialysis, hospitalization, or death. *JAMA*: 1946 – 53.
- Hill, N.R., Fatoba, S.T., Oke, J.L., Hirst, J.A., O’Challagan, C.A., Lasersson, D. *et al.*, 2016. Global Prevalence of Chronic Kidney Disease – A Systematic Review and Meta-Analysis. *PLoS ONE*: 1-18.
- Heimbürger, O., Wang, T., Lindholm, B., 1999. Alterations in water and solute transport with time on peritoneal dialysis. *Perit Dial Int. 19 Suppl 2*:583-590.
- Howell, C., Marshall. C., Opolski M., 2008. Management of recurrent depression. *Australian Family Physician Vol. 37, No. 9.*: 704-8.
- Indonesian Renal Registry (IRR), 2014. *7th Report Of Indonesian Renal Registry*: 1-36.
- Johnson, J., Weissman, M.M., Klerman, G.L., 1992. Service utilization and social morbidity associated with depressive symptoms in the . *JAMA*:1478-83.
- Jung, H., Jang, H., M., Kim, Y.W., Cho, S., Kim, H., *et al.*, 2016. Depressive Symptoms, Patient Satisfaction, and Quality of Life Over Time in Automated and Continuous Ambulatory Peritoneal Dialysis Patients. *Medicine Volume 95 Number 21*: 1-10.

- Kemenkes, 2017. *Situasi Penyakit Ginjal Kronis*. Jakarta : 1-9.
- Kidney Disease Improving Global Outcomes (KDIGO) Team, 2012. KDIGO 2012 clinical practice guideline for the evaluation and management of chronic kidney disease. *Kid. Internat. Suppl 3*: 5-14.
- KDOQI, 2002. Chronic Kidney Disease: Evaluation, Classification, and Stratification. diakses: http://www.kidney.org/professionals/kdoqi/guidelines_commentaries.cfm.
- Levey, A.S., Stevens, L.A., Schmid, C.H., Zhang, Y.L., Castro, A.F., Feldman, H.I. *et al.*, 2009. A new equation to estimate glomerular filtration rate. *Ann Intern Med*: 604–12.
- Linan, S., 2002. Dialysis training in the United States. *Am J Kidney Dis*: 208-209.
- Lo, W.K., Bargman, J.M., Burkart, J., 2006. Guideline on targets for solute and fluid removal in adult patients on chronic peritoneal dialysis. *Perit Dial* 26: 520-2.
- Lopes, A.A., Bragg, J., Young, E., Goodkin, D., Mapes, J., Combe, C., *et al.* 2002. Depression as a predictor of mortality and hospitalization among hemodialysis patients in the United States and Europe. *Kidney International*, Vol. 62 : 199–207.
- Mancini, G.B., 2013. Minimum clinically important differences identified for commonly used depression rating scales. *Journal of Clinical Epidemiology* 66: 802-7.
- Mittal, S.K., Ahern, L., Flaster, E., Mittal, V.S., Maesaka, J.K., Fishbane, S., 2001. Self-assessed quality of life in peritoneal dialysis patients. *Am J Nephrol*: 215-20.
- Palmer, S.C., Vecchio, M., Craig, C.J. 2013. Association between depression and death in people with CKD: a meta-analysis of cohort studies. *American journal of kidney disease* 62: 493-504.
- Pederson, J.E., Warkentin, L.M., Majumdar, S.R., McAlister, F.A., 2016. Depressive Symptoms Are Associated With Higher Rates of Readmission or Mortality After Medical Hospitalization: A Systematic Review and Meta-analysis. *Journal of Hospital Medicine* :373–80.
- Perhimpunan Nefrologi Indonesia (PERNEFRI), 2011. Konsensus peritoneal dialisis pada penyakit ginjal kronik. PERNEFRI, Jakarta.

- Pengurus Pusat Perhimpunan Dokter Spesialis Kedokteran Jiwa Indonesia (PP PDSKJI), 2012. Pedoman Nasional Pelayanan Kedokteran Jiwa/Psikiatri (PNPK Jiw/Psikiatri), PP PDSKJI, Jakarta. : 63-72.
- Ramalingam, A. T., Narayan, T., Limbasia, R. D., Ushma, M., Dimple, J., . Patel Dimple, P. *et al.*, 2013. Quality of life, level of physical activity and depression in dialysis patients. *IOSR Journal of Dental and Medical Sciences (IOSR-JDMS)* : 46-50.
- Safaruddin, S., 2012. *Hubungan pola terapi , nilai ureum-kreatinin plasma dan hemoglobin dengan kualitas hidup pasien hemodialisis di RSUD Dr. Soedarso Pontianak* (Thesis). Universitas Indonesia, Jakarta.
- Saraha, S. M., Kanine, E., Wowiling, F., 2013. Hubungan Dukungan Keluarga dengan Depresi pada PAsien Ginjal Kronik di Ruang Hemodialisa BLU RSUP Prof. Dr. R D. Kandaou. *ejournal Keperawatan (e-Kp) Volume 1. Nomor 1* : 1-6.
- Smit, W., 2003. Peritoneal function and assessment of reference values using a 3.86% glucose solution. *Perit Dial Int* 23(5):440-449.
- Smit, W., 2004. Analysis of the prevalence and causes of ultrafiltration failure during long-term peritoneal dialysis: a cross-sectional study. *Perit Dial Int* 24(6):562-570.
- Suwitra, K., 2014. Penyakit ginjal kronik. In: Setiati, S., Alwi, I., Sudoyo, A.W., Simadibrata, M., Setiyohadi, B., Syam, A.F. (Ed): *Buku Ajar Ilmu Penyakit Dalam*, pp: 2159-65. Interna Publishing, Jakarta.
- Silva, M.T., Galvao, T.F., Martins, S.S., Pereira, M.G., 2014. Prevalence of depression morbidity among Brazilian adults: a systematic review and meta-analysis. *Revista Brasileira de Psiquiatria* :262–70.
- Sinnakirouchennan, R., Holley, J.L., 2011. Peritoneal Dialysis Versus Hemodialysis: Risks, Benefits, and Access Issues. *Advances in Chronic Kidney Disease Vol. 18*, No 6: 428-32.
- Sorayah, 2014. Uji Validitas Konstruk Beck Depression Inventory – II (BDI – II). *JP3I Vol. IV*: 29-46.
- Stack, A.G., 2002. Determinants of modality selection among incident US dialysis patients: results from a national study. *J Am Soc Nephrol.*:1279-87.
- Thomas, R., Acharya, S., Shukla, S., 2014. Prevalance of Depression among Patients with Chronic Kidney Disease. *IOSR Journal of Dental and Medical Sciences* :19-22.

- Weinhandl, E.D., Foley, R.N., Gilbertson, D.T., 2010. Propensity-matched mortality comparison of incident hemodialysis and peritoneal dialysis patients. *J Am Soc Nephrol* :499-506.
- Van Dij, C.M., Ledesma, S.G., Teitelbaum, I., 2005. Patient characteristics associated with defects of the peritoneal cavity boundary. *Perit Dial Int* 25(4):367-73.
- Yu, Z.L., Yeoh, L.Y., Seow, Y.Y., Luo, X.C., Griva, K., 2012. Evaluation of adherence and depression among patients on peritoneal dialysis. *Singapore Med J* ;57(7) :474-8