

DAFTAR PUSTAKA

- Baik, et al. 2001. Laryngeal Mask Insertion During Target Controlled Infusion of Propofol, In: *Journal of Clinical Anesthesia*, vol 13, p 175-81
- Bajwa, et al. 2010. Comparison of two drug combinations in TIVA: propofol-ketamine and propofol-fentanyl. *Saudi Journal of Anaesthesia*. www.saudija.org
- Budiarto, E. 2002, *Metodologi Penelitian Kedokteran*, Jakarta : Penerbit Buku Kedokteran EGC.
- Butterworth, J.F, Mackey, D.C. & Wasnick, J.D. (ed). 2013, *Morgan & Mikhail's Clinical Anesthesiology*, New York: McGraw Hill Education.
- Casati A, et al., 1999. The target plasma concentration of propofol required to place LMA vs COPA, in: *Anesthesi Analgesia*, vol.88, p 917-20
- Cook T dan Howes B. 2011. Supraglottic Airway Devices: Recent Advances, in: *Continuing Education in Anaesthesia Critical Care & Pain*, 11(2), p. 56-61
- Fan, et al., 1995. Propofol Concentration Monitoring in Plasma or Whole Blood by Gas Chromatography and High Performance Liquid Chromatograph, in: *Anesthesia Analgesia*, vol. 81, p 175-8
- Gopinath et al., 2015. Estimation of Effect-Site Concentration of Propofol for Laryngeal Mask Airway Insertion Using Fentanyl or Morphine as Adjuvant, in: *Indian Journal of Anaesthesia*
- Hagberg, C.A. 2013, *Benumof and Hagberg Airway Management*, 3rd ed. Elsevier Saunders
- Haynes, et al., 1992. Arterial Oxygen Saturation During Induction of Anaesthesia and Laryngeal Mask Insertion : Prospective Evaluation of Four Techniques, in: *British Journal of Anaesthesia*, vol. 68, p 519-22
- Hosseinzadeh H et al., 2013. Hemodynamic Changes Following Anesthesia Induction and LMA Insertion With Propofol, Etomidate and Propofol + Etomidate, in: *Journal of Cardiovascular and Thoracic Research*, 5(3), 109-112, doi: 10.5681/jcvtr.2013.023

- Jayaram et al, 2013. Comparison of Dexmedetomidine Combined with Propofol vs Fentanyl Combined with Propofol for Laryngeal Mask Insertion. *J Clin Sci Res* 2014; 3:p228-36. DOI: <http://dx.doi.org/10.15380/2277-5706.JCSR.13.032>
- Joo HS., Perks WJ. 2000, Sevoflurane Versus Propofol for Anesthetic Induction: A Meta-Analysis, in: *Anesthesia & Analgesia*, 91(1): 213-19
- Khan, P., Afridi, Y., 2008. Comparison Between Propofol and Thiopentone Sodium for Laryngeal Mask Airway Insertion in Day Case Surgery, in: *Journal of Postgraduate Medical Institute*, 22(3)
- Krishnappa & Kundra, 2011. Optimal Anaesthetic Depth for LMA Insertion, in: *Indian Journal of Anaesthesia*, vol. 55, p 504-7
- Laso LF et al., 2016. Manual Vs. Target Controlled Infusion Induction with Propofol: An Observational Study, in: *Columbian Journal of Anesthesiology*, 44(4), p 272-77.
- Leslie, K., Clavisi, O., Hargrove, J. (2008). *Target-controlled infusion versus manually-controlled infusion of propofol for general anaesthesia or sedation in adults (Review)*, The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.
- Macqueire, V. Cantraine, F. Schmartz, D. Et al. (2002). Target-Controlled Infusion Of Propofol Induction With Or Without Plasma Concentration Constraint In High-Risk“. In: *Acta Anaesthesiologica Scandinavica*, vol. 8, p 1010–1016
- Parasa, M. 2014, Ideal Induction Agent for LMA Insertion : a Comparative Study Between Thiopental Sodium and Propofol, in: *International Journal of Scientific Research*, 3(11), p. 7-8, DOI: 10.15373/22778179.
- Ramaswamy & Shaikh, 2015. Comparison of Dexmedetomidine-Propofol Versus Fentanyl-Propofol for Insertion of Laryngeal Mask, in: *Journal of Anaesthesiology Clinical Pharmacology*, vol. 31, p 217-220.
- Sastroasmoro S. & Ismael, S, 2014. *Dasar-dasar Metode Penelitian Klinis*. Jakarta: Sagung Seto.
- Sintavanuruk K., et al, 2010. Comparative Study of Effective Site Target Controlled Infusion with Standard Bolus Induction of Propofol for LMA Insertion, in: *Asian Biomedicine*, vol. 4, p 177-82.

- Stanski, D.R., Shafer, S.L. 2005. Measuring Depth of Anesthesia, in: *Miller's Anesthesia*, 6th ed., Elsevier Churchill Livingstone.
- Stoelting, R. K., et al, 2015, *Handbook of Pharmacology and Physiology in Anesthetic Practise*, 3rd ed, Pennsylvania: Lippincot-William & Wilkins.
- Struys, MM et al., 2016. The History of Target-Controlled Infusion, in: *Anesthesia & Analgesia*, vol. 122, p56-69
- Sugiarto, Adhrie., 2012, *Panduan Praktis Total Intravenous Anesthesia dan Target Controlled Infusion*, PP PERDATIN
- Surahman, 2016. *Metodologi penelitian*, Pusat Pendidikan Sumber Daya Manusia Kesehatan, Badan Pengembangan dan Pemberdayaan Sumber Daya Manusia Kesehatan, Kementerian Kesehatan Republik Indonesia.
- Taylor N, 1998. Requirements of Target-Controlled Infusion of Propofol to Insert the Laryngeal Mask Airway, in: *Anaesthesia Journal*, 52 (3), p. 222-226
- Teleflex, 2013. *Instruction for Use-LMA Supreme*, LMA The Laryngeal Mask Company Limited
- Vuyk J., Sitsen E., Reekers M., 2015. Intravenous Anesthetics in: *Miller's Anesthesia*, 8th Ed, Philadelphia: Elsevier, p 821-832
- White, F. P., Eng, M. R., 2013. Intravenous Anesthesia in: Barash, et al (ed), *Clinical Anesthesia*, 7th ed, Philadelphia Lippincott Wiliams & Wilkins.
- Wong CM., et al, 2007. Fentanyl Dose-Response Curves when Inserting LMA Classic, in: *Anaesthesia*, vol. 62, p 654-60
- Zaballos M., et al, 2015. Effect-Site Concentration of Propofol Required for LMA-Supreme Insertion with and without Remifentanil: a Randomized Controlled Trial, in: *BMC Anesthesiology*. www.biomedcentral.com