

DAFTAR PUSTAKA

- Antinori, A., Arendt, G., Becker, J.T., Brew, B.J., Byrd, D.A., Cherner, M. *et al.* (2008). Updated research nosology for HIV-associated neurocognitive disorders. *Neurology*. 69: 1789-1799.
- Arora, Sunil; De Sousa, A. (2013). Plasma Viral Load, CD4 Count And Hiv Associated Dementia Sunil. *National Journal Of Medical Research Original* pp. 13–15.
- Balaini, N., Sharma, A., Sharma, A., Sharma, A. (2017). HIV associated neurocognitive dysfunction and its association with CD4 count in HIV positive patients-a hospital based study. *International Journal of Research in Medical Sciences*. Oct;5(10):4259-4266.
- Bartlett, JG, Gallant, JE. (2005). Medical management of HIV infection. *Baltimore: Johns Hopkins University Division of Infectious Diseases*; Available: www.hopkinsaids.edu/mmhiv/order.html
- Beyrer, C., Baral, S. D., van Griensven, F., Goodreau, S. M., Chariyalertsak, S., Wirtz, A. L. *et al.* (2012). Global epidemiology of HIV infection in men who have sex with men. *Lancet (London, England)*, 380(9839), 367-77.
- Bhakta, M. (2001). Anemia, Hematologi klinik ringkas. UPT Penerbit Universitas Udayana, p.12-13.
- Bottiggi, K.A., Chang, J.J., Schmitt, F.A., Avison, M.J., Mootoor, Y., Nath, A. *et al.* (2007). The HIV Dementia Scale: predictive power in mild dementia and HAART. *Journal of the neurological sciences*. 260:11-15.
- Brasileira, S., Troncoso, F., Conterno, L. (2015). Major Article Prevalence of neurocognitive disorders and depression in a Brazilian HIV population. 48(4), pp. 390–398.
- Budiarto, Eko. (2004). Metodologi Penelitian Kedokteran: Sebuah Pengantar. Jakarta: Penerbit Buku Kedokteran EGC.
- Chalermchai, T., Valcour, V., Sithinamsuwan, P., Pinyakorn, S. (2013). Trial making test A improves performance characteristics of the international HIV-dementia scale to identify symptomatic HAND. *Journal of neurovirology*. 99-106.
- Dahlan, M.S. (2014). *Statistik Untuk Kedokteran Dan Kesehatan : Deskriptif, Bivariat, Dan Multivariat Dilengkapi Aplikasi Menggunakan SPSS, 6th ed, 1.* Jakarta: Epidemiologi Indonesia.
- Dhadke, S.V., Dhadke, V.N., Mahajan, N.P., Korade, M.B. (2014) Clinical profile of neurological complications in HIV- reactive patients and their relation with CD4. *Int J Med Biomed Res* 2014;3(2):91-100.

- Ditjen PP & PL Kemenkes RI and Kementrian Kesehatan RI. (2014). *Data Statistik HIV di Indonesia 2014*. Kemenkes RI, pp. 1–3.
- Ditjen PP&PL Kemenkes RI. (2016). *Statistik kasus HIV/AIDS di Indonesia*.
- Duarsa, N. W. (2009). *Infeksi Menular Seksual*. 4th ed. Jakarta: Balai Penerbit FKUI, pp. 146-53.
- Ellis, R.J., Badiee, J., Vaida, F., Letendre, S., Heaton, R.K., Clifford, D. *et al.* (2011) CD4 nadir is a predictor of HIV neurocognitive impairment in the era of combination antiretroviral therapy. *AIDS*. 2011 September 10; 25(14)
- Estiasari, R., Imran, J., Diatri & Prawiroharjo, Pukovisa & Price, Patricia. (2015). Cognitive impairment among Indonesia HIV naïve patients. *Neurology Asia*. 20. 155-160.
- Fernandes, L. G. (2016). Hiv And Dementia : Prevalence And Risk Factors. *Rev Bras Promoç Saúde, Fortaleza*, 29(2): 212-218, abr./jun., 2016
- Fitri, F. I., Rambe, A. S., Fitri, A. (2018). Correlation between Lymphocyte CD4 Count , Treatment Duration , Opportunistic Infection and Cognitive Function in Human Immunodeficiency Virus-Acquired Immunodeficiency Syndrome (HIV-AIDS) Patients. *Open Access Maced J Med Sci*. 2018 Apr 15; 6(4):643-647.
- Ganasean, K.A., Fincham, D., Smit, J., Seedat, S., Stein, D. (2008). Utility of the HIV Dementia Scale (HDS) in identifying HIV dementia in a South African sample. *Journal of the neurological sciences*. 269:62-64.
- Gannon, P., Khan, M. Z., & Kolson, D. L. (2011). Current understanding of HIV-associated neurocognitive disorders pathogenesis. Current opinion in neurology, 24(3), 275-83.
- Ghafouri, M., Amini, S., Khalili, K., Sawaya, B.E. (2006). HIV-1 associated dementia: symptoms and causes. *Retrovirology*. 3:1-11.
- Grund, B., Wright, E.J., Brew, B.J. (2013). Improved neurocognitive test performance in both arms of the SMART study: impact of practice effect. *J. Neurovirol* 19:383–392.
- Haddow, L.J., Floyd, S., Copas, A., Gilson, R.J.C. (2013). A systematic review of the screening accuracy of the HIV Dementia Scale and International HIV dementia Scale. *Plos ONE*. 8(4):e61826.
- Hasbun, R., Eraso, J., Ramireddy, S., Wainwright, D., Salazar, L., Grimes, R. *et al.* (2012). Screening for neurocognitive impairment in HIV individuals: the utility of the montreal cognitive assessment test. *Journal AIDS clinic Res*. 3:10.
- Heaton, R.K., Franklin, D.R., Ellis, R.J., McCutchan, J.A., Letendre, S.L., Leblanc, S. *et al.* (2011). HIV-associated neurocognitive disorders before and

during the era of combination antiretroviral therapy: differences in rates, nature, and predictors. *J Neurovirol.*17:3-16.

- Hinkin, C.H., Castellon, S.A., Atkinson, J.H., Goodkin, K. (2001) Neuropsychiatric aspects of HIV infection among older adults. *Journal of clinical epidemiology.* 119:44-52.
- Hu, X., Zhou, Y., Long, J., Feng, Q., Wang, R., Su, L. *et al.* (2012) Diagnostic accuracy of the International HIV Dementia Scale and HIV Dementia Scale: A meta-analysis. *Experimental and therapeutic medicine.* 4:665-668.
- Johnson-Wimbley, T. D., Graham, D. Y. (2011). Diagnosis and management of iron deficiency anemia in the 21st century. *Therapeutic Advances in Gastroenterology,* 4(3), 177–184. <http://doi.org/10.1177/1756283X11398736>
- Joska, J.A., Gouse, H., Paul, R.H., Stein, D.J., Flisher, A.J. (2010). Does highly active antiretroviral therapy improve neurocognitive function? A systematic review. *Journal of neurovirology.* 1-14.
- Kalinowska, S., Trzeźniowska-Drukała, B., Samochowiec, J., (2013) HIV-associated neurocognitive disorders. *Psychiatr Pol.* May-Jun;47(3):453-63.
- Kallianpur, A., Wang, Q., Peilin, H., Zhao, Z., Letendre, S. (2016). Anemia and Red Blood Cell Indices Predict HIV-Associated Neurocognitive Impairment in the Highly Active Antiretroviral Therapy Era. *The Journal of Infectious Diseases,* 213(7), pp. 1065–1073. Available at: <http://dx.doi.org/10.1093/infdis/jiv754>.
- Kanmogne, GD., Kuate, CT., Cysique, LA. (2010). HIV-associated neurocognitive disorders in sub-Saharan Africa: a pilot study in Cameroon. *BMC Neurol.* 2010;10:60.
- Kaul, M., Garden, G. A., Lipton, S. A. (2001). Pathways to neuronal injury and apoptosis in HIV-associated dementia. *Nature,* 410, 988. Retrieved from <http://dx.doi.org/10.1038/35073667>
- Kaul, M., Zheng, J., Okamoto, S., Gendelman, H., Lipton, S. (2005). HIV-1 infection and AIDS: consequences for the central nervous system, Cell Death And Differentiation. *Nature Publishing Group,* 12, p. 878. Available at: <https://doi.org/10.1038/sj.cdd.4401623>.
- Kinuthia, RN., Gakinya, BN., Thigiti, JM. (2016). Relationship between HIV stage and psychomotor speed neurocognitive score at a Kenyan sub-county hospital. *Afr J Prim Health Care Fam Med.* 2016;8(1):e1-8.
- Kolegium Neurologi Indonesia. (2008). *Neurobehaviour bagian II: gangguan gerak.* Jakarta : PERDOSSI.



- Kresno, S.B. (2001). *Uji serologi infeksi HIV. Immunologi: diagnosis dan prosedur laboratorium*. Jakarta: Fakultas Kedokteran Universitas Indonesia, pp.369-377.
- Langford, S., Ananworanich, J., Cooper, D. (2007). Predictors of disease progression in HIV infection: a review. *AIDS Res Ther.* 2007; 4: 11.
- Lawler, K., Mosepele, M., Ratcliffe, S., Seloilwe, E., Steele, K., Nthobatsang, R. *et al.* (2010). Neurocognitive impairment among HIV-positive individuals in Botswana: a pilot study. *Journal of the international AIDS society.* 13:15.
- Levinson, P., Choi, RY., Cole, AL., Hirbod, T., Rhedin, S., Payne, B. (2012) HIV-Neutralizing Activity of Cationic Polypeptides in Cervicovaginal Secretions of Women in HIV-Serodiscordant Relationships. *PLoS ONE* 7(2): e31996. Available from <https://doi.org/10.1371/journal.pone.0031996>.
- Lindl, K., David, R., Marks, Dennis, L., Kolson, and, Sciutto J, Kelly L. (2010) HIV-Associated Neurocognitive Disorder: Pathogenesis and Therapeutic Opportunities. *J Neuroimmune Pharmacol.* 5:294-309.
- Maiga, Y., Diakite, S., Cissoko, Y., Diallo, F., Kaïoulou, HA. (2014). Neuropathic Pain in HIV / AIDS Patients on Antiretroviral Therapy and Followed as Outpatients in Bamako, Mali. *J Pain Relief* S3: 004.
- Maki PM, Rubin LH, Valcour V, Martin E, Crystal H, Young M. *et al.* (2014) Cognitive function in women with HIV: findings from the Women's Interagency HIV Study. *American Academy of Neurology.*
- Mamidi A., DeSimone JA., Pomerantz RJ. (2002) Central nervous system infections in individuals with HIV-1 infection. *J Neurovirol.* 8(3):158-67.
- Marcotte, T., Deutsch, R., Mccutchan, J., Moore, D., Letendre, S., Ellis, R. *et al.* (2003). Prediction of Incident Neurocognitive Impairment by Plasma HIV RNA and CD4 Levels Early After HIV Seroconversion. *Archives of neurology.* 60. 1406-12. 10.1001/archneur.60.10.1406.
- McArthur, J.C., Hoover, D.R., Bacellar, H., Miller, E.N., Cohen, B.A., Becker, J.T. *et al.* (1993) Dementia in AIDS patients: incidence and risk factors. *Neurology.* 43:2245-2252.
- McArthur, J.C., McClernon, D., Cronin, M., Nance-Sproson, T., Saah, A., Lanier, R., (1997) Relationship Between Human Immunodeficiency Virus-Associated Dementia and Viral Load in Cerebrospinal Fluid and Brain. *Neurology.* 42:689-698
- McCombe, JA., Vivithanaporn, P., Gill, MJ., Power, C., (2013) Predictors of symptomatic HIV-associated neurocognitive disorders in universal health care. *HIV Med.* 14(2):99-107.

- McGuire, J.L., Barrett, J.S., Vezina, H.E., Spitsin, S., Douglas, S.D., (2014). Adjuvant therapies for HIV-associated neurocognitive disorders. *Philadelphia: ANNALS of clinical and translational neurology*.
- Muñoz-Moreno JA, Pérez-Álvarez N, Muñoz-Murillo A, Prats A, Garolera M, Jurado MÀ, et al. (2014) Classification Models for Neurocognitive Impairment in HIV Infection Based on Demographic and Clinical Variables. *PLoS ONE* 9(9): e107625. <https://doi.org/10.1371/journal.pone.0107625>.
- Nakku, J., Kinyanda, E., Hoskins, S. (2013) Prevalence and factors associated with probable HIV dementia in an African population: a cross-sectional study of an HIV/AIDS clinic population. *BMC Psychiatry*. 13:126
- National Institute of Health. (2014) *The eighth report of the Joint National Committee on prevention, detection, evaluation, and treatment of high blood pressure*. United State: Department of Health and Human Services.
- Njamnshi, A.K., Djientcheu, V.P., Fonsah, J.Y., Yepnjio, F.N., Njamnshi, D.M., Muna, W.F., (2008) The international HIV Dementia Scale is a useful screening tool for HIV-associated dementia/cognitive impairment in HIV-infected adults in Yaounde-Cameroon. *Journal acquir immune defic syndr*. 49:393-397.
- O. Olajumoke Oshinaike, A. Akinsegun Akinbami, O. Oluwadamilola Ojo, I. Frank Ojini, U. Njideka Okubadejo, and A. Mustapha Danesi. (2012). Comparison of the Minimental State Examination Scale and the International HIV Dementia Scale in Assessing Cognitive Function in Nigerian HIV Patients on Antiretroviral Therapy. *AIDS Research and Treatment*, vol. 2012, Article ID 581531, 6 pages, 2012.
- Ogunrin, A. O., Odiase, F. E. and Ogunniyi, A. (2007) Reaction time in patients with HIV / AIDS and correlation with CD4 count : a case — control study. *Trans R Soc Trop Med Hyg* . doi: 10.1016/j.trstmh.2006.10.002.
- Oshinaike, O., Akinbami, AA., Ojo, O., Ojini, IF., Okubadejo, UN., Danesi, AM., (2012). Clinical Study Comparison of the Minimental State Examination Scale and the International HIV Dementia Scale in Assessing Cognitive Function in Nigerian HIV Patients on Antiretroviral Therapy. *AIDS Res Treat*. doi: 10.1155/2012/581531.
- Pantaleo, G., Graziosi, C., Demarest, JF., Butini, L., Montroni, M., Fox, CH. *et al*. (1993) HIV infection is active and progressive in lymphoid tissue during the clinically latent stage of disease. *Nature*. 362(6418):355-8.
- Pascal, M., Gaspard, T., Philomène, N., Emmanuel, Y., Avilah, AW., Hortense, H. (2016). Determinants of Neurocognitive Impairment in HIV in a Cohort of Patients on Antiretroviral Therapy Followed in Bangui (Central African Republic). *Neuroscience & Medicine*. 07. 1-9. 10.4236/nm.2016.71001.

- Perdossi. (2015). *Diagnosis dan penatalaksanaan demensia*. Kolegium Neurologi Indonesia.
- Perhimpunan Dokter Spesialis Saraf Indonesia (PERDOSSI). (2006) *Konsensus nasional penanganan trauma kaputid dan trauma spinal*. Jakarta: PERDOSSI.
- Perkumpulan Endokrinologi Indonesia (PERKENI). (2015). *Konsensus Pengelolaan dan Pencegahan Diabetes melitus tipe 2 di Indonesia*. Jakarta : PERKENI.
- Pinheiro, C., Mattos Souza, L., Motta, J., Kelbert, E., Martins, C. de S. R., Souza, M. (2016). Aging, neurocognitive impairment and adherence to antiretroviral therapy in human immunodeficiency virus-infected individuals. *The Brazilian Journal of Infectious Diseases*, 20(6), 599–604. <https://doi.org/https://doi.org/10.1016/j.bjid.2016.09.006>
- Power, C., Selnes O.A., Grim, J.A., McArthur, J.C. (1995) HIV Dementia Scale: a rapid screening test. *J Acquir Immune Defic Syndr Hum Retrovirol*. 8(3):273-278
- Ragin, A. B., D'Souza, G., Reynolds, S., Miller, E., Sacktor, N., Selnes, O. *et al.* (2011). Platelet decline as a predictor of brain injury in HIV infection. *Journal of neurovirology*, 17(5), 487-95.
- Resita, Satiti, S., Nuradyo, D. (2011). *Korelasi Angka CD4 dengan Jumlah Refleks Primitif pada penderita Human Immunodeficiency Virus (HIV)*. Yogyakarta: Universitas Gadjah Mada.
- Rowi, M. (2011). *Infeksi Oportunistik sistem saraf pusat pada pasien AIDS*. Jakarta: Fakultas Kedokteran Universitas Trisakti.
- Sacktor, N., Nakasujja, N., Skolasky, R., Robertson, K., Wong, M. (2006). Antiretroviral therapy improves cognitive impairment in HIV+ individuals in sub-Saharan Africa. *Neurology*. 67:311-314.
- Sari, D. K. (2007). *Brain cancer*. Yogyakarta : Fakultas Kedokteran Universitas Islam Indonesia.
- Sastroasmoro, Sudigdo, Ismael, Sofyan (2008). *Dasar-dasar Metodologi Penelitian Klinis*. Edisi ke-3. Jakarta: Sagung Seto.
- Satriawan, DN., Widyadharma, E., Adnyana, MO (2010). *Uji Reliabilitas Skala Demensia HIV Internasional pada penderita HIV di RSUP Sanglah*. Udayana.
- Schreibman, T., Friedland, G. (2004). Use of Total Lymphocyte Count for Monitoring Response to Antiretroviral Therapy. *Clinical Infectious Diseases* ; Volume 38, Issue 2, 15 January 2004, Pages 257–262, <https://doi.org/10.1086/380792>

- Setiawan, N.K., Satiti, K., Nuradyo, D. (2011). *Risiko prevalensi indeks massa tubuh rendah dengan gangguan kognitif pada penderita human immunodeficiency virus*. Tesis, Yogyakarta: Bagian/SMF neurologi FK UGM/RSUP Dr. Sardjito.
- Setyopranoto, I., (2012) *Odem otak pada pasien stroke iskemik akut*. Yogyakarta : Badan penerbit Fakultas Kedokteran Universitas Gadjah Mada.
- Shapshak, P., Minagar, A., Duran, E., Ziegler, F., Davis, W. (2005) Gene expression in HIV-associated dementia. *Totowa, NJ: Humana Press Inc*.
- Steckley, J.L. (2005). Age, anemia, opportunistic infection, and low CD4 counts increased the risk of HIV dementia. *The UWO evidence based neurology group*.
- Stirrup, OT., Copas, AJ., Phillips, AN., Gill, M., Geskus, R., Touloumi. *et al.* (2018). Predictors of CD4 cell recovery following initiation of antiretroviral therapy among HIV-1 positive patients with well-estimated dates of seroconversion. *HIV Medicine* pp. 184–194. doi: 10.1111/hiv.12567.
- Sumonu, T. A., Imarhiagbe, F., Owolabi, L. F., Ogunrin, O. A., Komolafe, M. A., & Ilesanmi, O. S. (2017). Cognitive functions in newly diagnosed patients with HIV infection in a tertiary health facility: Assessment using community screening interview for dementia. *eNeurologicalSci*, 9, 8-13. doi:10.1016/j.ensci.2017.10.001
- The IeDEA and COHERE Cohort Collaborations. (2018). *Global Trends in CD4 Cell Count at the Start of Antiretroviral Therapy : Collaborative Study of Treatment Programs*. 66, pp. 893–903. doi: 10.1093/cid/cix915.
- Tozzi, V., Balestra, P., Libertone, R., Antinori, A. (2010). Cognitive function in treated HIV patients. *Italy: Dovepress*.
- Tumbelaka, A.R., Riono, P., Sastroasmoro, S., Wirdjodiarjo, M., Pudjiastuti, P. and Firman, K. (2002). *Pemilihan uji hipotesis*. Dalam: Sastroasmoro, S., & Ismael, S, ed. 2011. *Dasar-dasar metodologi penelitian*, ed. 2. Jakarta: CV. Sagung Seto, pp.240-258.
- Tyagi, M., Karn, J. (2007). CBF-1 promotes transcriptional silencing during the establishment of HIV-1 latency. *The EMBO Journal*, 26(24), 4985–4995. <http://doi.org/10.1038/sj.emboj.7601928>
- Van Harten, B., Courant, M.N., Scheltens, P., Weinstein, H.C., (2004) Validation of the HIV dementia scale in an elderly cohort of patients with subcortical cognitive impairment caused by subcortical ischaemic vascular disease or a normal pressure hydrocephalus. *Dement Geriatr cogn disord*. 18(1):109-114.
- Vassallo, M., Fabre, R.M., Durant, J.T., Lebrun-Frénay, C., Joly, H., Ticchioni, M. *et al.* (2016). A decreasing CD4/CD8 ratio over time and lower CSF-

penetrating antiretroviral regimens are associated with a higher risk of neurocognitive deterioration, independently of viral replication. *Journal of NeuroVirology*, 23, 216-225.

- Vassallo, Matteo & Durant, Jaquisha & Lebrun-Frenay, Christine & Fabre, R & Ticchioni. (2015). Virologically suppressed patients with asymptomatic and symptomatic HIV-associated neurocognitive disorders do not display the same pattern of immune activation: The CD4:CD8 ratio and neurocognitive disorders. *HIV Medicine*. 16. 10.1111/hiv.12246.
- Vo, Q. T., Cox, C., Li, X., Jacobson, L. P., McKaig, R., Sacktor, N. *et al.* (2013). Neuropsychological Test Performance Before and After HIV-1 Seroconversion: The Multicenter AIDS Cohort Study. *Journal of Neurovirology*, 19(1), 24–31. <http://doi.org/10.1007/s13365-012-0136-8>
- Wachtman, LM., Skolasky, RL., Tarwater, PM. (2007). Platelet Decline: An Avenue for Investigation Into the Pathogenesis of Human Immunodeficiency Virus–Associated Dementia. *Arch Neurol*. 2007;64(9):1264–1272. doi:10.1001/archneur.64.9.1264
- Waldrop-Valverde, D., Nehra, R., Sharma, S., Malik, A., Jones, D., Kumar, A.M. *et al.* (2010) Education effects on the International HIV dementia scale. *Journal of neurovirology*. 16:264-267.
- Wang. (2013). High prevalence of HIV-associated neurocognitive disorder in HIV-infected patients with a baseline CD4 count \leq 350 cells/ μ L in Shanghai, China. *BioScience Trends*. 10.5582/bst.2013.v7.6.284.
- Wright, Ej., Grund, B., Cysique, L., Robertson, K., Brew, B., Collins. *et al.* (2015). Factors associated with neurocognitive test performance at baseline: a substudy of the INSIGHT Strategic Timing of AntiRetroviral Treatment (START) trial. *HIV medicine*. 16 Suppl S1. 97-108. 10.1111/hiv.12238.
- Zipursky, A.R., Gogolishvili, D., Rueda, S., Brunetta, J., Rueda, S., Carvalhal, A. *et al.* (2013) Evaluation of brief screening tools for neurocognitive impairment in HIV/AIDS: a systematic review of the literature. *AIDS*. 27:2385-2401.