

DAFTAR PUSTAKA

- Amiri-simkooei, A., Snellen, M. & Simons, D.G. 2009. Riverbed sediment classification using multi-beam echo-sounder backscatter data. *J. Acoust. Soc. Am.*, 126(4): 1724–1738.
- Anderson, J.T., Van Holliday, D., Kloser, R., Reid, D.G. & Simard, Y. 2008. Acoustic seabed classification: Current practice and future directions. *ICES Journal of Marine Science*, 65(6): 1004–1011.
- Biot, M.A. 1956. Theory of propagation of elastic waves in a fluid-saturated porous solid. I. Low-frequency range. *The Journal of the Acoustical Society of America*, 28(2): 168–178.
- Biot, M.A. 1962. Generalized theory of acoustic propagation in porous dissipative media. *The Journal of the Acoustic Society of America*, 34(5): 1254–1264.
- Blott, S.J. & Pye, K. 2001. GRADISTAT : A Grain Size Distribution and Statistics Package for The Analysis of Unconsolidated Sediments. *Earth Surface Processes and Landforms*, 26: 1237–1248.
- Brown, C., Schmidt, V., Malik, M. & Le Bouffant, N. 2015. Backscatter measurement by bathymetric echo sounders. X. Lurton & G. Lamarche, ed., *Backscatter measurements by seafloor-mapping sonars - Guidelines and Recommendations*. hal.79–105. Tersedia di <http://geohab.org/wp-content/uploads/2014/05/BSWGREPORT-MAY2015.pdf>.
- CARIS 2013. *CARIS HIPS and SIPS 8.1 User Guide*. Tersedia di https://www.cambridge.org/core/product/identifier/CBO9780511841538A006/type/book_part.
- Cervenka, P.S. & de Moustier, C. 1993. Sidescan Sonar image processing techniques. *IEEE Journal of Oceanic Engineering*, 18(2): 108–122.
- Clarke, J.E.H., Danforth, B.W. & Valentine, P. 1997. Areal Seabed Classification using Backscatter Angular Response at 95kHz Areal Seabed Classification using Backscatter Angular Response at 95kHz.
- Dufek, T. 2012. *Backscatter Analysis of Multibeam Sonar Data in the Area of the Valdivia Fracture Zone using Geocoder in CARIS HIPS & SIPS and IVS3D Fledermaus*. HafenCity Universität Hamburg.
- De Falco, G., Tonielli, R., Di Martino, G., Innangi, S., Simeone, S. & Parnum, I.M. 2010. Relationships between multibeam backscatter, sediment grain size and *Posidonia oceanica* seagrass distribution. *Continental Shelf Research*, 30(18): 1941–1950. Tersedia di <http://dx.doi.org/10.1016/j.csr.2010.09.006>.
- Fonseca, L., Brown, C., Calder, B., Mayer, L. & Rzhannov, Y. 2009. Angular range analysis of acoustic themes from Stanton Banks Ireland: A link between visual interpretation and multibeam echosounder angular signatures. *Applied Acoustics*, 70(10): 1298–1304. Tersedia di <http://dx.doi.org/10.1016/j.apacoust.2008.09.008>.

- Fonseca, L. & Calder, B. 2005. Geocoder: An efficient backscatter map constructor. *Proceedings of the U.S. Hydrographic Conference 2005*. San Diego, California, Amerika Serikat.
- Fonseca, L. & Calder, B. 2007. Clustering Acoustic Backscatter in the Angular Response Space. *Proceedings US Hydro 2007*, 1–12.
- Fonseca, L. & Mayer, L. 2007. Remote estimation of surficial seafloor properties through the application angular range analysis to multibeam sonar data. *Marine Geophysical Researches*, 119–126.
- Hammerstad, E., Pohner, F., Parthoit, F. & Bennet, J. 1991. Field testing of a new deep water multibeam echo sounder. *Proceedings IEEE*, 2: 743–749.
- Hellequin, L., Boucher, J. & Lurton, X. 2003. Processing of High-Frequency Multibeam Echo Sounder Data for Seafloor Characterization. *IEEE Journal of Oceanic Engineering*, 28(1): 78–89.
- Ivakin, A.N. 1998. A unified approach to volume and roughness scattering. *The Journal of the Acoustical Society of America*, 103(2): 827–837.
- Jackson, D., Winebrenner, D. & Ishimaru, I. 1986. Application of the composite roughness model to high-frequency bottom backscattering. *Journal of the Acoustic Society of America*, 79(5): 1410–1422.
- Jackson, D.R. & Richardson, M.D. 2007. *High-Frequency Seafloor Acoustics*. New York, Amerika Serikat: Springer Science+Business Media.
- de Jong, C.D., Lachapelle, G., Skone, S. & Elema, I.A. 2003. *Hydrography*. 2nd ed. *Nature*. Delft, Netherlands.
- Kowerko, D. 2015. *Numerical calculation of the relative overlap area of two normal distributions*. Tersedia di https://www.mathworks.com/matlabcentral/fileexchange/49823-calc_overlap_twonormal-s1-s2-mu1-mu2-xstart-xend-xinterval.
- Krumbein, W.C. 1934. Size frequency distribution of sediments. *Journal of Sedimentary Research*, 4(2): 65–77.
- Lucieer, V., Roche, M., Degrendele, K., Malik, M., Dolan, M. & Lamarche, G. 2017. User expectations for multibeam echo sounders backscatter strength data-looking back into the future. *Marine Geophysical Research*, 0(0): 0.
- Lurton, X. 2010. *An Introduction to Underwater Acoustics - Principles and Applications*. 2 ed. Berlin Heidelberg: Springer-Verlag.
- Masetti, G., Sacile, G. & Trucco, A. 2011. Remote characterization of seafloor adjacent to shipwrecks using mosaicking and analysis of backscatter response. *Italian Journal of Remote Sensing*, 43(2): 77–92.
- De Moustier, C. & Alexandrou, D. 1991. Angular dependence of 12-kHz seafloor acoustic backscatter. *Journal of the Acoustical Society of America*, 90(1): 522–531.
- Pace, N.G. & Gao, H. 1988. Swathe Seabed Classification. *IEEE Journal of Oceanic Engineering*, 13(2): 83–90.

- Penrose, J.D., Siwabessy, P.J.W., Gavrilov, A., Parnum, I., Hamilton, L.J., Bickers, A., Brooke, B., Ryan, D.A. & Kennedy, P. 2005. *Acoustic Techniques for Seabed Classification*. Cooperative Research Centre for Coastal Zone Estuary and Waterway Management.
- Snellen, M., Siemes, K. & Simons, D.G. 2011. Model-based sediment classification using single-beam echosounder signals. *J. Acoust. Soc. Am.*, 129(5): 2878–2888.
- Weber, T. & Lurton, X. 2015. Background and fundamentals. X. Lurton & G. Lamarche, ed., *Backscatter measurements by seafloor-mapping sonars - Guidelines and Recommendations*. hal.25–51. Tersedia di <http://geohab.org/wp-content/uploads/2014/05/BSWGREPORT-MAY2015.pdf>.
- Van Zuidam, R.A. 1983. *Guide to geomorphological aerial photographic interpretation and mapping*. Enschede, the Netherlands: ITC.