

DAFTAR PUSTAKA

- Aeberli, I., Gut-Knabenhans, M., Kusche-Ammann, R. S., Molinari, L., Zimmermann, M. B. (2013). A composite score combining waist circumference and body mass index more accurately predicts body fat percentage in 6 to 13year-old children. *European Journal of Nutrition*, 52(1), 247-253.
- Al-Bachir, M., & Ahmad, H. (2016). Validity of using waist and hip circumference measurements to determine body composition of young syrian men. *Journal of Biosocial Science*, 48(5), 647–657.
- Almatsier, S. (2009). *Prinsip Dasar Ilmu Gizi*. Jakarta: Gramedia.
- Badan Penelitian dan Pengembangan Kesehatan. (2013). *Riset Kesehatan Dasar; RISKESDAS*. Jakarta: Balitbang Kemenkes RI
- Bantas, K., Koesnanto, H., & Moelyono, B. (2013). Ukuran lingkar pinggang optimal untuk identifikasi sindrom metabolik pada populasi perkotaan di indonesia. *Jurnal Kesehatan Masyarakat Nasional*, 7(6), 284-288.
- Barba, C. et al., 2004. Appropriate body-mass index for Asian populations and its implications for policy and intervention strategies. *Lancet*, 363(9403), 157–163.
- Bassali, R., Waller, J., Gower, B., Allison, J. and Davis, C. (2010). Utility of waist circumference percentile for risk evaluation in obese children. *International Journal of Pediatric Obesity*, 5(1), 97-101.
- Batubara, J. R. L. (2010). Perkembangan remaja. *Sari Pediatri*, 12(1), 21-29.
- Biro, F. M., & Wien, M. (2010). Childhood obesity and adult morbidities. *The American Journal of Clinical Nutrition*, 91(1), 1499–1505.
- Bouchard, C. (1997). Genetic determinants of regional fat distribution. *Human Reproduction (Oxford, England)*, 12(1), 1–5.
- Brambilla, P., Bedogni, G., Moreno, L. A., Goran, M. I., Gutin, B., Fox, K. R., Pietrobelli, A. (2006). Crossvalidation of anthropometry against magnetic resonance imaging for the assessment of visceral and subcutaneous adipose tissue in children. *International Journal of Obesity*, 30(1), 23–30.
- Bray, A. G. (1976). *The obese patient*. Philadelphia: W. B. Saunders.
- Briefel, R., McDowell, M., Alaimo, K., Caughman, C., Bischof, A., Carroll, M., & Johnson, C. L. Total energy of the US population: the third National Health and Nutrition Examination Survey, 1988±1991. *The American Journal of Clinical Nutrition*, 62(5): 1072-1080.
- Cameron, N., Jones, L. L., Griffiths, P. L., Norris, S. A., & Pettifor, J. M. (2010). How well do waist circumference and body mass index reflect body composition in pre-pubertal children? *European Journal of Clinical Nutrition*, 63(9): 1065–1070.

- Centers of disease control. (2011). Body mass index: Considerations for practitioners. Retrieved from <http://scholar.google.com/scholar?hl=en&btnG=Search&q=intitle:Body+Mass+Index+:+Considerations+for+Practitioners#3%5Cnhttp://scholar.google.com/scholar?hl=en&btnG=Search&q=intitle:Body+mass+index+:+Considerations+for+practitioners#3>
- Centres for Disease Control and Prevention. (2007). Anthropometry procedures manual. *National Health and Nutrition Examination Survey (NHANES)*, 102. Retrieved from https://www.cdc.gov/nchs/data/nhanes/nhanes_07_08/manual_an.pdf%0Ahttp://www.cdc.gov/nchs/data/nhanes/nhanes_07_08/manual_an.pdf
- Chung, S. (2015). Body mass index and body composition scaling to height in children and adolescent. *Annals of Pediatric Endocrinology & Metabolism*, 20(3), 125-129.
- Daniels, S. R. (2009). The use of BMI in the clinical setting, *Pediatrics*, 124(1), 35-41.
- Deshmukh-Taskar, P., Nicklas, T. A., Morales, M., Yang, S. J., Zakeri, I., & Berenson, G. S. (2006). Tracking of overweight status from childhood to young adulthood: the Bogalusa Heart Study. *European Journal of Clinical Nutrition*, 60(1), 48-57.
- Druet, C., Baltakse, V., Chevenne, D., Dorgeret, S., Zaccaria, I., Wang, Y., & Levy-Marchal, C. (2008). Independent effect of visceral adipose tissue on metabolic syndrome in obese adolescents. *Hormone Research*, 70(1), 22-28.
- Duarte, M. O., Ruelas, Y. F., López-Alcaraz, F., Toro-Equihua, M. D., & Sánchez-Ramírez, C. A. (2014). Correlation between percentage of body fat measured by the Slaughter equation and bio impedance analysis technique in Mexican schoolchildren. *Nutricion Hospitalaria*, 29(1), 88-93.
- Duren, D. L., Sherwood, R. J., Czerwinski, S. A., Lee, M., Choh, A. C., Roger M. Siervogel, R. M., & Chumlea, W.C. (2008). Body composition methods: comparisons and interpretation. *Journal of Diabetes Science and Technology*, 2(6), 1139-1146.
- Eisenmann, J. (2005). Waist circumference percentiles for 7- to 15-year-old Australian children. *Acta Paediatrica*, 94(9), 1182-1185.
- Ellis, K. (2001). Selected body composition methods can be used in field studies. *The Journal of Nutrition*, 131(5), 1589-1595.
- Eissa, M. A., Dai, S., Mihalopoulos, N. L., Day, R. S., Harrist, R. B., Labarthe, D. R. (2009). Trajectories of fat mass index, fat free-mass index, and waist circumference in children project heartbeat! *American Journal of Preventive Medicine*, 37(1), 34-39.
- Fredriksen, P. E. R. M., Skår, A., & Mamen, A. (2018). Waist circumference in 6

– 12-year-old children : The Health Oriented Pedagogical Project (HOPP),
Scandinavian Journal of Public Health, 46(21), 12–20.

Fredriks, A. M., Van, B. S., Fekkes, M., Verloove-Vanhorick, S. P., & Wit, J.M. (2005). Are age references for waist circumference, hip circumference and waist-hip ratio in Dutch children useful in clinical practice? *European Journal of Clinical Nutrition*, 164(4), 216-222.

Freedland, E. S. (2004). Role of a critical visceral adipose tissue threshold (CVATT) in metabolic syndrome: Implications for controlling dietary carbohydrates: A review. *Nutrition and Metabolism*, 1(1), 1–24.

Geer, E. B., & Shen, W. (2009). Gender differences in insulin resistance, body composition, and energy balance. *Gender Medicine*, 6(1), 60–75.

Gilliam, T., Freedson, P., Grenen, D., & Shahraray, B. (1981). Physical activity patterns determined by heart rate monitoring in 6±7 year-old children. *Medicine and Science in Sports and Exercise*, 13(1): 65-67.

Goon, D. T., Toriola, A.L., Shaw, B. S., Shaw, L., Arogbonlo, S., & Wuam, S. (2009). Gender differences in waist circumference in *Nigerian* children. *African Journal for Physical Activity and Health Sciences*, 15(1), 91-101.

Goran, M. I. (1998). Measurement issues related to studies of childhood obesity : assessment of body composition, body fat distribution, physical activity, and food intake. *Pediatrics*, 101(2), 505–518.

Goran, M. I., Gower, B. A. (1999). Relation between visceral fat and disease risk in children and adolescents. *The American Journal of Clinical Nutrition*, 70(1), 149-156.

Guo, S. S., Chumlea, W. C. (1999). Tracking of body mass index in children in relation to overweight in adulthood. *The American Journal of Clinical Nutrition*, 70(1), 145–148.

Hassink, S. G., Sheslow, D. V., de Lancey, E., Opentanova, I., Considine, R.V., & Caro, J. F. Serum leptin in children with obesity: relationship to gender and development. *Pediatrics*, 98(2), 201-203.

He, Q., Horlick, M., Thornton, J., Wang, J., Pierson, R. N., Stanley Heshka, J.S., & Gallagher, D. (2002). Sex and race differences in fat distribution among Asian, African-American, and Caucasian prepubertal children. *The Journal of Clinical Endocrinology & Metabolism*, 87(5), 2164–2170.

He, X., Li, Z., Tang, X., Zhang, L., Wang, L., He, Y., Jin, T. and Yuan, D. (2018). Age- and sex-related differences in body composition in healthy subjects aged 18 to 82 years. *Medicine*, 97(25), 111-125.

Heymsfield, S., Peterson, C., Thomas, D., Heo, M. and Schuna, J. (2015). Why are there race/ethnic differences in adult body mass index-adiposity relationships? A quantitative critical review. *Obesity Reviews*, 17(3), 262-275.

- Houtkooper, L. B. (2000). Body composition. *Sport Nutrition for Health and Performance*, 13(1), 199–219.
- Hübers, M., Pourhassan, M., Braun, W., Geisler, C. and Müller, M. (2017). Definition of new cut-offs of BMI and waist circumference based on body composition and insulin resistance: differences between children, adolescents and adults. *Obesity Science & Practice*, 3(3), 272-281.
- Hussain, Z., Jafar, T., Zaman, M., Parveen, R. and Saeed, F. (2014). Correlations of skin fold thickness and validation of prediction equations using DEXA as the gold standard for estimation of body fat composition in Pakistani children. *BMJ Open*, 4(4), 41-94.
- Janz, K., nielsen, D., cassady, S., cook, J., Wu, Y. & Hansen, J. (1993). Cross-validation of the Slaughter skinfold equations for children and adolescents. *Medicine & Science in Sports & Exercise*, 25(9), 1070-1076.
- Kohrt, W. M. (1997). Dual-Energy X-Ray Absorptiometry: Research Issues and Equipment. *Emerging Technologies for Nutrition Research*, 6, 151-167.
- Küpper, J., Bartz, M., Schultink, J. W., Lukito, W., & Deurenberg, P. (1998). Measurements of body fat in Indonesian adults: Comparison between a three-compartment model and widely used methods. *Asia Pacific Journal of Clinical Nutrition*, 7(1), 49-54.
- Leeson, T. S., Paparo, A. A.. (1985). *Textbook of histology*. Philadelphia: W.B. Saunders.
- Maffeis, C., Grezzani, A., Pietrobelli, A., Provera, S., & Tato, L. (2001). Does waist circumference predict fat gain in children? *International Journal of Obesity*, 25(7), 978–983.
- Marshall, W. A., Tanner, J. M. (1969). Variations in pattern of pubertal changes in girls. *Archives of Disease in Childhood*, 44(235), 291–303.
- Marshall, W. A., Tanner, J. M. (1970). Variations in pattern of pubertal changes in boys. *Archives of Disease in Childhood*, 45(239), 13-23.
- Matsushita, Y., Nakagawa, T., Shinohara, M., Yamamoto, S., Takahashi, Y., Mizoue, T., Yokoyama, T., & Noda, M. (2014) How can waist circumference predict the body composition? *Diabetology & Metabolic Syndrome*, 6(1), 11-22.
- McCarthy, H. D., Jarrett, K. V., & Crawley, H. F. (2001). The development of waist circumference percentiles in British children aged 5.0-16.9 y. *European Journal of Clinical Nutrition*, 55(10), 902-907.
- Ndagire, C. T., Muyonga, J. H., Isabirye, D., Odur, B., Somda, S. M., Bukenya, R., & Nakimbugwe, D. (2018). Assessing the reliability of FTIR spectroscopy measurements and validity of bioelectrical impedance analysis as a surrogate measure of body composition among children and adolescents aged 8–19 years attending schools in Kampala, Uganda. *BMC Public Health*, 18(1), 687-696.

- Par'i, H. M., Wiyono, S., & Harjatmo, T. P. (2017). *Status Penilaian Gizi*. Jakarta: EGC
- Pramesti, R. A. A. Y. (2011). Hubungan lama pemberian ASI sampai 6 bulan pertama dengan komposisi tubuh pada ibu pasca melahirkan di Rumah Sakit Ibu dan Anak Happy Land Daerah Istimewa Yogyakarta. Skripsi. Tidak diterbitkan. Fakultas Kedokteran, Kesehatan Masyarakat, dan Keperawatan. Universitas Gadjah Mada: Yogyakarta.
- Roemmich, J. N., Rogol, A. D. (1999). Hormonal changes during puberty and their relationship to fat distribution. *American Journal of Human Biology*, 11(2), 209–224.
- Schwandt, P., & Haas, G. (2012). Waist circumference in children and adolescents from different ethnicities. *Childhood Obesity*, 79–95.
- Seino, S., Shinkai, S., Iijima, K., Obuchi, S., Fujiwara, Y., Yoshida, H., Kawai, H., Nishi, M., Murayama, H., Taniguchi, Y., Amano, H. and Takahashi, R. (2015). Reference values and age differences in body composition of community-dwelling older japanese men and women: a pooled analysis of four cohort studies. *PLOS ONE*, 10(7): e0131975.
- Shaw, N. J., Crabtree, N. J., Kibirige, M. S., & Fordham, J. N. (2007). Ethnic and gender differences in body fat in British schoolchildren as measured by DXA. *Archives of Disease in Childhood*, 92(10), 872–875.
- Sijtsma, A., Bocca, G., Labée, C., Liem, E. T., Sauer, P. J., & Corpeleijn, E. (2014). Waist-to-height ratio, waist circumference and BMI as indicators of percentage fat mass and cardiometabolic risk factors in children aged 3–7 years. *Clinical Nutrition*, 33(2), 311–315.
- Silva, D., Ribeiro, A., Pavão, F., Ronque, E., Avelar, A., Silva, A. and Cyrino, E. (2013). Validity of the methods to assess body fat in children and adolescents using multi-compartment models as the reference method: a systematic review. *Revista da Associação Médica Brasileira*, 59(5), 475–486.
- Skerlj, B., Brozek, J., Hunt, E. E., (1953). Subcutaneous fat and age changes in body build and body form in female. *The American Journal of Physical Anthropology*, 11(4), 577–600.
- Sochung Chung, MD, P. (2015). Body mass index and body composition scaling to height in children and adolescent. *Annals of Pediatric Endocrinology & Metabolism*, 20(3), 125–129.
- Šporin, D. Z. (2014). Waist circumference is appropriate for quantifying body fat in university female students. *Anthropological Notebooks*, 20(2), 145–153.
- Staiano, A. E., Broyles, S. T., Gupta, A. K., & Katzmarzyk, P. T. (2013). Ethnic and sex differences in visceral, subcutaneous, and total body fat in children and adolescents. *Obesity*, 21(6), 1251–1255.
- Sugianto, Fauzan M., Setyani, A., & Prihatini, M. (2013). *Riskesdas Dalam Angka*

- Daerah Istimewa Yogyakarta Tahun 2013*. Jakarta: Badan Penelitian dan Pengembangan Kesehatan Kementerian Kesehatan RI
- Supariasa, I. D. N., Bakri, B., Fajar, I. (2012). *Penilaian Status Gizi*. Jakarta: EGC
- Taylor, R. W., Gold, E., Manning, P., & Goulding, A. (1997). Gender differences in body fat content are present well before puberty. *International journal of obesity and related metabolic disorders*, 21(11), 1082-1084.
- Taylor, R. W., Jones, I. E., Williams, S. M., and A. G. (2000). Evaluation of waist circumference, waist to hip ratio, and the conicity index as screening tools for high trunk fat mass, as measured by dual energy x-ray absorptiometry in children aged 3-19y. *The American Journal of Clinical Nutrition*, 72(2), 490-495.
- Taylor, R., 1 , Grant, A. M., Williams, S. M., & Goulding, A.(2010). Sex differences in regional body fat distribution from pre- to postpuberty. *Obesity*, 18(7), 1410-1416.
- Verduin, W., Van Den Helder, R., Doodeman, H., Struijf, E. and Houdijk, A. (2016). Dexa body composition assessment in 10-11 year healthy children. *PLOS ONE*, 11(10), e0165275.
- Wang, J. J., Wang, H. J., Liu, J. S., & Jun, M. A. (2013). The association between body mass index, waist circumference with body fat percent, and abdominal fat rate in overweight and obese pupils. *Chinese Journal of Preventive Medicine*, 47(7): 603-607.
- Wells, J. C. K., & Fewtrell, M. S. (2006). Measuring body composition. *Archives of Disease in Childhood*, 91(7), 612-617.
- Wirawan, N. N. (2016). Sensitifitas dan spesifisitas imt dan lingkaran pinggang-panggul dalam mengklasifikasikan kegemukan pada wanita. *Indonesian Journal of Human Nutrition*, 3(1), 49 – 59.
- Wickramasinghe, V. P., Lamabadusuriay, S. P., 1 , Cleghorn, G. J., & Davies, P. S. (2008). Use of skin-fold thickness in Sri Lankan children : comparison of several prediction equations. *Indian Journal of Pediatrics*, 75(12), 1237-1242.
- World Health Organization. (2019). *BMI-for-age (5-19 years)*. Retrieved from: https://www.who.int/growthref/who2007_bmi_for_age/en/
- World Health Organization. (2008). *Waist Circumference and Waist-Hip Ratio: Report of a WHO Expert Consultation*. Geneva, World Health Organization.
- Xiong, K., He, H., Zhang, Y., & Ni, G. (2012). Analyses of body composition charts among younger and older Chinese children and adolescents aged 5 to 18 years. *BMC Public Health*, 12(1), 835-844.
- Yang, C., & Wang, L. (2017). Comparisons of waist circumference measurements at five different anatomical sites in chinese children, *BioMed Research International*, 2017, 1-8.