

REFERENCES

- A review on phytopharmaceutical importance of asiaticoside. (2016). *J. coast. life med*, 4(12), pp.1000-1007.
- ALLEN, A. (2000). Localization and function of angiotensin AT1 receptors. *American Journal of Hypertension*, 13(1), pp.S31-S38.
- Banday, A. and Lokhandwala, M. (2008). Oxidative stress-induced renal angiotensin AT1 receptor upregulation causes increased stimulation of sodium transporters and hypertension. *American Journal of Physiology-Renal Physiology*, 295(3), pp.F698-F706.
- Bartosz, M., Kedziora, J. and Bartosz, G. (1997). Antioxidant and Prooxidant Properties of Captopril and Enalapril. *Free Radic Biol Med*, 23(5), pp.729-735.
- Baynest, H. (2015). Classification, Pathophysiology, Diagnosis and Management of Diabetes Mellitus. *J Diabetes Metab*, 06(05).
- Campbell, K. and Yacoub, R. (2015). Inhibition of RAS in diabetic nephropathy. *Int J Nephrol Renovasc Dis*, p.29.
- DeFronzo, R. (2009). From the Triumvirate to the Ominous Octet: A New Paradigm for the Treatment of Type 2 Diabetes Mellitus. *Diabetes*, 58(4), pp.773-795.
- Gheith, O., Othman, N., Nampoory, N., Halimb, M. and Al-Otaibi, T. (2016). Diabetic kidney disease: difference in the prevalence and risk factors worldwide. *J Egypt Soc Nephrol Transplant*, 16(3), p.65.
- Giribabu, N., Srinivasarao, N., Swapna Rekha, S., Muniandy, S. and Salleh, N. (2014). Centella asiatica Attenuates Diabetes Induced Hippocampal Changes in Experimental Diabetic Rats. *Evidence-Based Complementary and Alternat Med*, 2014, pp.1-10.
- Goossens, G. (2012). The Renin-Angiotensin System in the Pathophysiology of Type 2 Diabetes. *Obesity Facts*, 5(4), pp.611-624.
- Guo, D., Uno, S. and Inagami, T. (1995). Steroid hormones upregulate rat angiotensin II type 1A receptor gene: Role of glucocorticoid responsive elements in rat angiotensin II type 1A promoter. *The Journal of Steroid Biochemistry and Molecular Biology*, 53(1-6), pp.69-73.
- Hsu, F., Lin, F., Ou, H., Huang, S. and Wang, C. (2017). Renoprotective Effect of Angiotensin-Converting Enzyme Inhibitors and Angiotensin II Receptor Blockers in Diabetic Patients with Proteinuria. *Kidney and Blood Press Res*, 42(2), pp.358-368.

- Kabir, A., Samad, M., D'Costa, N., Akhter, F., Ahmed, A. and Hannan, J. (2014). Anti-hyperglycemic activity of *Centella asiatica* is partly mediated by carbohydrase inhibition and glucose-fiber binding. *BMC Complementary and Altern Med*, 14(1).
- KWON, K., BAE, S., KIM, K., AN, I., AHN, K., AN, S. and CHA, H. (2014). Asiaticoside, a component of *Centella asiatica*, inhibits melanogenesis in B16F10 mouse melanoma. *Mol Med Rep*, 10(1), pp.503-507.
- Le, T., Kim, H., Allen, A., Spurney, R., Smithies, O. and Coffman, T. (2003). Physiological Impact of Increased Expression of the AT 1 Angiotensin Receptor. *Hypertension*, 42(4), pp.507-514.
- Lim, A. (2014). Diabetic nephropathy – complications and treatment. *Int J Nephrol Renovasc*, p.361.
- Lin, Y., Chang, Y., Yang, S., Wu, K. and Chu, T. (2018). Update of pathophysiology and management of diabetic kidney disease. *J Formos Med Assoc*, 117(8), pp.662-675.
- Loc, N. and Nhat, N. (2013). Production of asiaticoside from centella (*Centella asiatica* L. Urban) cells in bioreactor. *Asian Pac J Trop Biomed*, 3(10), pp.806-810.
- Mun'im, A., Ahmad, I., Yanuar, A. and Mulia, K. (2017). Review of angiotensin-converting enzyme inhibitory assay: Rapid method in drug discovery of herbal plants. *Pharmacognosy Reviews*, 11(21), p.1.
- Pharmacologic Approaches to Glycemic Treatment: Standards of Medical Care in Diabetes—2019. (2018). *Diabetes Care*, 42(Supplement 1), pp.S90-S102.
- Ogawa, S., Mori, T., Nako, K., Kato, T., Takeuchi, K. and Ito, S. (2006). Angiotensin II Type 1 Receptor Blockers Reduce Urinary Oxidative Stress Markers in Hypertensive Diabetic Nephropathy. *Hypertension*, 47(4), pp.699-705.
- Ozougwu, O. (2013). The pathogenesis and pathophysiology of type 1 and type 2 diabetes mellitus. *J. Physiol. Pathophysiol*, 4(4), pp.46-57.
- Punthakee, Z., Goldenberg, R. and Katz, P. (2018). Definition, Classification and Diagnosis of Diabetes, Prediabetes and Metabolic Syndrome. *Can J Diabetes*, 42, pp.S10-S15.
- Sen, S., Kanter, M., Ustundag, S., Aktas, C., Dogutan, H. and Yalcin, O. (2008). Effect of Angiotensin-Converting Enzyme Inhibition and Angiotensin II Type 1 Receptor Blockade on Streptozotocin-Induced Diabetic Nephropathy. *Renal Failure*, 30(10), pp.1023-1033.
- Siragy, H. (2009). The angiotensin II type 2 receptor and the kidney. *J of the Renin-Angiotensin-Aldosterone Syst*, 11(1), pp.33-36.

- STEWEN, P., MERVAALA, E., KARPPANEN, H., NYMAN, T., SAIJONMAA, O., TIKKANEN, I. and FYHRQUIST, F. (2003). Sodium Load Increases Renal Angiotensin Type 1 Receptors and Decreases Bradykinin Type 2 Receptors. *Hypertension Research*, 26(7), pp.583-589.
- Toth-Manikowski, S. and Atta, M. (2015). Diabetic Kidney Disease: Pathophysiology and Therapeutic Targets. *J Diabetes Res*, 2015, pp.1-16.
- Umanath, K. and Lewis, J. (2018). Update on Diabetic Nephropathy: Core Curriculum 2018. *Am J Kidney Dis*, 71(6), pp.884-895.
- Wang, Z., Liu, J. and Sun, W. (2013). Effects of asiaticoside on levels of podocyte cytoskeletal proteins and renal slit diaphragm proteins in adriamycin-induced rat nephropathy. *Life Sci*, 93(8), pp.352-358.
- Xu, H., Cheng, Y., Wang, W., Wu, H., Zhang, Y., Zang, C. and Xu, Z. (2016). 12-Lipoxygenase Inhibition on Microalbuminuria in Type-1 and Type-2 Diabetes Is Associated with Changes of Glomerular Angiotensin II Type 1 Receptor Related to Insulin Resistance. *Int J Mol Sc*, 17(5), p.684.
- Zhang, Z., Ma, J., Feng, R. and Wang, Z. (2018). Centella asiatica inhibits renal interstitial brosis by regulating Smad3 and Smad7 expression in the TGF β signaling pathway. *Int J Clin Exp Pathol*, 11(2), pp.1009-1017.