

DAFTAR PUSTAKA

- Aanta RE., Kanto JH., Scheinin M., Kallio AML., Scheinin H. Dexmedetomidine Premedication for Minor Gynecologic Surgery in *Anesth Analg*, 1990, 70(2):407-13.
- Abbot Information. Precedex TM Dexmedetomidine Hydrochloride Injection in *Draft Labeling Abbot Laboratories, USA*, 1999.
- Aono J., Ueda W., Mamiya K., Takemoto E., Manabe M. Greater incidence of delirium during recovery from sevoflurane anesthesia in preschool boys. *Anesthesiology*, 1997, 87: 1298-300.
- Aouad M.T., Nasr V.G., Emergence agitation in children: an update. *Curr Opin Anaesthesiology*, 2005, 18:614-9.
- Aouad, Marie T., Yazbeck-Karam, Vanda G., Nasr, Viviane G., El-Khatib, Mohamad F., Kanazi, Ghassan E., Bleik, Jamal H., A single dose of propofol at the end of surgery for the prevention of emergence agitation in children undergoing strabismus surgery during sevoflurane anesthesia. *Anesthesiology*, November, 2007, 107:733-738.
- Berde CB., Todres ID. Recovery from Anesthesia and the Postanesthesia Care Unit. Dalam: *A Practice of Anesthesia for Infants and Children*. Cote CJ, Ryan JF, Todres ID, Goudsouzian NG (Editor). WB Saunders Co. 1993, p. 471-82
- Bortone L., Ingelmo P., Grossi S., Grattagliano C., Bricchi C., Barantani D. Emergence agitation in preschool children: double-blind, randomized, controlled trial comparing sevoflurane and isoflurane anesthesia. *Paediatr Anaesth*, 2006, 16:1138-43.
- Bovill JG., Sebel PS., Blackburn CL., Heykants J. The Pharmacokinetics of Fentanyl (R 339209): a New Opioid Analgesic. *Anesthesiology*, 1982, 57:439.
- Buhrer M., Mappes A., Lauber R., 1994, Dexmedetomidine decreases Thiopenthal Dose Requirement and alters Distribution Pharmacokinetics in *Anesthesiology*, 1994, 80:6.
- Carey WB. The Effect of General Medical Illness and Its Treatment on Development and Behavior. Dalam: *Developmental-Behavioral Pediatrics*. Levine MD, Carey WB, Crocker AC, Gross RT (editor). WB Saunders Co., 1983, p. 447-55.
- Cohen I.T., Drewsen S., Hanallah R.S., Propofol or midazolam do not reduce the incidence of emergence agitation associated with desflurane anesthesia in children undergoing adenotonsilectomy. *Paediatric Anaesthesiology*, 2002, 12:604-9.
- Cohen IT., Hannallah RS., Hummer KA. The incidence of emergence agitation associated with desflurane anesthesia in children is reduced by fentanyl. *Anesth Analg*, 2001, 93:88-91.
- Cohen IT., Latta K., Weiner ES., Davis PJ. A Study of Ketorolac for Intraoperative and Postoperative Analgesia for Herniorrhaphy in Children. (Abstract). *Anest Analg*, 1993, p. 76: S50.

- Cole JW., Murray DJ., McAllister JD., Hirshberg GE. Emergence behaviour in children: defining the incidence of excitement and agitation following anaesthesia. *Paediatr Anaesth*, 2002, 12:442-7.
- Collins VJ., Boswell MV. Fluorinated Ether Anesthetics. In: *Physiologic and Pharmacologic Bases of Anesthesia*, Williams and Wilkins, Pennsylvania, 1996, p. 687-733.
- Corall IM., Moore AR., Strunin L. Plasma Concentrations of Fentanyl in Normal Surgical Patients and Those with Severe Renal and Hepatic Disease. *Br J Anaesth*, 1980, 52:101.
- Cote CJ., Preoperatif Preparation and Premedication. *Br J. Anaesth*, 1999, 83:16-28
- Cravero J., Surgenor S., Whalen K. Emergence agitation in paediatric patients after sevoflurane anaesthesia and no surgery: a comparison with halothane. *Paediatr Anaesth*, 2000, 10:419-24.
- Davis P., Greenberg J.A., Gendelman M., Recovery characteristics of sevoflurane and halothane in preschool aged children undergoing bilateral myringotomi and pressure equalization tube insertion. *Anesthesia Analgesia*, 1999, 88:34-8.
- Davis P.J., Newer inhaled pharmacologic agents in pediatric anesthesia and their effects on recovery. *ASA review course*. Chapter 3, vol 31. Philadelphia: Lippincott Williams & Wilkins, 2003.
- Demirbilek S., Tugal T., Cicek M., Aslan U., Sizanli E., Ersoy M.O., Effects of fentanyl on the incidence of emergence agitation in children receiving desflurane or sevoflurane anaesthesia. *Eur J Anaesthesiol*, 2004, 21:538-542.
- Dyck JB., Shafer SL. Dexmedetomidine Pharmacokinetics and Pharmacodynamics in *Anesthetic Pharmacology*, Review Vol. I, 1993, p.238-45.
- Ebert T., Hall JE. The Effect of increasing Plasma Concentration of Dexmedetomidine in Humans in *Anesthesiology*, August, Vol. 93, 2000, p. 2-6
- Eckenhoff J.E., Kneale D.H., Dripps R.D., The incidence and etiology of postanesthetic excitement. *Anesthesiology*, 1961, 22:667.
- Elitsur Y., Blankenship P., Lawrence Z., Propofol sedation for endoscopic Procedures In Children, *Endoscopy*, 2000, 32:788-791, [Http://www.Medscape.Com/Viewarticle/434090](http://www.Medscape.Com/Viewarticle/434090).
- Essam M.M., Ashraf A.A., Elsayed A.M., Fentanyl Versus Dexmedetomidine Effect On Agitation After Sevoflurane Anaesthesia, *Saudi Journal of Anaesthesia*, 2007, 1(2):57-61.
- Feld LH., Negus JB., White PF. Oral Midazolam Preanesthetic Medication in Pediatric Patients. *Anesthesiology*, 1990, 73:831-4.
- Freye E., Hartung E., Schenk GK. Bremazoxine: An Opiate that Induces Sedation and Analgesia without Respiratory Depression. *Anesth Analg*, 1983, 62:483.
- Galford RE. Problems in anesthesiology: approach to diagnosis. Boston, MA: Little, Brown & Company. 1992, p. 341 3.

- Hall JE., Toni UD., Shahbaz AR., Thomas EJ. 2000, Sedative, Amnestic and Analgesic properties of Small-Dose Dexmedetomidine Infusions in *Anesthesia & Analgesia*, 2000, 90(3): 699-705.
- Hanallah R.S., Ambulatory anesthesia for pediatric patients. ARS review course lectures. *Suppl Anesthesia Analgesia*, 1996, 67-73.
- Hayashi Y., Maze M. Alpha2 Adrenoceptor Agonists and Anaesthesia in *British Journal of Anaesthesia*, Vol. 71, 1993, p.108-18.
- Isik B., Arslan M., Tunga AD., Kurtipeç O. Dexmedetomidine decreases emergence agitation in pediatric patients after sevoflurane anesthesia without surgery. *Paediatr Anaesth*, 2006, 16:748-53.
- Kain ZN., Caldwell-Andrews AA., Maranets I., McClain B., Gaal D., Mayes LC. Preoperative anxiety and emergence delirium and postoperative maladaptive behaviors. *Anesth Analg*, 2004, 99:1648-54.
- Kain ZN., Mayes LC., Cicchetti DV., Bagnal AL., Finley JD., Hofstadter MB. The Yale Preoperative Anxiety Scale: How Does It Compare with a "Gold Standard"? *Anesth Analg*, 1997, 85:783-8.
- Kain ZN., Wang SM., Mayes LC., Caramino LA., Hofstadter MB. Distress During the Induction of Anesthesia and Postoperative Behavioral Outcomes. *Anesth Analg*, 1999a, 88:1042-7
- Kain ZN., Mayes LC., Wang SM., Hofstadter MB. Postoperative Behavioral Outcomes in Children: Effects of Sedative Premedication. *Anesthesiology*, 1999b, 90:758-65
- Kamibayashi T, Maze. Clinical uses of α_2 - adrenergic Agonists in *Anesthesiology*, November, Vol. 93, 2000, p. 5-16
- Lapin SL., Auden SM., Goldsmith LJ., Reynolds AM. Effects of sevoflurane anaesthesia on recovery in children: a comparison with halothane. *Paediatr Anaesth*, 1999, 9:299-304.
- Lichter JL., Collins VJ. Benzodiazepine. Dalam: *Physiologic and Pharmacologic Bases of Anesthesia*. William & Wilkins, Baltimore. 1996, p. 527-43.
- Liu S., Chiu AA., Neal JM., Carpenter RL., Bainton BG., Gerancher JC. Oral Clonidine Prolongs Lidocaine Spinal Anesthesia in Human Volunteer. *Anesthesiology*, 1995, 82:1353-1359
- Madiyono, B., Moeslichan, S., Sastroasmoro, S., Budiman, I., Purwanto, S.H., Perkiraan besar sampel. Dalam: Sastroasmoro, S., Ismael, S., editor. *Dasar-Dasar Metodologi Penelitian Klinis*. Edisi 2. Binarupa Aksara, Jakarta, 2002, p. 259-85.
- Martini DR. Commentary: The diagnosis of delirium in pediatric patients. *J Am Acad Child Adolesc Psychiatry*, 2005, 44:395-8.
- Mauricio E.I., Herná'n R. Muñoz., Verena Brandes., Anita L. Morales., Single-Dose Dexmedetomidine Reduces Agitation After Sevoflurane Anesthesia in Children. *Chile Anesth Analg*, 2004, 98:60-3
- Mc Clain DA., Hug CC. Intravenous Fentanyl Kinetics. *Clin Pharmacol Ther*, 1980, 28:106
- Mc Graw T. Preparing children for the operating room: psychological issues. *Can J Anesth*, 1994, 41:1094-103.

- Mc Millan CO., Spohr-Schopfer IA., Sikisch N., Hartley I., Lerman J. Premedication of children with oral midazolam, *Can Janesth*, 1992, 39:545-50.
- Mencia SB., López-Herce JC., Freddi N. Analgesia and sedation in children: practical approach for the most frequent situations. *JPediatr (Rio J)*, 2007, 83:S71-82.
- Meuldermans WEG., Hurkmans RNA., Heykants IJP. Plasma Protein Binding and Distribution of Fentanyl, Sufentanyl, Alfentanyl, Lofentanyl in Blood. *Arch Int Pharmacol Ther*, 1982, 257:4.
- Meyer R.R., Munster P., Werner C., Brambrink A.M., Isoflurane is associated with a similar incidence of emergence agitation/delirium as sevoflurane in young children-a randomized controlled study. *Pediatric Anesthesiology*, 2007, 17:56-60.
- Mikawa K., Maekawa N., Nishina K., Takao Y., Yaku H., Obara H. Efficacy of Oral Clonidine Premedication in Children. *Anesthesiology*, 1993, 79:926-31
- Morgan GE., Mikhail MS., Murray MJ. Clinical Pharmacology Inhalational Anesthetics in: *Clinical Anesthesiology*, 3rd ed, Lange Medical Books/McGraw-Hill, New York, 2002, p. 127-146
- Peden CJ., Prys-Robert C. Dexmedetomidine-A Powerful New Adjunct to Anaesthesia? In *British Journal of Anaesthesia*, February, Vol 6, No. 2, 1992, p. 123-5.
- Picard V., Dumont L., Pellegrini M. Quality of recovery in children: sevoflurane versus propofol. *Acta Anaesthesiol Scand*, 2000, 44:307-10.
- Przybylo HJ., Martini DR., Mazurek AJ., Bracey E., Johnsen L., Cote CJ. Assessing behaviour in children emerging from anaesthesia: can we apply psychiatric diagnostic techniques? *Paediatr Anaesth*, 2003, 13:609-16.
- Reich A., Beland B., VanAken H., Intravenous narcotics and analgesics agents. In: *Bisonette, penyunting. Pediatric anesthesia: principles and practice*. Edisi ke-1. McGraw-Hill, New York, 2002, p. 264-5.
- Reves JG., Glass PSA., Lubarsky DA. Nonbarbiturate Intravenous Anesthetics in *Anesthesia*, 5th Ed by Miller RD, Churchill Livingstone, 2000, p. 259-61.
- Sarner J.B., Levine M., Davis P.J., Lerman J., Cook D.R., Motoyama E.K., Clinical characteristic of sevoflurane in children; a comparison with halothane (clinical investigation). *Anesthesiology*, 1995, 82(1):38-46.
- Scheinin H., Aantaa R., Markku A. Reversal of the Sedative Effect of Dexmedetomidine with a Specific α_2 -Adrenoceptor Antagonist Atipamezole: A Pharmacodynamic and Kinetic Study in Healthy Volunteers in *Anesthesiology*, 1998, Vol.89, No.3.
- Shibata S., Shigeomi S., Sato W., Enzan K. Nitrous oxide administration during washout of sevoflurane improves postanesthetic agitation in children. *J Anesth*, 2005, 19:160-3.
- Sikich N., Lerman J. Development and psychometric evaluation of the pediatric anesthesia emergence delirium scale. *Anesthesiology*, 2004, 100:1138-45.
- Stoekel HH., Hengstmann JH., Schuttler. Pharmacokinetics of Fentanyl As a Possible Explanation for Reurrence of Respiratory Depression. *Br J Anaesth*, 1979, 51:741.

- Talke. Postoperative Pharmacokinetics and Sympatholytic Effects of Dexmedetomidine in *Anesthesia Analgesia*, Vol. 85, 1997, p. 1136-42.
- Tantani S. Deksmetomidin: Farmakologi dan Penggunaan dalam Praktek Anestesiologi dalam *Anestesiologi Indonesia*, Vol. I No.3, November, 2000, p. 119-123.
- Thaib MR. Perbandingan Pemberian Klonidin dan Lidokain terhadap Respon Kardiovaskuler selama Laringoskopi dan Intubasi dalam *Anestesiologi Indonesia*, November, Vol 1, No. 3, 2000, p. 131-7
- Thaib MR. Deksmetomidin, α_2 -Adrenoseptor Agonis Bani dalam *Scientific Programme 1 st Symposium on Obstetric Anesthesia*, Bandung, 2001, p. 118-30.
- Tripi PA., Palermo TM., Thomas S., Goldfinger MM., Florentino-Pineda I. Assessment of risk factors for emergence distress and postoperative behavioral changes in children following general anaesthesia. *Pediatr Anaesth*, 2004, 14:235-40.
- Venn RM., Hell J., Grounds RM. Respiratory Effect of Dexmedetomidine in the Surgical Patient Requiring Intensive Care in *Critical Care*, 2000, 4(5):302-308.
- Veyckemans F., Excitation phenomena during sevofluran anaesthesia in children. *Curr Opin Anaesthesiology*, 2001, 14:339-43.
- Vitanen H., Anilla P., Viitanen M., Yli-Hankala A. Midazolam Premedication Delays Recovery from Propofol Induced Sevoflurane Anesthesia in Children 1-3 yr. *Can J Anesth*, 1999, 46:766-771.
- Voepel-Lewis T., Malviya S., Tait A.R., A prospective cohort study of emergence agitation in the pediatric postanesthesia care unit. *Anesthesia Analgesia*, 2003, 96:1625-30.
- Welborn L.G., Hanallah R.S., Norden J.M., Ruttimann U.E., Callan C.M., Comparison of emergence and recovery characteristics of sevoflurane, desflurane, and halothane in pediatric ambulatory patients. *Anesthesia Analgesia*, 1996, 83:917-20.
- Weldon B.C., Bell M., Craddock T., The effect of caudal analgesia on emergence agitation in children after sevoflurane versus halothane anesthesia. *Anesthesia Analgesia*, 2004, 98:321-6.