

DAFTAR PUSTAKA

- Alberti, K. (2009). Harmonizing the metabolic syndrome: a joint interim statement of the International Diabetes Federation Task Force on Epidemiology and Prevention; National Heart, Lung, and Blood Institute; American Heart Association; World Heart Federation; International. *Circulation*, 120(16), 1640-1645.
- Asif, M., Aslam, M., & Altaf, S. (2018). Mid-upper-arm circumference as a screening measure for identifying children with elevated body mass index: a study for Pakistan. *Korean Journal of Pediatrics*, 61(1), 6-11.
- Avram, A. S., & Avram, M. M. (2005). Subcutaneous fat in normal and diseased states. *Journal of The American Academy of Dermatology*, 56(3), 472-492.
- Barba, G., Casullo, C., Russo, M. D., Russo, P., Nappo, A., Lauria, F., et al. (2008). Gender-related differences in the relationships between blood pressure, age, and body size in prepubertal children. *American Journal of Hypertension*, 21(9), 1007-1010.
- Bassareo, P. P., Marras, A. R., Barbanti, C., & Mercuro, G. (2013). Comparison between waist and mid-upper arm circumferences in influencing systolic blood pressure in adolescence: the SHARP (Sardinian Hypertensive Adolescent Research Programme) study. *Journal of Pediatric and Neonatal Individualized Medicine*, 2(2), 202-207.
- Berenson, G., Srinivasan, S., Bao, W., Newman, W., Tracy, R., & Wattigney, W. (1998). Association between multiple cardiovascular risk factors and atherosclerosis in children and young adults: The Bogalusa Heart Study. *New England Journal of Medicine*, 338(23), 1650-1656.
- Bergheanu, S., Bodde, M., & Jukema, J. (2017). Pathophysiology and treatment of atherosclerosis. *Netherlands Heart Journal*, 25(4), 231-242.
- Birmingham, B., Dyer, A., & Shekelle, R. B. (1993). Subscapular and triceps skinfold thickness, body mass index, and cardiovascular risk factors in a cohort of middle-aged employed men. *Journal of Clinical Epidemiology*, 46(3), 289-302.
- Brozek, J. (1956). *Body Measurements and Human Nutrition* (Vol. 14). Detroit: Wayne University Press.
- Bucher, B. S., & Ferrarini, A. (2013). Primary hypertension in childhood. *Pediatric Hypertension*, 15(5), 444-452.
- Burkinshaw, L., Jones, P. R., & Krupowicz, D. W. (1973). Observer error in skinfold thickness measurements. *Human Biology*, 45(2), 273-279.
- Center for Disease Control and Prevention. (2007). Anthropometry procedures manual. *National Health and Nutrition Examination Survey*, 1(3), 15-16.

- Center for Disease Control and Prevention. (2016, October 20). *Overweight and Obesity*. Retrieved April 19, 2018, from Defining Childhood Obesity: <https://www.cdc.gov/obesity/childhood/defining.html>
- Coleman, E. (2017, Desember 6). *SF Gate*. Retrieved Mei 22, 2018, from The Recommended Amount & Percent of Carbohydrates Per Day: <http://healthyeating.sfgate.com/recommended-amount-percent-carbohydrates-per-day-7287.html>
- Cowell, C., Briody, J., Llyod-Jones, S., Smith, C., Moore, B., & Howman-Giles, R. (1997). Fat distribution in children and adolescents– the influence of sex and hormones. *Hormonal Research*, 48(5), 93-100.
- de Almeida, C. A., Del Ciampo, L. A., Ricco, R. G., & Silva Jr, S. M. (2003). Assessment of mid-upper arm circumference as a method for obesity screening in preschool children. *Journal of Pediatric (Rio J)*, 79(5), 455-460.
- de Swiet, M., Fayers, P., & Shinebourne, E. (1992). Blood pressure in first 10 years of life: the Brompton study. *British Medical Journal*, 304(6818), 23-26.
- Diaz, K. M., & Shimbo, D. (2013). Physical activity and prevention of hypertension. *Current Hypertension Report*, 15, 659-658.
- Dietz, W. (1998). Health consequences of obesity in youth: childhood predictors of adult disease. *American Academy of Pediatrics*, 101(2), 521-522.
- Elwald, R., & Hadelman, L. (2016). Risk factors in adolescent hypertension. *Global Pediatric Health*, 3(1), 1-26.
- Freedman, D., & Wang, J. (2007). The prediction of body fatness by BMI and skinfold thicknesses among children and adolescents. *Annals of Human Biology*, 34(2), 183-194.
- Freedman, D., Katzmarzyk, P., Dietz, W., Srinivasan, S., & Berenson, G. (2009). Relation of body mass index and skinfold thicknesses to cardiovascular disease risk factors in children: the Bogalusa Heart Study. *The American Journal of Clinical Nutrition*, 90(1), 210-216.
- Frisancho, A. (1974). Triceps skin fold and upper arm muscle size norms for assesment of nutritional status. *The American Journal of Clinical Nutrition*, 27(10), 1052-1058.
- George, S. M. (2016). Reliability of 24-hour dietary recalls as a measure of diet in African American youth. *Journal of Academy of Nutrition and Diet*, 116(10), 1551-1559.
- Grimes, W. B., & Franzini, L. R. (1977). Skinfold measurement techniques for estimating percentage body fat. *Journal of Behavioural Therapy and Experimental Psychiatry*, 8(1), 65-69.
- Gruskin, A. (1995). Factors affecting blood pressure. *Pediatric Nephrology: Pediatric and Adolescent Medicine*, 3, 1047.

- Guillery, E., & Robillard, J. (1993). Guillery, E. N., & Robillarde the renin-angiotensin system and blood pressure regulation during infancy and childhood. *Pediatric Clinics of North America*, 40(1), 61-79.
- Guyton, A. C., & Hall, J. E. (2011). *Guyton and Hall Textbook of Medical Physiology*. Philadelphia: Elsevier.
- Haris, Syafruddin; Dimiati, Herlina; Anwar, M. Sidqi. (2013). Profil hipertensi pada anak di RSUD Zaenal Abidin Banda Aceh. *Sari Pediatri*, 15(2), 105-110.
- Henson, M. (2014, Januari 3). *Kaiser Permanente*. Retrieved Mei 23, 2018, from Balancing Carbs, Protein, and Fat: <https://wa.kaiserpermanente.org/healthAndWellness?item=%2Fcommon%2FhealthAndWellness%2Fconditions%2Fdiabetes%2FfoodBalancing.html>
- Hong, Y. (2010). Atherosclerotic cardiovascular disease beginning in childhood. *Korean Circulation Journal*, 40(1), 1-9.
- Jaiswal, M., Bansal, R., & Agarwal, A. (2017). Role of mid-upper arm circumference for determining overweight and obesity in children and adolescents. *Journal of Clinical and Diagnostic Research*, 11(8), SC05-SC08.
- Järvisalo, M., Putto-Laurila, A., Jartti, L., Lehtimäki, T., Solakivi, T., Rönnemaa, T., et al. (2002). Carotid artery intima-media thickness in children with type 1 diabetes. *Diabetes*, 51(2), 493-498.
- Joglekar, C., Fall, C., Deshpande, V., Joshi, N., & al, e. (2007). Newborn size, infant and childhood growth, and body composition and cardiovascular disease risk factors at the age of 6 years: the Pune Maternal Nutrition Study. *International Journal of Obesity (London)*, 31(10), 1534-1544.
- Joint National Committee VIII. (2014). *2014 Evidence-based guideline for the management of high blood pressure in adults: report from the panel members appointed to the Eighth Joint National Committee (JNC 8)*. JAMA.
- Kaur, J. (2014). A Comprehensive Review on Metabolic Syndrome. *Cardiology Research and Practice*.
- Kementerian Kesehatan Republik Indonesia. (2013). *Riset Kesehatan Dasar*. Kesehatan.
- Kurukutasuriya, L., Stas, S., Lastra, G., Manrique, C., & Sowers, J. (2011). Hypertension in obesity. *Medical Clinics of North America*, 95(5), 903-917.
- Ledwaba, K. R., Nkalanga, F., & Monyeke, K. D. (2014). The Relationship Between Mid-Upper Arm Circumference and Blood Pressure of Private School Children Aged 6-13 Years, in Polokwane, Limpopo Province, South Africa. *Annals of Pediatric Child Health*, 2(4), 1026.
- Liu, Y., Li, Y., He, J., Ma, P., Yu, L., Zheng, Q., et al. (2018). Gender stratified analyses of the association of skinfold thickness with hypertension: a cross-sectional study in general

northeastern chinese residents. *International Journal of Environmental Research and Public Health*, 15(12), 2748.

- Ma, C. m., Li, Y., Gao, G. q., Yin, F. z., Wang, R., & Liu, X. l. (2015). Mid-upper arm circumference as a screening measure for identifying children with hypertension. *Blood Pressure Monitoring*, 20(4), 189-193.
- Mayer, J. (1968). Overweight, causes, cost, and control. *Journal of Pediatrics*, 44(1), 196-211.
- Mazicioglu, M., Hatipoglu, N., Ozturk, A., Cicek, B., Ustunbas, B., & Kurtoglu, S. (2010). Waist circumference and mid-upper arm circumference in evaluation of obesity in children between 6 and 17 years. *Journal of Clinical Research on Pediatric Endocrinology*, 2(4), 144-150.
- Mazigoglu, M., Hatipolu, N., Ozturk, A., Cicek, B., Ustunbafl, H., & Kurtolu, S. (2010). Waist circumference and mid-upper arm circumference in evaluation of obesity in children aged between 6 and 17 years. *Journal of Clinical Research in Pediatric Endocrinology*, 2, 144-150.
- National High Blood Pressure Education Program. (2004). The fourth report on the diagnosis, evaluation, and treatment of high blood pressure in children and adolescents. *Pediatrics*, 114(2), 555-576.
- National Research Council (US) Committee on Diet and Health. (1989). *6Calories: Total Macronutrient Intake, Energy Expenditure, and Net Energy Stores*. Washington DC: National Academies Press.
- Norton, K., & Olds, T. (2004). *Anthropometrica: a textbook of body measurement for sports and health courses*. Sydney: University of New South Wales Press.
- O'Brien, E. T., & O'Malley, K. (1979). ABC of blood pressure measurement. *British Medical Journal*, 322(7292), 981-985.
- Olsen, M. (2016). A call to action and a lifecourse strategy to address the global burden of raised blood pressure on current and future generation. *Lancet Commision on Hypertension*, 388(10060), 2665-2712.
- Pinto, E. (2007). Blood pressure and ageing. *Postgraduate Medical Journal*, 83(976), 109-114.
- Poissonnet, C., & Lavelle, M. (1988). Growth and development of adipose tissue. *Journal of Pediatric*, 113(1), 1-9.
- Pungky, A., & Damanik, M. (2006). Hipertensi pada anak di RS Dr. Sardjito Yogyakarta. *Berita Kedokteran Masyarakat*, 15(2), 105-109.
- Rao, S. (1983). Blood pressure and triceps skinfold thickness. *Annals of Human Biology*, 10(2), 191-193.

- Rochmah, W., Wilopo, S., & Aswin, S. (1984). Hubungan antara tekanan darah dan ukuran antropometrik sekelompok pelajar sekolah tingkat pertama di kotamadya Yogyakarta. *Berkala Ilmu Kedokteran*, 16, 95-99.
- Rosner, B., Prineas, R., Loggie, J., & Daniels, S. (1993). Blood pressure normograms for children and adolescents, by height, sex, and age, in the United States. *Journal of Pediatrics*, 123(6), 871-886.
- Seltzer, C. C., & Mayer, J. (1965). A simple criterion of obesity. *Postgraduate Medicine*, 38(2), 101-107.
- Seltzer, C., & Mayer, J. (1965). A Simple Criterion of Obesity. *Postgraduate Medicine*, 101-107.
- Sharma, A. (2004). Is there a rationale for angiotensin blockade in the management of obesity hypertension? *Hypertension*, 44(1), 12-19.
- Sherwood, L. (2013). *Human Physiology : From Cells to Systems* (8th ed.). (S. Alexander, Ed.) Belmont, California: Brooks/Cole.
- Shimokata, H., Tobin, J., Muller, D., Elahi, D., Coon, P., & Andres, R. (1989). Studies in distribution of body fat : effects of age, sex, and obesity. *Journal of Gerontology*, 44(2), M66-M73.
- Siervogel, R. (1983). Genetic and familial factors in essential hypertension and related traits. *American Journal of Physical Anthropology*, 26 (S1), 37-63.
- Sims, E. (1973). Endocrine and metabolic effects of experimental obesity in man. *Recent Progress on Hormonal Research*, 29, 457-496.
- Sloan, A., & Saphiro, M. (1972). A comparison of skinfold measurement with three standard calipers. *Human Biology*, 44(1), 29-36.
- Soeliman, F. (2014). Weight loss maintenance: A review on dietary related strategies. *Journal of Research in Medical Science*, 19(3), 268-275.
- Sorof, J., & Daniels, S. (2002). Obesity hypertension in children : a problem of epidemic proportions. *Hypertension*, 40(4), 41-447.
- Sowers, J. (2013). Diabetes mellitus and vascular disease. *Hypertension*, 61(5), 943-947.
- Swift, D. L. (2014). The Role of Exercise and Physical Activity in Weight Loss and Maintenance. *Cardiovascular disease*.
- Tortora, G. J., & Derrickson, B. (2009). *Principles of Anatomy and Physiology*. Hoboken: John Wiley & Sons, Inc.
- Turton, P. (1986). The use of mid upper arm circumference in the assessment of nutritional status: The Mursi. *Midwife Health Visit Community Nurse*, 21, 81-86.
- Wahdah, N. (2011). *Menaklukan Hipertensi dan Diabetes (Mendeteksi, Mencegah, dan Mengobati dengan Cara Medis dan Herbal)*. Yogyakarta: Multipress.

- Whitlock, E., & Williams, S. (2010). Screening and interventions for childhood overweight: a summary of evidence for the US Preventive Services Task Force. *Pediatrics*, 116(1), 125-144.
- Williams, D. (1992). Body fatness and risk for elevated blood pressure, total cholesterol, and serum lipoprotein ration in children and adolescents. *American Journal on Public Health*, 82(3), 358-363.
- World Health Organization. (2004). *A Glossary of Terms for Community Health Care and Services for Older Persons Volume 5*. Technical Report, Centre for Health Development.
- Xi, B., Zhang, T., Zhang, M., Liu, F., Zong, X., Zhao, M., et al. (2015). Trends in elevated blood pressure among US children. *American Journal of Hypertension*, 29(2), 217-225.
- Ying-Xiu, Z., & Shu-Rong, W. (2011). Distribution of skinfold thickness and blood pressure among children and adolescents in Shandong, China. *Journal of Tropical Pediatrics*, 57(4), 258-262.
- Yuhasz, M. (1974). *Physical Fitness and Sports Appraisal Laboratory Manual*. Canada: University of Western Ontario.
- Zhang, Y., & Wang, S. (2011). The relationship of body mass index distribution to relatively high blood pressure among children and adolescents in Shandong, China. *Annals of Human Biology*, 38, 630-634.
- Zhang, Y., & Wang, S. (2013). The relationship of waist circumference distribution to blood pressure levels among children and adolescents in Shandong, China. *International Journal of Cardiology*, 168(2), 1516-1520.