

## REFERENCES

- Abitbol CL, Moxey-Mims M (2016). Low birth weight and the global burden of kidney disease: Chronic kidney disease. *Nat Rev Nephrol*, 12(4):199–200.
- Andayasari L, Opitasari C (2016). Parity and risk of low birth weight infant in full term pregnancy. *Health Sci J Indones*, 7(1): 13–16.
- Andriani H, Kuo H-W (2014). Adverse effects of parental smoking during pregnancy in urban and rural areas. *BMC Pregnancy Childbirth*, 14(414):1–15.
- Anggrahini SM, Notobroto HB, Triyoga RS (2018). Socio-Demographic and Cultural Factors Associated with The Occurrence of Low Birth Weight in Kupang. *Humanist Netw Sci Technol*, 2(2): 169–173.
- Aryastami NK, Shankar A, Kusumawardani N, Besral B, Jahari AB, Achadi E (2017). Low birth weight was the most dominant predictor associated with stunting among children aged 12–23 months in Indonesia. *BMC Nutr*, 3(16):1–6.
- Astuti S, Susanti AI, Elista R (2016). Gambaran Paparan Asap Rokok Pada Ibu Hamil Berdasarkan Usia Kehamilan di Desa Cintamulya Kecamatan Jatinangor Kabupaten Sumedang. *J. Sist. Kesehat*, 2.
- Bacci S, Bartolucci F, Chiavarini M, Minelli L, Pieroni L (2014). Differences in Birthweight Outcomes: A Longitudinal Study Based on Siblings. *Int. J. Environ. Res. Public Health*, 11: 6472 – 6484.
- Boggess KA (2005). Pathophysiology of Preterm Birth: Emerging Concepts of Maternal Infection. *Clin. Perinatol*, 32(3): 561–569.
- Cantarutti A, Franchi M, Monzio MC, Merlino L, Corrao G (2017). Mother's education and the risk of several neonatal outcomes: an evidence from an Italian population-based study. *BMC Pregnancy Childbirth*, 17(221):1–10.
- Chaiton M, Diemert L, Cohen JE, Bondy SJ, Selby P, Philipneri A, Schwartz R (2016). Estimating the number of quit attempts it takes to quit smoking successfully in a longitudinal cohort of smokers. *BMJ Open*, 6(6): 1–11.
- Cui H, Gong TT, Liu CX, Wu QJ (2016). Associations between Passive Maternal Smoking during Pregnancy and Preterm Birth: Evidence from a Meta-Analysis of Observational Studies. *PLOS ONE* 11(1):1–18.
- De Boo HA, Harding JE (2006). The developmental origins of adult disease (Barker) hypothesis. *Aust N Z J Obstet Gynaecol*, 46(1): 4–14.
- Dhillon PK, Jeemon P, Arora NK, Mathur P, Maskey M, Sukirna RD, Prabhakaran D (2012). Status of epidemiology in the WHO South-East Asia region: burden of disease, determinants of health and epidemiological research, workforce and training capacity. *Int J Epidemiol*, 41(3):847–860.

- Eftekhar M, Purmasumi S, Sabeti P, Mirhosseini F (2016). Relation of Second Hand Smoker and Effect on Pregnancy Outcome and Newborns Parameters. *Womens Health Gynecol*, 2(2):1–6.
- Foulstone AR, Kelly AB, Kifle T (2017). Partner influences on smoking cessation: A longitudinal study of couple relationships. *J Subst Use*, 22(5):501–506.
- Health Ministry of Republic Indonesia (2013). Riset Kesehatan Dasar RI 2013. Jakarta, Badan Penelitian dan Pengembangan Kesehatan
- Hudić I, Stray-Pedersen B, Tomić V (2015). Preterm Birth: Pathophysiology, Prevention, Diagnosis, and Treatment. *BioMed Res Int*, 1.
- Inoue S, Naruse H, Yorifuji T, Kato T, Murakoshi T, Doi H, Subramanian S (2016). Association between Short Maternal Height and Low Birth Weight: a Hospital-based Study in Japan. *J Korean Med Sci*, 31(3):353–359.
- Khader YS, Al-Akour N, AlZubi IM, Lataifeh, I (2011). The Association Between Second Hand Smoke and Low Birth Weight and Preterm Delivery. *Matern. Child Health J*. 15: 453–459.
- Ko TJ, Tsai LY, Chu LC, Yeh SJ, Leung C, Chen CY, Chou HC, Tsao PN, Chen PC, Hsieh WS (2014). Parental Smoking During Pregnancy and Its Association with Low Birth Weight, Small for Gestational Age, and Preterm Birth Offspring: A Birth Cohort Study. *Pediatr Neonatol*, 55(1): 20–27.
- Koo YJ, Ryu HM, Yang JH, Lim JH, Lee JE, Kim MY, Chung JH (2012). Pregnancy outcomes according to increasing maternal age. *Taiwan J Obstet Gynecol*, 51(1):60–65
- Kovac JR, Labbate C, Ramasamy R, Tang D, Lipshultz LI (2015). Effects of cigarette smoking on erectile dysfunction. *Andrologia*, 47(10):1087–1092.
- Krishnamurthy AV, Chinnakali P, Dorairajan G, Sundaram SP, Sarveswaran G, Sivakumar M, Krishnamoorthy K, Dayalane H, Sinouvassan V (2018). Tobacco use, exposure to second-hand smoke among pregnant women and their association with birth weight: A retrospective cohort study. *J Fam Med Prim Care*, 7(4):728–733.
- Kumar S, Gonzalez F (2018). Effects of health insurance on birth weight in Mexico: Insurance and Birth Weight in Mexico. *Health Econ*, 27(8):1149–1159.
- Kusumawardani N, Tarigan I, Suparmi, Schlottheuber A (2018). Socio-economic, demographic and geographic correlates of cigarette smoking among Indonesian adolescents: results from the 2013 Indonesian Basic Health Research (RISKESDAS) survey. *Glob Health Action*, 11(Suppl 1):1–9.
- Lima SAM, El Dib RP, Rodrigues MRK, Ferraz GAR, Molina AC, Neto CAP, de Lima MAF, Rudge MVC (2018). Is the risk of low birth weight or preterm labor greater when maternal stress is experienced during pregnancy? A systematic review and meta-analysis of cohort studies. *PLOS ONE*, 13(7):1–15.

- Mahumud RA, Sultana M, Sarker AR (2017). Distribution and Determinants of Low Birth Weight in Developing Countries. *J Prev Med Pub Health*, 50(1):18–28.
- Momenabadi V, Kaveh MH, Mousavi SM, Alizadeh S (2017). Maternal Risk Factors Associated with Low Birth Weight. *Iran J Health Sci*, 5(3):58–64
- Metgud, CS, Naik VA, Mallapur MD (2012). Factors Affecting Birth Weight of a Newborn – A Community Based Study in Rural Karnataka, India. *PLOS ONE*, 7: 1-4.
- Niu Z, Xie C, Wen X, Tian F, Yuan S, Jia D, Chen W-Q (2016). Potential pathways by which maternal second-hand smoke exposure during pregnancy causes full-term low birth weight. *Sci Rep*, 6(24987):1–8.
- Nobile CGA, Raffaele G, Altomare C, Pavia M (2007). Influence of maternal and social factors as predictors of low birth weight in Italy. *BMC Public Health*, 7(192):1–9.
- Norsa'adah B, Salinah O (2014). The Effect of Second-Hand Smoke Exposure during Pregnancy on the Newborn Weight in Malaysia. *Malays J Med Sci*, 21(2):44–53.
- Özaltın E, Hill K, Subramanian SV (2010). Association of Maternal Stature With Offspring Mortality, Underweight, and Stunting in Low- to Middle-Income Countries. *JAMA*, 303(15):1507–1516.
- Pathirathna M, Abeywickrama H, Sekijima K, Sadakata M, Fujiwara N, Muramatsu Y, Wimalasiri K, Jayawardene U, de Silva D, Dematawewa C (2017). Effects of Prenatal Tobacco and Wood-Fuel Smoke Exposure on Birth Weight in Sri Lanka. *Healthcare*, 5(4):1–10.
- Pei L, Kang Y, Zhao Y, Cheng Y, Yan H (2016). Changes in Socioeconomic Inequality of Low Birth Weight and Macrosomia in Shaanxi Province of Northwest China, 2010 – 2013: A Cross-sectional Study. *Medicine*, 95: 1 – 8.
- Restrepo-Méndez MC, Lawlor DA, Horta BL, Matijasevich A, Santos IS, Menezes AMB, Barros FC, Victora CG (2015). The Association of Maternal Age with Birthweight and Gestational Age: A Cross-Cohort Comparison: The association of maternal age with birthweight and gestational age. *Paediatr Perinat Epidemiol*, 29(1):31–40.
- Rokhmah NL (2013). Hubungan Usia Ibu Dengan Kejadian Bayi Berat Lahir Rendah di RS PKU Muhammadiyah Yogyakarta Tahun 2012. *Sekol. Tinggi Ilmu Kesehatan. Aisyiyah*, 1: 1-10.
- Schane RE, Ling PM, Glantz SA (2010). Health Effects of Light and Intermittent Smoking: A Review. *Circulation*, 121(13):1518–1522.
- Schwartz IL (1990). Low-Birth-Weight Effects of Demographic and Socioeconomic Variables and Prenatal Care in Pima County, Arizona. *West J Med*, 152(6):725–728.

- Silva AA, Silva LM, Barbieri MA, Bettiol H, Carvalho LM, Ribeiro VS, Goldani MZ (2010). The epidemiologic paradox of low birth weight in Brazil. *Rev Saude Publica*, 44(5):767–775.
- Silvestrin S, da Silva CH, Hiraqueta VN, Goldani AAS, Silveira PP, Goldani MZ (2013). Maternal education level and low birth weight: a meta-analysis. *J Pediatr (Rio J)*, 89(4):339–345.
- Simbolon D (2014). Pengaruh Kepemilikan Jaminan Kesehatan Masyarakat Miskin Terhadap Status Kelahiran dan Kejadian Stunting Pada Baduta Indonesia (Analisis Data IFLS 1993-2007). *J. Kebijak. Kesehat. Indones*, 3: 56–65.
- Singh A, Sahoo N (2013). Urban–rural differentials in the factors associated with exposure to second-hand smoke in India. *BMJ Open*, 3(11):1–8.
- Stratton K, 2015. *Public Health Implications of Raising the Minimum Age of Legal Access to Tobacco Products*. National Academies Press: Washington, D.C.
- Syed W, Kamathi VC (2012). Maternal short stature: A risk factor for low birth weight in neonates. *J Med Allied Sci*, 2(2):62–65.
- Tan TYT, Yeo GSH (2005). Intrauterine growth restriction. *Curr Opin Obstet Gynecol*, 17(2):135–142.
- Tsai Y-W, Tsai T-I, Yang C-L, Kuo KN (2008). Gender Differences in Smoking Behaviors in an Asian Population. *J Womens Health*, 17(6):971–978.
- Vettore MV, Gama SG, Lamarca Gde A, Schilithz AO, Leal Mdo C (2010). Housing conditions as a social determinant of low birthweight and preterm low birthweight. *Rev Saude Publica*, 44(6):1021–1031.
- Wahabi HA, Alzeidan RA, Fayed AA, Mandil A, Al-Shaikh G, Esmail SA (2013). Effects of secondhand smoke on the birth weight of term infants and the demographic profile of Saudi exposed women. *BMC Public Health* 13: 341-346.
- Wang L, Yang Y, Liu F, Yang A, Xu Q, Wang Q, Shen H, Zhang Y, Yan D, Peng Z, He Y, Wang Y, Xu J, Zhao J, Zhang H, Zhang Y, Dai Q, Ma X (2018). Paternal smoking and spontaneous abortion: a population-based retrospective cohort study among non-smoking women aged 20–49 years in rural China. *J Epidemiol Community Health*, 72(9):783–789.
- Wardlaw, T.M., World Health Organization, UNICEF (Eds.), 2004. *Low birthweight: country, regional and global estimates*. WHO ; UNICEF, Geneva : New York.
- World Health Organization, Bloomberg Philanthropies, 2017. *WHO report on the global tobacco epidemic, 2017: monitoring tobacco use and prevention policies*.
- Yerushalmy J (2014). The relationship of parents' cigarette smoking to outcome of pregnancy—implications as to the problem of inferring causation from observed associations. *Int J Epidemiol*, 43(5):1355–1366.