

REFERENCES

- Aminoff, M., Greenberg, D. and Simon, R. (2014). *Clinical Neurology*. 9th ed. McGraw-Hill.
- Asadollahi, K., Beeching, N. and Gill, G. (2010). Leukocytosis as a predictor for non-infective mortality and morbidity. *QJM*, 103(5), pp.285-292.
- Chen, R., Ovbiagele, B. and Feng, W. (2016). Diabetes and Stroke: Epidemiology, Pathophysiology, Pharmaceuticals and Outcomes. *The American Journal of the Medical Sciences*, 351(4), pp.380-386.
- Counsell, C., Dennis, M., McDowall, M. and Warlow, C. (2002). Predicting Outcome After Acute and Subacute Stroke: Development and Validation of New Prognostic Models. *Stroke*, 33(4), pp.1041-1047.
- Di Carlo, A., Lamassa, M., Baldereschi, M., Pracucci, G., Basile, A., Wolfe, C., Giroud, M., Rudd, A., Ghetti, A. and Inzitari, D. (2003). Sex Differences in the Clinical Presentation, Resource Use, and 3-Month Outcome of Acute Stroke in Europe. *Stroke*, 34(5), pp.1114-1119.
- Bagg s., Pombo A.P., Hopman W. 2002. Effect of Age on Functional Outcomes After Stroke Rehabilitation. *Stroke*. 33:179-185.
- Doehner, W., Schenkel, J., Anker, S., Springer, J. and Audebert, H. (2012). Overweight and obesity are associated with improved survival, functional outcome, and stroke recurrence after acute stroke or transient ischaemic attack: observations from the TEMPiS trial. *European Heart Journal*, 34(4), pp.268-277.
- Bang O.Y., Park H.Y., Lee P.H., Kim G.M., Chung C.S., Lee K.H., 2007. Improved Outcome after atherosclerotic stroke in male smoker. *Journal of the Neurological Sciences*. 260(1-2) :43-48
- Ducrocq, G., Amarenco, P., Labreuche, J., Alberts, M., Mas, J., Ohman, E., Goto, S., Lavallée, P., Bhatt, D. and Steg, P. (2013). A History of Stroke/Transient Ischemic Attack Indicates High Risks of Cardiovascular Event and Hemorrhagic Stroke in Patients With Coronary Artery Disease. *Circulation*, 127(6), pp.730-738.
- Echouffo-Tcheugui, J., Xu, H., Matsouaka, R., Xian, Y., Schwamm, L., Smith, E., Bhatt, D., Hernandez, A., Heidenreich, P. and Fonarow, G. (2018). Diabetes and long-term outcomes of ischaemic stroke: findings from Get With The Guidelines-Stroke. *European Heart Journal*, 39(25), pp.2376-2386.
- Eckel, R. (2011). *Metabolic risk for cardiovascular disease*. Dallas, TX: American Heart Association.
- Gabriel, F., Dannug, A. and Noble, P. (2017). Leukocytosis as a predictor of poor outcomes in acute ischemic stroke among adults: A meta analysis. *Journal of the Neurological Sciences*, 381, pp.505-506.
- Guyton, A. and Hall, J. (2006). *Textbook of medical physiology*. Philadelphia: Elsevier Saunders.

- Intercollegiate Stroke Working Party. National clinical guideline for stroke. (2016). 5th ed. London: Royal College of Physicians.
- Kumar, A., Boehme, A., Siegler, J., Gillette, M., Albright, K. and Martin-Schild, S. (2013). Leukocytosis in Patients with Neurologic Deterioration after Acute Ischemic Stroke is Associated with Poor Outcomes. *Journal of Stroke and Cerebrovascular Diseases*, 22(7), pp.e111-e117.
- Kumar, V., Abbas, A., Aster, J. and Perkins, J. (n.d.). *Robbins basic pathology*.
- Kusuma, Y., Venketasubramanian, N., Kiemas, L. and Misbach, J. (2009). Burden of Stroke in Indonesia. *International Journal of Stroke*, 4(5), pp.379-380.
- Lee, J.-H., Lee, J.Y., Ahn, S.H., Jang, M.U., Oh, M.S., Kim, C.-H., Yu, K.-H. and Lee, B.-C., 2015. Smoking is Not a Good Prognostic Factor following First-Ever Acute Ischemic Stroke. *Journal of Stroke*, 17(2), p.177.
- Liao, C., Shih, C., Yeh, C., Chang, Y., Hu, C., Lin, J. and Chen, T. (2015). Impact of Diabetes on Stroke Risk and Outcomes. *Medicine*, 94(52), p.e2282.
- Mallmann, A., Fuchs, S., Gus, M., Fuchs, F. and Moreira, L. (2012). Population-Attributable Risks for Ischemic Stroke in a Community in South Brazil: A Case-Control Study. *PLoS ONE*, 7(4), p.e35680.
- Nardi, K., Milia, P., Eusebi, P., Paciaroni, M., Caso, V. and Agnelli, G. (2012). Admission Leukocytosis in Acute Cerebral Ischemia: Influence on Early Outcome. *Journal of Stroke and Cerebrovascular Diseases*, 21(8), pp.819-824.
- Niewada, M., Kobayashi, A., Sandercock, P., Kamiński, B. and Członkowska, A. (2005). Influence of Gender on Baseline Features and Clinical Outcomes among 17,370 Patients with Confirmed Ischaemic Stroke in the International Stroke Trial. *Neuroepidemiology*, 24(3), pp.123-128.
- Sacco, R., Kasner, S., Broderick, J., Caplan, L., Connors, J., Culebras, A., Elkind, M., George, M., Hamdan, A., Higashida, R., Hoh, B., Janis, L., Kase, C., Kleindorfer, D., Lee, J., Moseley, M., Peterson, E., Turan, T., Valderrama, A. and Vinters, H. (2013). An Updated Definition of Stroke for the 21st Century: A Statement for Healthcare Professionals From the American Heart Association/American Stroke Association. *Stroke*, 44(7), pp.2064-2089.
- Semplicini, A., Maresca, A., Boscolo, G., Sartori, M., Rocchi, R., Giantin, V., Forte, P.L. and Pessina, A.C., 2003. Hypertension in Acute Ischemic Stroke. *Archives of Internal Medicine*, 163(2), p.211.
- Vemmos, K., Ntaios, G., Spengos, K., Savvari, P., Vemmou, A., Pappa, T., Manios, E., Georgiopoulos, G. and Alevizaki, M. (2011). Association Between Obesity and Mortality After Acute First-Ever Stroke. *Stroke*, 42(1), pp.30-36.



Zhou, J., Zhang, Y., Arima, H., Zhao, Y., Zhao, H., Zheng, D., Tian, Y., Liu, Y., Huang, Q. and Yang, J., 2014. Sex differences in clinical characteristics and outcomes after intracerebral haemorrhage: results from a 12-month prospective stroke registry in Nanjing, China. *BMC Neurology*, 14(1).