

DAFTAR PUSTAKA

- Anderson, S., Halter, B.J., Hazzard, R.W., Himmelfarb, J. 2009. Prediction, Progression, and Outcomes of Chronic Kidney Disease in Older Adults. *J Am Soc Nephrol* 20: 1199–1209.
- Andrews. 1999. Disorders of Iron Metabolism. *New England J Med*, 341(26): 1986–1995.
- Anonim1. 2010. No Title Reticulocyte and Their Significance. Sysmex Xtra Online. http://www.sysmex.ru/files/articles/Xtra_online_reticulocytes.pdf.
- Anonim2. 2010. Reticulocyte Hemoglobin in Anemia Diagnosis and Management. *America, Sysmex. Inc.*
- Bahrainwala, J. & Berns, J.S. 2016. Diagnosis of Iron-Deficiency Anemia in Chronic Kidney Disease. *Semin Nephrol*, 36(2): 94–98.
- Bandiara, R. 2003. Penatalaksanaan anemi defisiensi besi pada pasien yang menjalani hemodialisis. : 1–9.
- Brugnara, C. 2002. A Hematologic “Gold Standard” for Iron-deficient States? In *Clin Chem*. Boston: 981–982.
- Brugnara, C., Schiller, B. & Moran, J. 2006. Reticulocyte hemoglobin equivalent (Ret He) and assessment of iron-deficient states. *Clin Lab Haematol*, 28(5): 303–308.
- Chang, C.C. & Kass, L. 1997. Clinical significance of immature reticulocyte fraction determined by automated reticulocyte counting. *American Journal of Clinical Pathology*, 108(1): 69–73.
- Coyne, D.W., Goldsmith, D. & Macdougall, I.C. 2017. New options for the anemia of chronic kidney disease. *Kidney Int Suppl*, 7(3): 157–163.
- Dalimunthe, N.N. & Lubis, A.R. 2016. Usefulness of Reticulocyte Hemoglobin Equivalent in Management of Regular Hemodialysis Patients with Iron Deficiency Anemia. *Rom J Intern Med*, 54(1): 31–36.
- Denggo, S.D., Muhadi, D. & Arif, M. 2013. *Anemia pada Penyakit Ginjal Kronik*.
- Fischbach, F.A., Gregory, D.W., Harrison, P.M., Hoy, T.G. & Williams, J.M. 1971. On the structure of hemosiderin and its relationship to ferritin. *J Ultrastruct Res*, 37(5–6): 495–503.
- Fishbane, S., Pollack, S., Feldman, H.I. & Joffe, M.M. 2009. Iron indices in chronic kidney disease in the National Health and Nutritional Examination Survey 1988–2004. *Clin J Am soc Nephrol*, 4(1): 57–61.
- Fishbane, S., Valenzuela, O.F., Shapiro, W., Faubert, J. & Dutka, P. 2001. A randomized trial of iron deficiency testing strategies in hemodialysis patients. *Int Soc Nephrol*, 60(6): 2406–2411.
- Gafor, A.H.A., Subramaniam, R., Hadi, F., Cader, R., Yen, K., Mohd, R. & Shah, S.A. 2018. The Role of Reticulocyte Hemoglobin Content in the Management of Iron Deficiency Anemia in Patients on Hemodialysis. *Nephro-Urol Mon*, 10(3): 1–5.
- Goodnough, L.T., Skikne, B. & Brugnara, C. 2000. Erythropoietin, iron, and erythropoiesis. *Am Soc Haematol*, 96(3): 823–833.

- Hikmawati, D.A. 2017. *Internet*. Repository.unair.ac.id. 2017.
- Humphries, J.E. 1992. Anemia of renal failure: Use of erythropoietin. *Medical Clinics of North America*, 76(3): 711–725.
- IRR. 2017. 10 th Report Of Indonesian Renal Registry.
- IRR. 2016. 9 th Report Of Indonesian Renal Registry. : 1–46.
- Karagülle, M., Gündüz, E., Mutlu, F.Ş. & Akay, M.O. 2013. Clinical Significance of Reticulocyte Hemoglobin Content in the Diagnosis of Iron Deficiency Anemia. *Turk J Haematol*, 30(2): 153–156.
- KDOQI & Review, E. 2006. KDOQI Clinical Practice Guidelines and Clinical Practice Recommendations for Anemia in Chronic Kidney Disease. Executive Summary. *American journal of kidney diseases : the official journal of the National Kidney Foundation*, 47(5): S11-70.
- Kim, J.M., Ihm, C.H. & Kim, H.J. 2008. Evaluation of reticulocyte haemoglobin content as marker of iron deficiency and predictor of response to intravenous iron in haemodialysis patients. *Int J Lab Haematol*, 30(1): 46–52.
- Koca, E., Cetiner, D.A., Buyukasik, Y., Uner, A., Sayinalp, N. & Haznedaroglu, I.C. 2013. Bone Marrow Iron Staining is a Reliable Test for Elimination of Iron Deficiency Anemia Rather than its Diagnosis. *Int J Haematol Oncol*, 23(4): 260–263.
- Lameshow, S., Hosmer, D.W., Klar, J., Lwanga, K.S. 1990. Adequacy of sample size in health studies. The World Health Organization. New York.:37-38.
- Lankhorst, C.E. & Wish, J.B. 2010. Anemia in renal disease: Diagnosis and management. *Blood Reviews*, 24(1): 39–47.
- Legovini, P., Osis, J., Giacomini, A. & Beghi, L. 2000. Anemia Control in Chronic Renal Failure. *IFAC*, 33(3): 267–272.
- Lubis, A.. & Siregar, J.. 2013. Anemia pada Penyakit Ginjal Kronik. *UNSU*: 1–20.
- MacOni, M., Cavalca, L., Danise, P., Cardarelli, F. & Brini, M. 2009. Erythrocyte and reticulocyte indices in iron deficiency in chronic kidney disease: Comparison of two methods. *Scand J of Clin Labor Invest*, 69(3): 365–370.
- Mambatta, A.K., Alagesan, M., Meeran, M. & Rein, J.L. 2017. Evaluation of iron status in patients with end stage renal disease. *International Journal of Advances in Medicine*, 4(5): 1415–1421.
- Mast, A.E., Blinder, M.A. & Dietzen, D.J. 2008. Reticulocyte hemoglobin content. *Am J Hematol*, 83(4): 307–310.
- Miers, M.K., Exton, M.G. & Chapman, D.H. 2002. *Cell Counting Instrumentation in Rodak Hematology Clinical principles and Applications*. Second Edi. W.B Saunders Company, Philadelphia.
- Mikhail, A., Brown, C., Williams, J.A., Mathrani, V., Shrivastava, R., Evans, J., Isaac, H. & Bhandari, S. 2017. Renal association clinical practice guideline on Anaemia of Chronic Kidney Disease. *BMC Nephrology*, 18(1): 1–29.
- Mittman, N., Sreedhara, R., Mushnick, R., Chattopadhyay, J., Zelmanovic, D., Vaseghi, M. & Avram, M.M. 1997. Reticulocyte hemoglobin content predicts functional iron deficiency in hemodialysis patients receiving rHuEPO. *Am J Kidney Dis*, 30(6): 912–922.
- Miwa, N., Akiba, T., Hamaguchi, Y., Arakawa, Y., Tamura, T. & Nitta, K. 2010.

- Usefulness of measuring reticulocyte hemoglobin equivalent in the management of haemodialysis patients with iron deficiency. *Int J Lab Haematol*, 32(2): 248–255.
- NKF-KDOQI. 2002. *Clinical Practice Guidelines For Chronic Kidney Disease: Evaluation, Classification and Stratification*. New York.
- NKF-KDOQI. 2006. Executive Summary. *Am J Kidney Dis*, 47(5): s11–s15.
- Nurko, S. 2006. Anemia in chronic kidney disease : Causes , diagnosis , treatment. *Cleveland Clin J Med*, 73(3): 289–297.
- Ombuh, C., Rotty, L. & Palar, S. 2011. *Status Besi Pada Pasien Penyakit Ginjal Kronik Yang Sedang Menjalani Hemodialisis Di RSUD Prof.Dr.R.D Kandou Manado*. Manado.
- Pali, D.V., Moeis, E.S. & Rotty, L.W.A. 2012. Gambaran Anemia pada Penderita Penyakit Ginjal Kronik di BLU. RSUP. PROF. DR. R. D. KANDOU. : 1–7.
- PERNEFRI. 2011a. *Konsensus Manajemen Anemia Pada Penyakit Ginjal Kronik*. II. Jakarta.
- PERNEFRI. 2011b. *Konsensus Manajemen Anemia Pada Penyakit Ginjal Kronik*. II. Jakarta.
- PERNEFRI 2011. 2011. *PERNEFRI 2011.pdf*.
- Polin, V., Coriat, R., Perkins, G., Dhooge, M., Abitbol, V., Leblanc, S., Prat, F. & Chaussade, S. 2013. Iron deficiency: From diagnosis to treatment. *Dig Liver Dis*, 45(10): 803–809.
- Rafi, A., Karkar, A. & Abdelrahman, M. 2007. Monitoring Iron status in End-Stage Renal Disease Patients on Hemodialysis. *Saudi J Kidney Dis Transplant*, 18(1): 73–78.
- Ratcliffe, L.E.K., Thomas, W., Glen, J., Padhi, S., Pordes, B.A.J., Wonderling, D., Connell, R., Stephens, S., Mikhail, A.I., Fogarty, D.G., Cooper, J.K., Dring, B., Devonald, M.A.J., Brown, C. & Thomas, M.E. 2016. Diagnosis and management of iron deficiency in CKD: A summary of the NICE guideline recommendations and their rationale. *Am J Kidney Dis*, 67(4): 548–558.
- Reddy, C., Devak, R. & Rao, P. 2013. Iron Indices in Patients with Functional Anemia in Chronic Kidney Disease. *J Int Federat Clin Chem Labor Med*, 24(3): 129–136.
- Saputra, D.H. 2017. Perkembangan Terapi Erythropoietin Stimulating Agent untuk Anemia Penyakit Ginjal Kronik. *CDK*, 44: 37–42.
- Schaefer, R. & Schaefer, L. 1998. Iron monitoring and supplementation: how do we achieve the best results? *Nephrology Dialysis Transplantation*, 13(Suppl 2): 9–12. http://ndt.oxfordjournals.org/content/13/suppl_2/9.short.
- Schrier, S.L. & Auerbach, M. 2018. Causes and diagnosis of iron deficiency and iron deficiency anemia in adults. *UpToDate*.
- Silaban, B., Sugeng, C. & Waleleng, B.J. 2016. Gambaran status besi pada pasien penyakit ginjal kronik stadium 5 dengan anemia yang menjalani hemodialisis reguler. *e-Cl*, 4(1–7).
- Stauffer, M.E. & Fan, T. 2014. Prevalence of chronic kidney disease in the United States. *J Am Med Assoc*, 9(1): 1–4.
- Suega, K. 2010. Aplikasi Klinis Retikulosit. *J Peny Dalam*, 11(3): 191–201.
- Sysmex. 2012. *Sysmex Educational Enhancement & Development*.

- Szigeti, R.G. 2014. Reticulocyte Count and Reticulocyte Hemoglobin Content. emedicine.medscape.com/article/2086146-overview#showall.
- Tessitore, N., Solero, G. Pietro, Lippi, G., Bassi, A., Faccini, G.B., Bedogna, V., Gammaro, L., Brocco, G., Restivo, G., Bernich, P., Lupo, A. & Maschio, G. 2001. The role of iron status markers in predicting response to intravenous iron in haemodialysis patients on maintenance erythropoietin. *Nephrol Dial Transplant*, 16(7): 1416–1423.
- Thomas, D.W., Hinchliffe, R.F., Briggs, C., Macdougall, I.C., Littlewood, T. & Cavill, I. 2013. Guideline for the laboratory diagnosis of functional iron deficiency. *British J Haematol*, 161(5): 639–648.
- Ullrich, C., Wu, A., Armsby, C., Rieber, S., Wingerter, S., Brugnara, C., Deficiency, R.O.N., The, I.S. & Com, M. 2005. Screening Healthy Infants for Iron Deficiency Using Reticulocyte Hemoglobin Content. *JAMA*, 294(8): 924–930.
- Urrechaga, E., Boveda, O., Aguayo, F.J., de la Hera, P., Muñoz, R.I., Gallardo, I. & Escanero, J.F. 2016. Percentage of hypochromic erythrocytes and reticulocyte hemoglobin equivalent predictors of response to intravenous iron in hemodialysis patients. *Int J Lab Haematol*, 38(4): 360–365.
- Wang, W., Knovich, M.A., Coffman, L.G., Torti, F.M. & Torti, S. V. 2010. Serum ferritin: Past, present and future. *Biochim Biophys Acta*, 11(2): 760–780.
- WHO. 2007. Assessing the iron status of populations. *CDC*: 1–112.
- Wirawan, R., Tedja, A.T., Henrika, F. & Lydia, A. 2017. Concordance between Reticulocyte Hemoglobin Equivalent and Reticulocyte Hemoglobin Content in CKD Patients Undergoing Hemodialysis. *Indones J Intern Med*, 49(1): 34–40.
- Wish, J.B. 2006. Assessing iron status: beyond serum ferritin and transferrin saturation. *Clin J Am soc Nephrol*, 1: S4–S8.
- Wollmann, M., Gerzson, B.M.C., Schwert, V., Figuera, R.W. & de Oliveira Ritzel, G. 2014. Reticulocyte maturity indices in iron deficiency anemia. *Rev Bras Hematol Hemoter*, 36(1): 25–28.
- Yuh, Y.J., Kim, S.R. & Han, T.H. 2005. Immature Reticulocyte Fraction after Iron Therapy for Iron Deficiency Anemia. *Am Soc Haematol*.