

## DAFTAR PUSTAKA

- Badan Penelitian dan Pengembangan Kesehatan (2013) ‘Riset Kesehatan Dasar (RISKESDAS) 2013’, *Laporan Nasional 2013*, pp. 1–384. doi: 1 Desember 2013.
- Baghianimoghadam, M. H. *et al.* (2015) ‘Risk factors of low birth weight and effect of them on growth pattern of children up to sixth months of life: A cross-sectional study.’, *Journal of education and health promotion*, 4(May), p. 40. doi: 10.4103/2277-9531.157226.
- Committee to Study the Prevention of Low Birthweight; Division of Health Promotion and Disease Prevention; Institute of Medicine. (1985) *Preventing Low Birthweight*, Washington (DC); National Academies Press (US). doi: 10.17226/511.
- Fukuda, S. *et al.* (2015) ‘Mother’s Physique Strongly Affects Male Infant’s Physique Maternal Body Mass Index Correlates with the Neonatal Physique of Male Infants’, *Tohoku J. Exp. Med.* *Tohoku J. Exp. Med*, 237(2371), pp. 69–75. doi: 10.1620/tjem.237.69.
- Godin, G. and Kok, G. (1996) ‘The theory of planned behavior: a review of its applications to health-related behaviors.’, *American journal of health promotion : AJHP*, 11(2), pp. 87–98. doi: 10.4278/0890-1171-11.2.87.
- Goisis, A. *et al.* (2017) ‘Original Contribution Advanced Maternal Age and the Risk of Low Birth Weight and Preterm Delivery : a Within-Family Analysis Using Finnish Population Registers’, 186(11), pp. 1219–1226. doi: 10.1093/aje/kwx177.
- Hawsawi, A. M., Bryant, L. O. and Goodfellow, L. T. (2015) ‘Association Between Exposure to Secondhand Smoke During Pregnancy and Low Birthweight: A Narrative Review’, *Respiratory Care*, 60(1), pp. 135–140. doi: 10.4187/respcare.02798.
- Kader, M. and Perera, N. K. P. (2014) ‘Socio-economic and nutritional determinants of low birth weight in India’, *North American Journal of Medical Sciences*, 6(7), pp. 302–308. doi: 10.4103/1947-2714.136902.
- Kemendes RI (2013) ‘10 Perilaku hidup bersih dan sehat di rumah tangga’, *Pusat Promosi Kesehatan*.
- Kementerian Kesehatan Republik Indonesia (2016) *Profil Kesehatan Indonesia, Profil Kesehatan Provinsi Bali*.
- Kementrian Kesehatan RI (2018) ‘Profil Kesehatan Indonesia 2017

- Kiserud, T. *et al.* (2017) *The World Health Organization Fetal Growth Charts : A Multinational Longitudinal Study of Ultrasound Biometric Measurements and Estimated Fetal Weight*. doi: 10.1371/journal.pmed.1002220.
- Kurtz, M. E. *et al.* (2003) 'Knowledge and attitudes of economically disadvantaged women regarding exposure to environmental tobacco smoke', *European Journal of Public Health*, pp. 171–176.
- Mojibyan, M. *et al.* (2013) 'Exposure to Second-hand Smoke During Pregnancy and Preterm Delivery', 1(4), pp. 149–153. doi: 10.5812/ijhrba.7630.
- Nichter, M. *et al.* (2010) 'Developing a smoke free household initiative: an Indonesian case study', *Acta Obstetricia et Gynecologica Scandinavica*, 89(4), pp. 578–581. doi: 10.3109/00016340903578893.
- Norsa'Adah, B. and Salinah, O. (2014) 'The effect of second-hand smoke exposure during pregnancy on the newborn weight in Malaysia', *Malaysian Journal of Medical Sciences*, 21(2), pp. 44–53.
- Padmawati, R. S. *et al.* (2018) 'Establishing a community-based smoke-free homes movement in Indonesia', *European Network for Smoking and Tobacco Prevention*.
- Peneva, I. and Mavrodiiev, S. (2013) 'A Historical Approach to Assertiveness', *Psychological Thought; Vol 6, No 1 (2013)*. doi: 10.5964/psyct.v6i1.14.
- Peraturan Menteri Kesehatan Republik Indonesia No 97 Tahun 2014 (2014) 'Tentang Pelayanan Kesehatan Masa Sebelum Hamil, Masa Hamil, Persalinan, dan Masa Sesudah Melahirkan, Penyelenggaraan Pelayanan Kontrasepsi, serta Pelayanan Kesehatan Seksual', *Menteri Kesehatan Republik Indonesia*, (Jakarta).
- Rajantran, S. (2015) 'Level of Knowledge About Passive Smoking and Assertiveness among Passive Smokers (Women) in City of Yogyakarta'.
- Shmuelib, A. and Cullena, M. R. (2000) 'Birth Weight , Maternal Age , and Education : New Observations from Connecticut and Virginia', 72(1999), pp. 245–258.
- Soltani, H. *et al.* (2017) 'Pre-pregnancy body mass index and gestational weight gain and their effects on pregnancy and birth outcomes : a cohort study in West Sumatra , Indonesia'. *BMC Women's Health*, pp. 1–12. doi: 10.1186/s12905-017-0455-2.
- Wahabi, H. A. *et al.* (2013) 'Effects Of Environmental tobacco smoke On The Birth Weight Of Term Infants And The Demographic Profile Of Saudi Exposed Women'. *BMC Public Health Journal*. 13: 341., *BMC Public Health Journal*, 13, p. 341. Available at: <https://bmcpublichealth.biomedcentral.com/articles>.

- Weise, A. (2012) 'WHA Global Nutrition Targets 2025: Low Birth Weight Policy Brief', *W.H.O Publication*, pp. 1–7. doi: WHO/NMH/NHD/14.3.
- Wickstrom, R. (2007) 'Effects of Nicotine During Pregnancy: Human and Experimental Evidence', *Current Neuropharmacology*, 5(3), pp. 213–222. doi: 10.2174/157015907781695955.
- World Health Organization (2017) 'WHO report on the global tobacco epidemic India WHO Framework Convention on Tobacco Control (WHO FCTC) status'. Available at: [http://www.who.int/tobacco/surveillance/policy/country\\_profile/idn.pdf%0Ahttp://www.who.int/tobacco/surveillance/policy/country\\_profile/ind.pdf](http://www.who.int/tobacco/surveillance/policy/country_profile/idn.pdf%0Ahttp://www.who.int/tobacco/surveillance/policy/country_profile/ind.pdf).
- Zhang, L. *et al.* (2015) 'Exposure to Secondhand Tobacco Smoke and Interventions Among Pregnant Women in China : A Systematic Review', pp. 1–11.
- APA, 2017. *American Pregnancy Association*. [Online] Available at: <http://americanpregnancy.org/pregnancy-complications/second-hand-smoke-and-pregnancy/> [Accessed 10 4 2018].
- CDC, 2017. *Centers for Disease Control and Prevention*. [Online] Available at: [https://www.cdc.gov/tobacco/data\\_statistics/fact\\_sheets/health\\_effects/effects\\_cig\\_smoking/index.htm](https://www.cdc.gov/tobacco/data_statistics/fact_sheets/health_effects/effects_cig_smoking/index.htm) [Accessed 12 4 2018]
- QTI, 2018. *Quit Tobacco International* [Online] Available at: <https://www.quittobaccointernational.org> [Accessed 5 2018]