

## DAFTAR PUSTAKA

- Beltramo, F., Menteeer, J., Razavi, A., Khemani, R.G., Szmuszkovicz, J., Newth, C.J.L., et al. 2016. Validation of an ultrasound cardiac output monitor as a bedside tool for pediatric patients. *Pediatr. Cardiol.*, 37(1): 177–183.
- Bujang, M. and Nurakmal, B. 2017. Guidelines of the minimum sample size requirements for Cohen’s Kappa. *Epidemiol. Biostat. Public Heal.*, 14(2): 1–10.
- Cattermole, G.N., Leung, P.Y.M., Ho, G.Y.L., Lau, P.W.S., Chan, C.P.Y., Chan, S.S.W., et al. 2017. The normal ranges of cardiovascular parameters measured using the ultrasonic cardiac output monitor. *Physiol. Rep.*, 5(6): e13195.
- Cattermole, G.N., Leung, P.Y.M., Ho, G.Y.L., Lau, P.W.S., Chan, C.P.Y., Chan, S.S.W., et al. 2017. The normal ranges of cardiovascular parameters measured using the ultrasonic cardiac output monitor. *Physiol. Rep.*, 5(6): e13195.
- Chan, J.S., Segara, D. and Nair, P. 2006. Measurement of cardiac output with a non-invasive continuous wave Doppler device versus the pulmonary artery catheter: a comparative study. *Crit. Care Resusc.*, 8(4): 309–314.
- Chong, S.W. and Peyton, P.J. 2012. A meta-analysis of the accuracy and precision of the ultrasonic cardiac output monitor (USCOM). *Anaesthesia*, 67(11): 1266–1271.
- Corley, A., Barnett, A.G., Mullany, D. and Fraser, J.F. 2009. Nurse-determined assessment of cardiac output. Comparing a non-invasive cardiac output device and pulmonary artery catheter: A prospective observational study. *Int. J. Nurs. Stud.*, 46(10): 1291–1297.
- Dey, I. and Sprivilis, P. 2005. Emergency physicians can reliably assess emergency department patient cardiac output using the USCOM continuous wave Doppler cardiac output monitor. *Emerg Med Australas*, 17(3): 193–199.
- Dhanani, S., Barrowman, N.J., Ward, R.E. and Murto, K.T. 2011. Intra- and inter-observer reliability using a noninvasive ultrasound cardiac output monitor in healthy anesthetized children. *Paediatr. Anaesth.*, 21(8): 858–864.
- Flack, V.F., Afifi, A.A., Lachenbruch, P.A. and Schouten, H.J.A. 1988. Sample size determinations for the two rater kappa statistic. *Psychometrika*, 53(3): 321–325.
- Gidwani, U.K., Mohanty, B. and Chatterjee, K. 2013. The pulmonary artery catheter. a critical reappraisal. *Cardiol. Clin.*, 31(4): 545–565.
- Gnaegi, A., Feihl, F. and Perret, C. 1997. Intensive care physicians’ insufficient knowledge of right-heart catheterization at the bedside: time to act? *Crit. Care Med.*, 25(2): 213–20.
- Gwet, K.L. 2014. *Handbook of inter-rater reliability: the definitive guide to measuring the extent of agreement among raters; [a handbook for researchers, practitioners, teachers & students]*. 4. ed. Gaithersburg, MD: Advanced Analytics, LLC.
- Hodgson, L.E., Venn, R., Forni, L.G., Samuels, T.L. and Wakeling, H.G. 2015.

- Measuring the cardiac output in acute emergency admissions: use of the non-invasive ultrasonic cardiac output monitor (USCOM) with determination of the learning curve and inter-rater reliability. *J. Intensive Care Soc.*, 17(2): 122–128.
- Horster, S., Stemmler, H.J., Strecker, N., Brettner, F., Hausmann, A., Cnossen, J., et al. 2012. Cardiac output measurements in septic patients: Comparing the accuracy of USCOM to PiCCO. *Crit. Care Res. Pract.*, 2012.
- House, A.E., House, B.J. and Campbell, M.B. 1981. Measures of interobserver agreement: Calculation formulas and distribution effects. *J. Psychopathol. Behav. Assess.*, 3(1): 37–57.
- Huygh, J., Peeters, Y., Bernards, J. and Malbrain, M.L.N.G. 2016. Hemodynamic monitoring in the critically ill: an overview of current cardiac output monitoring methods. *F1000Research*, 5(0): 1–9.
- Jain, S., Allins, A., Salim, A., Vafa, A., Wilson, M.T. and Margulies, D.R. 2008. Noninvasive Doppler ultrasonography for assessing cardiac function: can it replace the Swan-Ganz catheter? *Am. J. Surg.*, 196(6): 961–968.
- Joffe, R., Duff, J., Garcia Guerra, G., Pugh, J. and Joffe, A.R. 2016. The accuracy of blood pressure measured by arterial line and non-invasive cuff in critically ill children. *Crit. Care*, 20(1): 177.
- de Jonge, R.C.J., Polderman, K.H. and Gemke, R.J.B.J. 2005. Central venous catheter use in the pediatric patient: Mechanical and infectious complications. *Pediatr. Crit. Care Med.*, 6(3): 329–339.
- Karras, D.J. 1997. Statistical methodology: II. Reliability and variability assessment in study design, Part A. *Acad. Emerg. Med.*, 4(1): 64–71.
- Knirsch, W., Kretschmar, O., Tomaske, M., Stutz, K., Nagdyman, N., Balmer, C., et al. 2008. Cardiac output measurement in children: Comparison of the Ultrasound Cardiac Output Monitor with thermodilution cardiac output measurement. *Intensive Care Med.*, 34(6): 1060–1064.
- Lemson, J., Nusmeier, A. and van der Hoeven, J.G. 2011. Advanced Hemodynamic Monitoring in Critically Ill Children. *Pediatrics*.
- Lichtenthal, P.R., Phillips, R.A., Sloniger, J.A. and Copeland, J.G. 2006. USCOM-Non Invasive Doppler – Are Cardiac Output Measurements Accurate in Both Infants and Adults? *Anesthesiology*, (January): 2006.
- McHugh, M. 2012. Interrater reliability: the kappa statistic. *Biochem. Medica*, 22(3): 276–282.
- Meyer, S., Todd, D. and Shadboldt, B. 2009. Assessment of portable continuous wave Doppler ultrasound (ultrasonic cardiac output monitor) for cardiac output measurements in neonates. *J. Paediatr. Child Health*, 45(7–8): 464–468.
- Nguyen, H.B., Banta, D.P., Stewart, G., Kim, T., Bansal, R., Anholm, J., et al. 2010. Cardiac index measurements by transcutaneous doppler ultrasound and transthoracic echocardiography in adult and pediatric emergency patients. *J. Clin. Monit. Comput.*, 24(3): 237–247.
- Nguyen, H.B., Losey, T., Rasmussen, J., Oliver, R., Guptill, M., Wittlake, W.A., et al. 2006. Interrater reliability of cardiac output measurements by transcutaneous Doppler ultrasound: implications for noninvasive

- hemodynamic monitoring in the ED. *Am. J. Emerg. Med.*, 24(7): 828–835.
- Osman, D., Ridel, C., Ray, P., Monnet, X., Anguel, N., Richard, C., et al. 2007. Cardiac filling pressures are not appropriate to predict hemodynamic response to volume challenge\*. *Crit. Care Med.*, 35(1): 64–68.
- Scheer, B., Perel, A. and Pfeiffer, U.J. 2002. Clinical review: complications and risk factors of peripheral arterial catheters used for haemodynamic monitoring in anaesthesia and intensive care medicine. *Crit. Care*, 6(3): 199–204.
- de Simone, G., Devereux, R.B., Daniels, S.R., Mureddu, G., Roman, M.J., Kimball, T.R., et al. 1997. Stroke volume and cardiac output in normotensive children and adults. *Circulation*, 95(7): 1837 LP-1843.
- Smith, B.E. 2013. *The uscom and haemodynamics a guide for junior medical and nursing staff*. Bathurst, Australia.
- Smith, B.E. 2013. *The uscom in clinical practice : a guide for junior medical and nursing staff*. Sidney.
- Stewart, G.M., Nguyen, H.B., Kim, T.Y., Jauregui, J., Hayes, S.R. and Corbett, S. 2008. Inter-rater reliability for noninvasive measurement of cardiac function in children. *Pediatr. Emerg. Care*, 24(7): 433–437.
- Tibby, S.M., Hatherill, M., Marsh, M.J. and Murdoch, I. a. 1997. Clinicians' abilities to estimate cardiac index in ventilated children and infants. *Arch. Dis. Child.*, 77(6): 516–518.
- Trieschmann, U., Cate, U.T.F. and Sreeram, N. 2007. Central venous catheters in children and neonates - what is important? *Images Paediatr. Cardiol.*, 9(4): 1–8.
- Ward, M. and Langton, J.A. 2007. Blood pressure measurement. *Contin. Educ. Anaesthesia, Crit. Care Pain*, 7(4): 122–126.
- Webster, C.S., Merry, A.F., Emmens, D.J., Van Cotthem, I.C. and Holland, R.L. 2003. A prospective clinical audit of central venous catheter use and complications in 1000 consecutive patients. *Anaesth. Intensive Care*, 31(1): 80–6.
- Wiguna, P.A., Murni, I.K. and Rusmatiningtyas. 2017. Interrater agreement of Ultrasound Cardiac Output Monitor (USCOM) between pediatric intensivist and resident in Pediatric Intensive Care Unit (PICU) (Abstract). In *Abstract of the 17th Indonesian COngress of Pediatrics*. Yogyakarta: Indonesian Pediatric Society: 200.
- Zorko, D.J., Choong, K., Gilleland, J., Agar, B., Baker, S., Brennan, C., et al. 2013. Urgent Ultrasound Guided Hemodynamic Assessments by a Pediatric Medical Emergency Team: A Pilot Study. *PLoS One*, 8(6): 6–11.