

REFERENCE

- Abu-Baker, N., Haddad, L., & Savage, C. (2010). The Influence of Secondhand Smoke Exposure on Birth Outcomes in Jordan. *International Journal of Environmental Research and Public Health*, 7(2), 616–634.
- Badan Pusat Statistik (2019). “Pendidikan”, <https://www.bps.go.id/subject/28/pendidikan.html#subjekViewTab1>, accessed in 12 April 2019 at 20:19.
- Badan Pusat Statistik (2019). “Tenaga Kerja”, <https://www.bps.go.id/subject/6/tenaga-kerja.html#subjekViewTab3>, accessed in 12 April 2019 at 19:20.
- Bernstein, I. M., Horbar, J. D., Badger, G. J., Ohlsson, A., & Golan, A. (2000). Morbidity and mortality among very-low-birth-weight neonates with intrauterine growth restriction. *American Journal of Obstetrics and Gynecology*, 182(1), 198–206.
- Bhanji, S., Andrades, M., Taj, F., Khuwaja, A.K., 2011. Factors related to knowledge and perception of women about smoking: a cross sectional study from a developing country. *BMC Women's Health* 11, 16.
- Chi, Y.-C., Wu, C.-L., Chen, C.-Y., Lyu, S.-Y., Lo, F.-E., & Morisky, D. E. (2015). Randomized trial of a secondhand smoke exposure reduction intervention among hospital-based pregnant women. *Addictive Behaviors*, 41, 117–123.
- Commar, A., Prasad, V. K., Tursan d'Espaignet, E., Wolfenden, L., & Weltgesundheitsorganisation. (2018). *WHO global report on trends in prevalence of tobacco smoking 2000-2025*.
- Cui H, Gong TT, Liu CX, Wu QJ (2016). Associations between Passive Maternal Smoking during Pregnancy and Preterm Birth: Evidence from a Meta- Analysis of Observational Studies. *PLOS ONE* 11(1):1–18.
- Dherani, M., Zehra, S. N., Jackson, C., Satyanaryana, V., Huque, R., Chandra, P., ... Siddiqi, K. (2017). Behaviour change interventions to reduce second-hand smoke exposure at home in pregnant women – a systematic review and intervention appraisal. *BMC Pregnancy and Childbirth*, 17(1).
- Gharaibeh, H., Haddad, L., Alzyoud, S., El-Shahawy, O., Baker, N. A., & Umlauf, M. (2011). Knowledge, Attitudes, and Behavior in Avoiding Secondhand Smoke Exposure Among Non-Smoking Employed Women with Higher Education in Jordan. *International Journal of Environmental Research and Public Health*, 8(11), 4207–4219.
- Gilpin, EA., Pierce, JP., Cavin, SW., Berry, CC., Evans, NJ., Johnson, M., Bal, DG. (1994). Estimates of population smoking prevalence: self-vs proxy reports of smoking status. *American Journal of Epidemiology*, 84(10), 1576-9.
- Global Burden of Disease Collaborative Network (2016). “Global Burden of Disease Study 2016 (GBD 2016) Population Estimates 1950-2016”, <https://tobaccoatlas.org/topic/consumption>, accessed in 15 April 2019 at 15:02.
- Hawsawi, A. M., Bryant, L. O., & Goodfellow, L. T. (2015). Association Between Exposure to Secondhand Smoke During Pregnancy and Low Birthweight: A Narrative Review. *Respiratory Care*, 60(1), 135–140.

- Hornik, R.C., 2002. Exposure: Theory and Evidence about All the Ways it Matters. *Social Marketing Quarterly* 8, 31–37.
- Hoyt, A. T., Canfield, M. A., Romitti, P. A., Botto, L. D., Anderka, M. T., Krikov, S. V., & Feldkamp, M. L. (2018). Does Maternal Exposure to Secondhand Tobacco Smoke During Pregnancy Increase the Risk for Preterm or Small-for-Gestational Age Birth? *Maternal and Child Health Journal*, 22(10), 1418–1429.
- Hyland, A., Cummings, KM., Lynn, WR., Corle, D., Giffen, CA. (1997). Effect of proxy-reported smoking status on population estimates of smoking prevalence. *American Journal of Epidemiology*, 145(8), 746-51.
- Jackson, C., Huque, R., Satyanarayana, V., Nasreen, S., Kaur, M., Barua, D., ... Chandra, P. (2016). “He Doesn’t Listen to My Words at All, So I Don’t Tell Him Anything”—A Qualitative Investigation on Exposure to Second Hand Smoke among Pregnant Women, Their Husbands and Family Members from Rural Bangladesh and Urban India. *International Journal of Environmental Research and Public Health*, 13(11), 1098.
- Kazemi, A., Ehsanpour, S., & Nekoei-Zahraei, N. S. (2012). A randomized trial to promote health belief and to reduce environmental tobacco smoke exposure in pregnant women. *Health Education Research*, 27(1), 151–159.
- Lee, A. H. (2008). A pilot intervention for pregnant women in Sichuan, China on passive smoking. *Patient Education and Counseling*, 71(3), 396–401.
- Lee, J., Lee, D.-R., Lee, D.-H., Paek, Y.-J., & Lee, W.-C. (2015). Influence of Maternal Environmental Tobacco Smoke Exposure Assessed by Hair Nicotine Levels on Birth Weight. *Asian Pacific Journal of Cancer Prevention*, 16(7), 3029–3034.
- Loke, A. Y., & Lam, T. H. (2005). A randomized controlled trial of the simple advice given by obstetricians in Guangzhou, China, to non-smoking pregnant women to help their husbands quit smoking. *Patient Education and Counseling*, 59(1), 31–37.
- Ma, G. X., Tan, Y., Fang, C. Y., Toubbeh, J. I., & Shive, S. E. (2005). Knowledge, attitudes and behavior regarding secondhand smoke among Asian Americans. *Preventive Medicine*, 41(2), 446–453.
- Mao, A., Bristow, K., & Robinson, J. (2013). Caught in a dilemma: why do non-smoking women in China support the smoking behaviors of men in their families? *Health Education Research*, 28(1), 153–164.
- Mojibyan, M., Karimi, M., Bidaki, R., Rafiee, P., & Zare, A. (2013). Exposure to Second-hand Smoke During Pregnancy and Preterm Delivery. *International Journal of High Risk Behaviors and Addiction*, 1(4).
- Mund, M., Louwen, F., Klingelhofer, D., & Gerber, A. (2013). Smoking and Pregnancy — A Review on the First Major Environmental Risk Factor of the Unborn. *International Journal of Environmental Research and Public Health*, 10(12), 6485–6499.
- Nihayah, Z., 2014. *Hubungan asertif dengan kebahagiaan pada mahasiswa Fakultas Psikologi Angkatan 2013 Universitas Islam Negeri Maulana Malik Ibrahim Malang* (Doctoral dissertation, Universitas Islam Negeri Maulana Malik Ibrahim).
- Norsa’adah B, Salinah O (2014). The Effect of Second-Hand Smoke Exposure during Pregnancy on the Newborn Weight in Malaysia. *The Malaysian Journal of Medical Sciences : MJMS*, 21(2):44–53.

- Onyeizugbo, E. U. (2003). Effects of Gender, Age, and Education on Assertiveness in a Nigerian Sample. *Psychology of Women Quarterly*, 27(1), 12–16.
- Organisation mondiale de la santé. (2017). *WHO report on the global tobacco epidemic, 2017: monitoring tobacco use and prevention policies*.
- Öberg, M., Woodward, A., Jaakkola, M. S., Peruga, A., & Prüss-Ustün, A. (n.d.). Global estimate of the burden of disease from secondhand smoke, 81.
- Passey, M. E., Longman, J. M., Robinson, J., Wiggers, J., & Jones, L. L. (2016). Smoke-free homes: what are the barriers, motivators and enablers? A qualitative systematic review and thematic synthesis. *BMJ Open*, 6(3), e010260.
- Prabandari, Y. S., & Dewi, A. (2016). How do Indonesian youth perceive cigarette advertising? A cross-sectional study among Indonesian high school students. *Global Health Action*, 9(1), 30914.
- Pusat Promosi Kesehatan (2011). *Pedoman Pembangunan Kawasan Tanpa Rokok*. Jakarta: Kementerian Kesehatan Republik Indonesia.
- Rakos, R. F. (1979). Content Consideration in the Distinction between Assertive and Aggressive Behavior. *Psychological Reports*, 44(3), 767–773.
- Samlee Plianbanchang, Ministry of Health, R. of I., CDC Foundation, World Health Organization, & Regional Office for South-East Asia. (2012). *Global adult tobacco survey: Indonesia report 2011*. New Delhi: World Health Organization, Regional Office for Southeast Asia.
- Saw, A., Tang, H., Tsoh, J. Y., Chen, M. S., & Tong, E. K. (2017). Non-smoker assertive behaviour against smoke exposure: Chinese and Korean American non-smokers: Non-smoker assertive behaviour. *Drug and Alcohol Review*, 36(6), 779–787.
- Tedeschji, T., Smith, R. B., & Brown, C. A. (1974). A reinterpretation of research on aggression. *Psychological Bulletin*, 81, 540-562.
- Tobacco Atlas (2019). “Consumption”, <https://tobaccoatlas.org/topic/consumption>, accessed in 15 April 2019 at 15:02.
- Willemsen, M. C., & de Vries, H. (1996). Saying “No” to Environmental Tobacco Smoke: Determinants of Assertiveness among Nonsmoking Employees. *Preventive Medicine*, 25(5), 575–582.
- World Health Organization. (2015). *WHO global report on trends in prevalence of tobacco smoking, 2015*. Retrieved from http://apps.who.int/iris/bitstream/10665/156262/1/9789241564922_eng.pdf
- World Health Organization. (2016). *Prevalence of tobacco smoking: Age-standardized prevalence of current tobacco smoking among persons aged 15 years and older (%), 2015: Male*. [Online] Available at: http://gamapserver.who.int/gho/interactive_charts/tobacco/use/atlas.html [Accessed 1 February 2019]
- World Health Organization. (2018). *World Health Statistics and data visualisation dashboard*. [Online] Available at: <http://apps.who.int/gho/data/node.sdg.3-a-viz?lang=en> [Accessed 1 February 2019]
- World Health Organization. (2019) *WHO Framework convention on tobacco control 2003*. [Online] Available at: <http://whwlibdoc.who.int/publication/2003/9241591013.pdf> [Accessed 13 April 2019]

- World Health Organization. (2017). WHO report on the global tobacco epidemic, 2017. Country profile Indonesia. http://www.who.int/tobacco/surveillance/policy/country_profile/idn.pdf, accessed in 14 April 2019 at 14:07.
- Yang, L., Tong, E. K., Mao, Z., & Hu, T. (2010). Exposure to secondhand smoke and associated factors among non-smoking pregnant women with smoking husbands in Sichuan province, China. *Acta Obstetrica et Gynecologica Scandinavica*, 89(4), 549–557.
- Yerushalmy, J. (2014). The relationship of parents' cigarette smoking to outcome of pregnancy—implications as to the problem of inferring causation from observed associations¹. *International Journal of Epidemiology*, 43(5), 1355–1366.
- Zhang, L., Hsia, J., Tu, X., Xia, Y., Zhang, L., Bi, Z., ... Stanton, B. (2015). Exposure to Secondhand Tobacco Smoke and Interventions Among Pregnant Women in China: A Systematic Review. *Preventing Chronic Disease*, 12.