

DAFTAR PUSTAKA

- Adibah. (2013). Aplikasi Penginderaan Jauh dan Sistem Informasi Geografis untuk Analisis Daerah Resapan Air (Studi kasus:Kota Pekalongan). *Jurnal Geodesi Undip*, 1-13.
- Alberth, D. P. (2005). *Spatial Analysis, GIS and Remote Sensing Applications in The Health Sciences*. Taylor & Francis e-Library.
- Alexandera, M. H. (2003). Spatial Variation of Anopheles Transmitted Wuchereria Bancrofti and Plasmodium Falciparum Infection Densities in Papua New Guinea. *Filaria Journal 2003* , 2:14.
- Amelia, r. (2014). Analisis Faktor Risiko Kejadian Penyakit Filariasis . *Unnes Journal of Public Health Vol.3 No.1*.
- Anand, G. K. (2018). Prediction of land use change based on Land Change Modeler and Attribution of changes in the water balance of Ganga basin to land use change using the SWAT model. *Science of The Total Environment*, 644.
- Anonim. (2015). *Pedoman Pengelolaan Data Penginderaan jauh Landsat 8 untuk Mangrove*. Jakarta: Pusat Pemanfaatan PJ, LAPAN.
- Bao, Q. Z. (2014). NDVI-based Long-Term Vegetation, Dynamics and Its Response to Climatic Change in the Mongolian Plateau. *Remote Sensing ISSN*.
- Beck, L. W. (2000). Remote Sensing and Human Health: New Sensors and New Opportunities. *Emerging Infectious Diseases, California, USA*, Vol.6 No.3.
- Benie, G., Kabore, S., Goita, K., & Courel, M. (2005). *Remote sensing-based spatio-temporal modeling to predict*. Canada: Elsevier.
- Bergquist. (2001). Vector-Borne Parasitic Diseases: New Trends in Data Collection and Risk Assessment. *Acta Tropica 79 (2001)* , 13-20.
- Brugnot. (2008). *Geographical Information System Series*. London: John Wiley & Sons, Inc.
- Campbell, W. (2011). *Introduction to Remote Sensing, Fifth Edition*. New York: Division of Guilford Publications.

- Caprarelli, F. (2013). A Brief review of spatial analysis concepts and tools used for mapping, cantainment and risk modelling of infectious diseases and other illness . *Parasitology (2014), Cambridge University Press 2013*, 141, 581-601.
- Cardo, V. A. (2012). Oviposition stratgies of temporary pool mosquitoes in relation to weather, tidal regime and land use in a temperate wetland. *Bulletin of Entomological Rsearc*, 651-662.
- Catolino, G., Palomba, F., De Lucia, A., & Ferrucci, F. (2019). *Enhancing change prediction models using developer-related factors*. Itali: Elsevier Inc.
- Chandra, G. (2008). Nature Limits Filarial Transmission. *Chandra: Licenses Biomed Central Ltd*.
- Chen, N.-T. S.-B.-Y. (2013). Multi-Decadal Mangrove Forest CHange Detection and Prediction in Honduras, Cenral America, with Landsat Imagery and a Markov Chain Model. *Remote Sensing*, 6408-6426.
- Corner, R. J., M Dewan, A., & Hashizume, M. (2013). *Modelling typhoid risk in Dhaka Metropolitan* (Vol. 12). Perth, Australia: International Journal of Health Geographic. Retrieved from <http://www.ij-healthgeographics.com/content/12/1/13>
- Danoedoro. (2012). *Pengantar Pengindraan Jauh Digital*. Yogyakarta: Andi Offset.
- Doninck V J, B. D. (2014). Modelling the Spatial Distribution of *Culicoides imicola*: Climatic versus Remote Sensing Data. *Remote Sensing*, 6:6604-6619.
- Febrianto, B. (2008). Faktor Risiko Filariasis di Desa Samborejo Kecamatan Tirto kabupaten Pekalongan Jawa Tengah . *Penelitian Vektor dan Reservoir Penyakit, Salatiga. Buletin Penelitian Kesehatan Vol.36 No.2*.
- Fortin, D. (2005). *Spatial ANalysis A Guide for Ecologists*. New York: Published inthe United States of America by Cambridge University Press.
- Gao, B.-C. (1996). *NDWI - A Normalized Difference Water Index from Remote Sensing of Vegetation Liquid Water From Space*. New York, Amerika: Elsevier Science Inc.

- Gomes E, N. O. (2012). Schistosomiasis transmission and environmental change: a spatio temporal analysis in Porto de alihas, Pernambuco Brazil. *International Journal of Health Geographics*, 11:51.
- Gopal, G. a. (2013). *Essentials of Medical Geology : Revised Edition*. Springer Science.
- Guermond. (2008). *The Modelling Process in Geography (form Determinism to Complexity)*. John Wiley & Sons, Inc.Ltd.
- Hamid, A. (2012). Spatial distribution and abundance of culicine mosquitoes in relation to the risk of filariasis transmission in E Sharqiya Governorate.
- Hamid, S. d. (2009). Spatial Distribution and Abundance of culicine mosquitoes in realtion to the risk of filariasis transmission in El Sharqiya Governote, Egypt. *Acad J.Biologi Sci* , (1) : 39-48.
- Hardoyo, S. (2014). *Aspek Sosial Banjir Genangan (ROB) di Kawasan Pesisir*. Yogyakarta: Gadjah Mada University Press.
- Hasan. (1998). Prediction of Villages at Risk for Filariasis Transmission in the Nile Delta Using Remote Sensing and Geography Information System Technologies.
- Hasan, N. K. (2013). Characterization of Landscape features Associated with Mosquito breeding in Urban Cairo using remote sensing, Institute of environmental Studies & Researches, Ain Shams University, Cairo Egypt Stanton . *Baseline drivers of lymphatic Filariasis in Burkina Faso, Geospatial Health*8(1), pg 159-173.
- Huang Y, N. P. (2015). The Prediction of interregional land use differences in Beijing: a Markov Model. *Environment Earth Sci* , 73: 4077-4090.
- Jateng, D. (2013). *Buku Profil Kesehatan Provinsi Jawa Tengah Tahun 2012*. Semarang: Dinas Kesehatan Jawa Tengah.
- Jensen. (2007). *Remote Sensing of the Environment and Earth Resource Prespective*. License of Pearson Education in South Asia: Dorling Kindersley (India) Pvt.Ltd.
- Jensen. (2016). *Introductory Digital Image Processing, A Remote Sensing Perspective 4th Edition*. Pearson Education Inc.

- Jose. (2005). *Spatial Analysis: A Guide for Ecologists*. London: United States of America by Cambridge University.
- Juriastuti. (2010). Faktor Risiko Kejadian Filariasis di Kelurahan Jati Sampurna, Bekasi. *Makara, Kesehatan Vol.14 No.1*, 31-36.
- Kammerer, J. S., Shang, N., Althomsons, S., Haddad, M., Grant, J., & Navin, T. (2013). *Using statistical methods and genotyping to detect tuberculosis outbreaks*. Atlanta, USA: BioMed Central Ltd. .
- Kelly M, B. R. (2011). Terrsetrial Remotely Sensed Imagery in Support of Public Health : New Avenues of Researches using object based image analysis. *Remote sensing*, 2321-2345.
- Klei, T. R., & Rajan, T. (2002). *The Filaria*. USA: Kluwer Academic.
- Lardeux, C. J. (1997). Temperature thresholds and statical modelling of larval *Wucherera bancrofti* developmenta rates. *Parasitology (1997)*, 123-124.
- Lee, D. (2001). *Statistical Analysis with ArcView GIS*. John Wiley & Sons, Inc.
- Lee, D., & Mitchell, R. (2014). *Controlling for localised spatio-temporal autocorrelation in long-term air pollution and health studies*. England: SAGE.
- Leyk, S., McCormick, B., & Nuckols, J. (2011). *Effects of varying temporal scale on spatial models of mortality patterns attributed to pediatric diarrhea*. USA: Elsevier Ltd.
- Liang, S. (2008). *Advance in Land Remote Sensing System, Modelling, Inversion and Application, Springer*.
- Lillesand, K. a. (2004). *Remote Sensing and Image Interpretation*. John Wiley & Sons, Inc. All Right Reserved.
- Liping C, Y. S. (2018). Monitoring and predcting and use and land cover changes using remote sensing and GIS techniques- A case studyof a hilly area, Jianle China. *PLoS ONE:137*, 1-23.
- Liu, H., & Weng, Q. (2011). *Enhancing temporal resolution of satellite imagery for public health studies: A casestudy of West Nile Virus outbreak in Los Angeles in 2007*. Virginia, united states: Elsevier Inc.

- Lunetta dan Lyon, 2004, *Remote Sensing and GIS Accuracy Assessment*, Washington: CFC Press
- Maheswara R, C. M. (2004). *GIS in Public Health Practice*. CRC Press LLC.
- Manheje, M.-P. M. (2013). Socio-environmental variables and transmission risk of lymphatic filariasis in central and northern Mozambique. *Geospatial Health* 7(2), pg 391-398.
- Mardiatno, D., Marfai, M., Rahmawati, K., Tanjung, R., Sianturi, R., & Mutiarni, Y. (2012). *Penilaian Multirisiko Banjir dan Rob di Kecamatan Pekalongan Utara*. Yogyakarta: MPPDAS Program S-2 Geografi, Fakultas Geografi Universitas Gadjah Mada.
- Matějčíček, L., Engst, P., & Jaňour, Z. (2006). A GIS-based approach to spatio-temporal analysis of environmental pollution in urban areas: A case study of Prague's environment extended by LIDAR data. *Ecological Modelling*, 199, 261–277. doi:doi:10.1016/j.ecolmodel.2006.05.018
- McCoy, R. M. (2005). *Field Methods in Remote Sensing*. New York: The Guilford Press.
- Mukhopadhyay A, M. A. (2014). Forest cover change prediction using hybrid methodology of eoinformatic and Markov chain model : A case study on sub Himalayan town Gangtok India. *J.Earth Syst, Sci* 123, 1349-1360.
- Mwase, S. N.-S. (2014). Mapping The Geographical Distribution of Lymphatic Filariasis in Zambia Feb 2014. *PLOS Neglected Tropical Diseases*, Vol.8, Issue 2, e2714.
- Naggar A, E. S. (2013). *GIS application in Prediction of Diseases transmitted by mosquitoes*. Egypt: Suez Canal University.
- Netzband, S. a. (2007). *Applied Remote Sensing for Urban Planning, Governance and Sustainability*. Springer-Verlag Berlin Heidelberg.
- Nouri J, G. A. (2014). Predicting Urban Land Use Changes Using a CA-Markov Model. *Arab J Sci Eng*, 39:5565-5573.
- Oladejo. (2014). Epidemiological study of Tropical Diseases using Geographic Information Systems (GIS) and Remote Sensing (RS). *Sci-Afric Journal of Scientific Issues, Research and Essays Vol 2(5), May 2014*, Pg 238-247.

- Ozturk. (2015). Urban Growth Simulation of Atakum (Samsn,Turkey) Using Celular Automata-Markov Chain and Multi Layer Perceptron-Markov Chain Model. *Remote Sensing*, 7: 5918-5950.
- Palaniyandi. (2014). A Geospatial modelling for Mapping of Filariasis Transmission Risk in India, using remote sensing and GIS . *International Journal of Mosquito Research* , Vol.1 Issue 1.
- Prisbitari, L. D. (2014). *Laporan Kuliah Kerja Lapangan*. Yogyakarta: Graduate of School, Gadjah Mada University.
- Prisbitari, L. D., Nurroh, S., Hadadi, O., Saimu, A., & Saputra, A. (2014). *Laporan Kuliah Kerja Lapangan*. Yogyakarta: Graduate School of Environment Science Magister Program of Environmental Management.
- Putranto, D. (2015). Studi Kerentanan Airtanah Terhadap Pemompaan di Kota Pekalongan, Provinsi Jawa Tengah. *Seminar Nasional Kebumihan ke-8*. Yogyakarta.
- Ratmanov, M. a. (2013). Vector Bone diseases in West Africa: Geographic distribution and Geospatial characteristics. *Royal Society of Tropical Medicine and Hygiene Advance Access* .
- Reddy, A. (2008). *Remote Sensing and Geographical Information Systems*. India: Institute of Science and Technology Jawaharlal Nehru Technological University Kukatpal/y, Hyderabad-72(A.P).
- Rose. (2011). Modelling the impact of climate change on spatial patterns of disease risk: sheep blowfly strike by *Lucilia sericata* in Great Britain. *International Journal for Parasitology*, 739-746.
- Sabesan, R. S. (2013). Lymphatic Filariasis Transmission Risk Map of India. *Based on a Geo Environmental Risk Model, Vector Bone and Zoonatic Diseases*, *Mary Ann Liebert Inc*, Vol.13 No.9 .
- Shen C, L. C. (2016). Spatio-tepora autocorrelation measures for nonstationary series : a new temporaly detrended spatio temporal Moran's index. *Physics Letter A* 380, 106-116.
- Shunlin, L. (2008). System Modelling, Inversion and Application. *Springer Science + Business Media B.V*.

- Simooga, B. V. (2009). Remote Sensing, Geographical Information System and Spatial Analysis for Schistosomiasis Epidemiology and Ecology in Africa, Parasitology. *Cambridge University Press*, 136.
- Skidmore. (2003). *Environmental Modelling with GIS and Remote Sensing*. London EC4P 4EE: Taylor & Francis 11 New Fetter Lane.
- Soemirat. (2010). *Epidemiologi Lingkungan*. Yogyakarta: Gadjah Mada University Press.
- Srividya, M. p. (2002). A Geostatistical analysis of the Geographic Distribution of Lymphatic Filariasis Prevalence In Southern India. *Journal Tropical Medicine and Hygiene by The American Society of Tropical Medicine and Hygiene*, 67(5), pg 480-489.
- Stanton M,. (2013). Baseline Drivers of Lymphatic Filariasis in Burkina Faso. *Geospatial Health* 8(1), 159-173.
- Stein, V. d. (2002). *Spatial Statistics For Remote Sensing, Interbational Institute for Aerospace Survey and Eath Sciences*. The Netherlands: Kluwer Academic Publisher.
- Sugimori. (2004). Remote Sensing, GIS and Public Health. *Remote Sensing and Earth Sciences*, 1:1.
- Sullivan, U. (2010). *Geographic Information Analysis Second Edition*. New Jersey: Published by John Wiley & Sons Inc, Hoboken.
- Sunaryo. (2008). Distribusi Filariasis Limfatik di Kelurahan pabean, Kecamatan Pekalongan Utara Kota Pekalongan Provinsi Jawa Tengah. *Balaba Ed.2007 No.2 Des 2008*, pg 2-6.
- Susilo. (2015). *Pemodelan Spasial Probabilistik ntegrasi Markov Chai dan Cellular Automata untuk Kajian Perubahan Penggunaan Lahan Skala Regional di Provins DIY*. Yogyakarta: Universitas Gadjah Mada.
- Sutanto. (2013). *Metode Penelitian Penginderaan Jauh*. Yogyakarta: Universitas Gadjah Mada.
- Timmreck. (2005). *Epidemiologi Suatu Pengantar* . Jakarta: EGC.
- Tran, G. w. (2002). Mapping Diseases Incidence in Suburban Areas using Remotely Sensed Dara. *American Journal of Epidemilogy Vol.156 No.7*.

- Ugarte, M. D., Adin, A., Goicoa, T., & Militino, A. (2014). *On fitting spatio-temporal disease mapping models using approximate Bayesian inference*. Spain: Sage.
- Utomo, A. H., Mulyaningsih, B., & Hadianto, T. (2002, Januari). Prevalensi Filariasis dan Perilaku Penduduk Desa Rami Pasai dan Betung dalam Pengobatan Filariasis. *Mutiara Medika*, 2.
- Wahyono, T. Y., Purwastyastuti, & Supali, T. (2010). *Filariasis Di Indonesia*. Jakarta: Pusat Data dan Survelans Epidemiologi KemenkesRI.
- Wahyono, T. Y., Purwastyastuti, & Supali, T. (2010). *Filariasis Di Indonesia*. Jakarta: Buletin Jendela Epidemiologi.
- Wang, J.-f., Guo, Y.-S., Christakos, G., Yang, W.-Z., Liao, Y.-L., & Li, Z.-J. (2011). *Hand, foot and mouth disease: spatiotemporal transmission and climate*. Beijing: BioMed Central Ltd.
- Wei, L., Qian, Q., Wang, Z.-Q., & Glass, G. (2011). *Using Geographic Information System-based Ecologic Niche Models to Forecast the Risk of Hantavirus Infection in Shandong Province, China*. Cina: The American Society of Tropical Medicine and Hygiene.
- Wen T, L. N. (2010). Spatial temporal patens of dengue in areas at risk of dengeu hemorrhagic fever in Kaosiung Taiwan 2002. *International Journal of Infectious Diseases*, 334-343.
- Weng. (2010). *Remote Sensing and GIS Integration Theories, Methods and Application* . McGraw-Hill Companies,Inc.
- WHO. (2010). *Lymphatic Filariasis*. Switzerland : WHO Press.
- WHO. (2013). *Lymphatic filariasis: a handbook of Practical Entomology for National Lymphatic Filariasis Elimination Programs*. Switzerland: WHO Press, World Health Organization, 20 Avenue Appia, 1211 Geneva 27.
- Wibowo A, S. J. (2017). Spatial Temporal Analysis of Urban Heat Hazard on Education Area (University of Indonesia). *International Journal of Geography*, 49:1-10.

Widayani. (2016). *Pemodelan Spasial Kerentanan Wilayah Terhadap Penyakit Menular Terkait Lingkungan Bebas Basis Pengindraan Jauh*. Yogyakarta: Gadjah Mada University .

Windiastuti. (2013). Hubungan Kondisi Lingkungan Rumah, Sosial Ekonomi dan Perilaku Masyarakat dengan Kejadian Filariasis di kecamatan Pekalongan Selatan kota Pekalongan. *Jurnal Kesehatan Lingkungan Indonesia* , Vol.12 No.1.