

DAFTAR PUSTAKA

- American Diabetes Association. "2. Classification and diagnosis of diabetes." *Diabetes care* 40.Supplement 1 (2017): S11-S24.
- A. Ghasemi, S. Khalifi, S. J. (2015) 'Streptozotocin-nicotinamide-induced rat model of type 2 diabetes of type 2 diabetes', (December 2014). doi: 10.1556/APhysiol.101.2014.4.2.
- Agus, A. *et al.* (2016) 'Western diet induces a shift in microbiota composition enhancing susceptibility to Adherent-Invasive E. coli infection and intestinal inflammation', *Scientific Reports*. Nature Publishing Group, 6(July 2015), pp. 1–14. doi: 10.1038/srep19032.
- Andersen, A. and Knop, F. K. (2018) 'Glucagon-like peptide 1 in health and disease', 1. doi: 10.1038/s41574-018-0016-2.
- Apolinário, A. C. *et al.* (2014) 'Inulin-type fructans: A review on different aspects of biochemical and pharmaceutical technology', *Carbohydrate Polymers*, 101(1), pp. 368–378. doi: 10.1016/j.carbpol.2013.09.081.
- Athayde, M. L. and Merrill, G. L. (2014) 'Effect of fermented soybean condiment supplemented diet on α -amylase and α -glucosidase activities in Streptozotocin-induced diabetic rats', *Journal of Functional Foods*. Elsevier Ltd, 9, pp. 1–9. doi: 10.1016/j.jff.2014.04.003.
- Aydin, Ö., Nieuwdorp, M. and Gerdes, V. (2018) 'The Gut Microbiome as a Target for the Treatment of Type 2 Diabetes', *Current Diabetes Reports*. Current Diabetes Reports, 18(8). doi: 10.1007/s11892-018-1020-6.
- Banwell, J. G. *et al.* (1983) 'Phytohemagglutinin derived from red kidney bean (*Phaseolus vulgaris*): A cause for intestinal malabsorption associated with bacterial overgrowth in the rat', *Gastroenterology*. Elsevier Masson SAS, 84(3), pp. 506–515. doi: 10.1016/0016-5085(83)90074-4.
- Birt, D. F. *et al.* (2013) 'Resistant starch: promise for improving human health.', *Advances in nutrition (Bethesda, Md.)*, 4(6), pp. 587–601. doi: 10.3945/an.113.004325.
- Canfora, E. E., Jocken, J. W. and Blaak, E. E. (2015) 'Short-chain fatty acids in control of body', *Nature Publishing Group*. Nature Publishing Group. doi: 10.1038/nrendo.2015.128.
- Chambers, E. S. *et al.* (2018) 'Role of Gut Microbiota-Generated Short-Chain Fatty Acids in Metabolic and Cardiovascular Health', *Current Nutrition Reports*. Current Nutrition Reports, 7(4), pp. 198–206. doi: 10.1007/s13668-018-0248-8.
- Degruttola, A. K. *et al.* (2016) 'HHS Public Access', 22(5), pp. 1137–1150. doi: 10.1097/MIB.0000000000000750.Current.
- Duda, A. and Tomasz, C. (2015) 'Interaction of dietary compounds , especially polyphenols , with the intestinal microbiota : a review', pp. 325–341. doi: 10.1007/s00394-015-0852-y.
- Dwivedi, S. *et al.* (2014) 'Plant prebiotics and human health: Biotechnology to breed prebiotic-rich nutritious food crops', *Electronic Journal of Biotechnology*. Elsevier B.V., 17(5), pp. 238–245. doi: 10.1016/j.ejbt.2014.07.004.
- Esquivel-Elizondo, S. *et al.* (2017) 'Insights into Butyrate Production in a Controlled Fermentation System via Gene Predictions', *mSystems*, 2(4), pp.

- 1–13. doi: 10.1128/msystems.00051-17.
- Feng, W., Ao, H. and Pe, C. (2018) 'Gut microbiota, sort-chain fatty aciand herbal medicines', *Frontiers in Pharmacology*, 9(NOV), pp. 1–12. doi: 10.3389/fphar.2018.01354.
- Flamm, G. *et al.* (2001) 'Inulin and oligofructose as dietary fiber: A review of the evidence', *Critical Reviews in Food Science and Nutrition*, 41(5), pp. 353–362. doi: 10.1080/20014091091841.
- Ganesan, K. and Xu, B. (2017) 'Polyphenol-rich dry common beans (*Phaseolus vulgaris* L.) and their health benefits', *International Journal of Molecular Sciences*, 18(11). doi: 10.3390/ijms18112331.
- González-Gallego, J. *et al.* (2013) 'Anti-Inflammatory and Immunomodulatory Properties of Dietary Flavonoids', *Polyphenols in Human Health and Disease*, 1(II), pp. 435–452. doi: 10.1016/B978-0-12-398456-2.00032-3.
- Hasim *et al.* (2015) 'Bacillus subtilis natto fermentation to improve aglycone isoflavones content of black soybean varieties detam 2', *International Food Research Journal*, 22(6), pp. 2558–2564.
- Hidalgo, M. *et al.* (2012) 'Metabolism of anthocyanins by human gut microflora and their influence on gut bacterial growth', *Journal of Agricultural and Food Chemistry*, 60(15), pp. 3882–3890. doi: 10.1021/jf3002153.
- Hu, Y. *et al.* (2010) 'Characterization of fermented black soybean natto inoculated with Bacillus natto during fermentation', *Journal of the Science of Food and Agriculture*, 90(7), pp. 1194–1202. doi: 10.1002/jsfa.3947.
- Karakaya, S. (2010) 'Changes in Total Phenols , Total Flavonoids , and Antioxidant Activities of Common Beans and Pinto Beans after Soaking , Cooking , and', (June). doi: 10.1007/s10068-010-0089-8.
- Kim, Y. A., Keogh, J. B. and Clifton, P. M. (2017) 'Probiotics, prebiotics, synbiotics and insulin sensitivity', *Nutrition Research Reviews*, (2018), pp. 1–17. doi: 10.1017/S095442241700018X.
- Kiss, A. and Forgo, P. (2011) 'Investigations on inulin-type oligosaccharides with regard to HPLC analysis and prospective food applicability', *Monatshefte fur Chemie*, 142(6), pp. 547–553. doi: 10.1007/s00706-011-0485-7.
- Kriukova, Y. *et al.* (2018) 'Chain length distribution of inulin from dahlia tubers as influenced by the extraction method', *International Journal of Food Properties*. Taylor & Francis, 20(3), pp. 3112–3122. doi: 10.1080/10942912.2017.1357043.
- Kumar, S. *et al.* (2013) 'Clinical complications of kidney bean (*Phaseolus vulgaris* L.) consumption', *Nutrition*. Elsevier Inc., 29(6), pp. 821–827. doi: 10.1016/j.nut.2012.11.010.
- Kusmiyati, N., Wahyuningsih, T. D. and Widodo (2018) 'Prebiotic effect of inulin extract from dahlia tubers (*Dahlia pinnata* L.) on the growth performance of intestinal-origin Lactobacillus casei AP', *Pakistan Journal of Nutrition*, 17(8), pp. 405–410. doi: 10.3923/pjn.2018.405.410.
- Li, M. *et al.* (2018) 'Pro- and anti-inflammatory effects of short chain fatty acids on immune and endothelial cells', *European Journal of Pharmacology*, 831(May), pp. 52–59. doi: 10.1016/j.ejphar.2018.05.003.
- Limón, R. I. *et al.* (2015) 'Fermentation enhances the content of bioactive compounds in kidney bean extracts', *Food Chemistry*. Elsevier Ltd, 172, pp. 343–352. doi: 10.1016/j.foodchem.2014.09.084.
- Lingyun, W. *et al.* (2007) 'Studies on the extracting technical conditions of inulin

- from Jerusalem artichoke tubers', *Journal of Food Engineering*, 79(3), pp. 1087–1093. doi: 10.1016/j.jfoodeng.2006.03.028.
- Louis, P. and Flint, H. J. (2017) 'Formation of propionate and butyrate by the human colonic microbiota', *Environmental Microbiology*, 19(1), pp. 29–41. doi: 10.1111/1462-2920.13589.
- Luca, C. De and Olefsky, J. M. (2009) 'NIH Public Access', 582(1), pp. 97–105.
- Macy, J. M., Ljungdahl, L. G. and Gottschalk, G. (1978) 'Pathway of succinate and propionate formation in *Bacteroides fragilis*', *Journal of Bacteriology*, 134(1), pp. 84–91.
- Melanie, H. *et al.* (2015) 'Characterization of Inulin from Local Red Dahlia (*Dahlia* sp. L) Tubers by Infrared Spectroscopy', *Procedia Chemistry*, 16, pp. 78–84. doi: 10.1016/j.proche.2015.12.027.
- Mensink, M. A. *et al.* (2015) 'Inulin, a flexible oligosaccharide. II: Review of its pharmaceutical applications-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>)', *Carbohydrate Polymers*. Elsevier Ltd., 134, pp. 418–428. doi: 10.1016/j.carbpol.2015.08.022.
- Morrison, D. J. and Preston, T. (2016) 'Formation of short chain fatty acids by the gut microbiota and their impact on human metabolism', *Gut Microbes*. Taylor & Francis, 7(3), pp. 189–200. doi: 10.1080/19490976.2015.1134082.
- Nair, A. B. and Jacob, S. (2016) 'A simple practice guide for dose conversion between animals and human.', *Journal of basic and clinical pharmacy*, 7(2), pp. 27–31. doi: 10.4103/0976-0105.177703.
- Naito, Y., Uchiyama, K. and Takagi, T. (2018) 'Dual antiplatelet therapy does not affect the Society for Free Radical Research Japan 10.3164/j6cb .18-16 1880 50860912-0009 JJCBN Original Article Kyj bn18-1 c oto, Japan ournal of Clinical Biochemistry and Nutrition the incidence of low?dose aspirin?ind', *Journal of Clinical Biochemistry and Nutrition*, 63(June), pp. 2016–2019. doi: 10.3164/jcbn.18.
- Nciri, N. and Cho, N. (2018) 'New research highlights: Impact of chronic ingestion of white kidney beans (*Phaseolus vulgaris* L. var. Beldia) on small-intestinal disaccharidase activity in Wistar rats', *Toxicology Reports*. Elsevier, 5(December 2017), pp. 46–55. doi: 10.1016/j.toxrep.2017.12.016.
- Neilson, A. P., Goodrich, K. M. and Ferruzzi, M. G. (2017) *Bioavailability and Metabolism of Bioactive Compounds From Foods*. Fourth Edi, *Nutrition in the Prevention and Treatment of Disease*. Fourth Edi. Elsevier Inc. doi: 10.1016/B978-0-12-802928-2.00015-1.
- Ohira, H., Tsutsui, W. and Fujioka, Y. (2017) 'Are Short Chain Fatty Acids in Gut Microbiota Defensive Players for Inflammation and Atherosclerosis?', *Journal of Atherosclerosis and Thrombosis*, 24(7), pp. 660–672. doi: 10.5551/jat.RV17006.
- Ozidal, T. *et al.* (2016) 'The reciprocal interactions between polyphenols and gut microbiota and effects on bioaccessibility', *Nutrients*, 8(2), pp. 1–36. doi: 10.3390/nu8020078.
- Preti, R., Rapa, M. and Vinci, G. (2017) ' Effect of Steaming and Boiling on the Antioxidant Properties and Biogenic Amines Content in Green Bean (*Phaseolus vulgaris*) Varieties of Different Colours ', *Journal of Food Quality*, 2017, pp. 1–8. doi: 10.1155/2017/5329070.
- Puddu, A. *et al.* (2014) 'Evidence for the Gut Microbiota Short-Chain Fatty Acids as Key Pathophysiological Molecules Improving Diabetes', *Mediators of*

- Inflammation*, 2014, pp. 1–9. doi: 10.1155/2014/162021.
- Q. Wei, C. W. and K. C. C. (2001) 'Natto Characteristics as Affected by Steaming Time , Bacillus Strain , and Fermentation Time', pp. 167–173.
- Rivière, A. *et al.* (2016) 'Bifidobacteria and Butyrate-Producing Colon Bacteria : Importance and Strategies for Their Stimulation in the Human Gut', 7(June). doi: 10.3389/fmicb.2016.00979.
- Röhrborn, D., Wronkowitz, N. and Eckel, J. (2015) 'DPP4 in diabetes', 6(July), pp. 1–20. doi: 10.3389/fimmu.2015.00386.
- Saltzman, E. T. *et al.* (2018) 'Intestinal Microbiome Shifts , Dysbiosis , Inflammation , and Non-alcoholic Fatty Liver Disease', 9(January), pp. 1–11. doi: 10.3389/fmicb.2018.00061.
- Savage, G. P. (2004) 'Hemagglutinins (Haemagglutinins)', *Encyclopedia of Food Sciences and Nutrition*, pp. 3057–3060. doi: 10.1016/b0-12-227055-x/00935-4.
- Shrestha, A. K., Dahal, N. R. A. J. and Ndungutse, V. (2010) 'Bacillus Fermentation of Soybean : A Review', 6, pp. 1–9.
- Siah, S. *et al.* (2014) 'Effects of soaking, boiling and autoclaving on the phenolic contents and antioxidant activities of faba beans (*vicia faba* L.) differing in seed coat colours', *Food Chemistry*. Elsevier Ltd, 142, pp. 461–468. doi: 10.1016/j.foodchem.2013.07.068.
- Singh, A. *et al.* (2018) 'Inulin fiber dose-dependently modulates energy balance, glucose tolerance, gut microbiota, hormones and diet preference in high-fat-fed male rats', *Journal of Nutritional Biochemistry*. Elsevier Inc, 59, pp. 142–152. doi: 10.1016/j.jnutbio.2018.05.017.
- Sugiharto, S. *et al.* (2018) 'Effect of two-step fermentation by *Chrysonilia crassa* and *Bacillus subtilis* on nutritional values and antioxidative properties of agro-industrial by-products as poultry feed ingredients', 7710(December), pp. 472–480.
- Tamaroh, S. *et al.* (2018) 'Perubahan Antosianin dan Aktivitas Antioksidan Tepung Uwi Ungu selama Penyimpanan', *Jurnal Aplikasi Teknologi Pangan*, 7(1), pp. 31–36. doi: 10.17728/jatp.2224.
- Thursby, E. and Juge, N. (2017) 'Introduction to the human gut microbiota.', *The Biochemical journal*, 474(11), pp. 1823–1836. doi: 10.1042/BCJ20160510.
- De Vuyst, L. *et al.* (2014) 'Summer Meeting 2013: Growth and physiology of bifidobacteria', *Journal of Applied Microbiology*, 116(3), pp. 477–491. doi: 10.1111/jam.12415.
- Wang, Jun *et al.* (2012) 'A metagenome-wide association study of gut microbiota in type 2 diabetes', *Nature*. Nature Publishing Group, 490(7418), pp. 55–60. doi: 10.1038/nature11450.
- Wassenaar, T. M. and Zimmermann, K. (2018) 'Lipopolysaccharides in food, food supplements, and probiotics: should we be worried?', *European Journal of Microbiology and Immunology*, 8(3), pp. 63–69. doi: 10.1556/1886.2018.00017.
- Weitkunat, K. *et al.* (2015) 'ScienceDirect Effects of dietary inulin on bacterial growth , short-chain fatty acid production and hepatic lipid metabolism in gnotobiotic mice ☆', *The Journal of Nutritional Biochemistry*. Elsevier Inc., 26(9), pp. 929–937. doi: 10.1016/j.jnutbio.2015.03.010.
- Wilson, B. and Whelan, K. (2017) 'Prebiotic inulin-type fructans and galactooligosaccharides: definition, specificity, function, and application in gastrointestinal disorders', *Journal of Gastroenterology and Hepatology*

- (Australia), 32, pp. 64–68. doi: 10.1111/jgh.13700.
- Worku, A. and Sahu, O. (2017) 'Significance of fermentation process on biochemical properties of Phaseolus vulgaris (red beans)', *Biotechnology Reports*. Elsevier, 16(April), pp. 5–11. doi: 10.1016/j.btre.2017.09.001.
- Yen, G., Chen, H. and Science, F. (1995) 'Antioxidant Activity', pp. 27–32.
- Yoon, H. J. *et al.* (2015) 'Effect of fermentation by Bacillus subtilis on antioxidant and cytotoxic activities of black rice bran', *International Journal of Food Science and Technology*, 50(3), pp. 612–618. doi: 10.1111/ijfs.12693.
- Zhang, C. *et al.* (2014) 'Cooked navy and black bean diets improve biomarkers of colon health and reduce inflammation during colitis', pp. 1549–1563. doi: 10.1017/S0007114513004352.
- Zhang, Q. *et al.* (2018) 'Inulin-type fructan improves diabetic phenotype and gut microbiota profiles in rats', pp. 1–24. doi: 10.7717/peerj.4446.
- Zhang, X. *et al.* (2016) 'The Modulatory Effect of Anthocyanins from Purple Sweet Potato on Human Intestinal Microbiota in Vitro', *Journal of Agricultural and Food Chemistry*, 64(12), pp. 2582–2590. doi: 10.1021/acs.jafc.6b00586.
- Zubaidah, E. and Akhadiana, W. (2013) 'Comparative Study of Inulin Extracts from Dahlia, Yam, and Gembili Tubers as Prebiotic', 2013(November), pp. 8–12.