

BIBLIOGRAPHY

- Abhary, A. and Hassani, H. (2016) 'Using PCA and BR methods on Landsat- 8 OLI Images for Mapping Hydrothermal Altered Mineral Deposits in Baft 1 : 100 , 000 Geological', (8), pp. 1433–1438.
- Abrams, M. and Hook, S. (2002) 'ASTER User Handbook Version 2', *Jet Propulsion*, 2003(23/09/2003), p. 135.
- Arribas, A. (1995) 'Characteristics of high-sulfidation epithermal deposits, and their relation to magmatic fluid', *Mineralogical Association of Canada Short Course Series*, 23, pp. 419–454.
- Ashley, R. P. (1982) 'Occurrence model for Enargite-gold Deposit', pp. 144–147.
- Berlo, K. Van Hinsberg, V.J, Vigouroux. N Gagnon J.E and Williams-Jones, A.E. (2014) 'Sulfide breakdown controls metal signature in volcanic gas at Kawah Ijen volcano, Indonesia', *Chemical Geology*. Elsevier B.V., 371, pp. 115–127.
- Caudron, C. Kamil,D. and Lecocq,T (2015) 'Kawah Ijen volcanic activity : a review'.
- Caudron, C. Mauri.G and Williams-jones,G(2016) 'New insights into the Kawah Ijen hydrothermal system from geophysical data New insights into the Kawah Ijen hydrothermal system from geophysical data ' as *Renovables* , Universidad Nacional Auto', (January).
- Clark, D. A. (2014) 'Magnetic effects of hydrothermal alteration in porphyry copper and iron-oxide copper-gold systems: A review', *Tectonophysics*. Elsevier B.V., 624–625, pp. 46–65.
- Corbett, G. . (2002) 'Epithermal Gold for Explorationists', *AIG Journal, Applied*

Geoscientific Practice and Research in Australia, (February), p. 26 p.

Ducart, D. F. Silva, A. and Toledo, C. (2016) 'Mapping iron oxides with Landsat-8/OLI and EO-1/Hyperion imagery from the Serra Norte iron deposits in the Carajás Mineral Province, Brazil', *Brazilian Journal of Geology*, 46(3), pp. 331–349.

Estornell, J. Cro, A. and Azevedo, F. (2015) 'Principal component analysis applied to remote sensing', *Modelling in Science Education and Learning*, 6(7), p. 83.

Fatima, K. (2017) 'Minerals identification and mapping using ASTER satellite image Khunsa Fatima', (October).

Filho, C. R. D. E. S.. (2003) 'Targeting key alteration minerals in epithermal deposits in Patagonia, Argentina, using ASTER imagery and principal component analysis'.

Frutuoso, R. . da C. (2015) 'Mapping hydrothermal gold mineralization using Landsat 8 data. A case of study in Chaves license, Portugal'.

Hamilton, W. (1979) 'Tectonics of the Indonesian Region Tectonics of the Indonesian Region', *U.S. GEOLOGICAL SURVEY PROFESSIONAL PAPER*, 1078, pp. 1–65.

Harrison, R. (2012) 'The Geology, Alteration and Mineralisation of the Tumpangpitu Porphyry Cu-Au and High-Sulfidation Epithermal Au-Ag Deposit', (November), pp. 273–281.

Harrison, R. L. Maryono, A., and Norris, M. (2018) 'Geochronology of the Tumpangpitu Porphyry Au-Cu-Mo and High-Sulfidation Epithermal Au-Ag-Cu Deposit: Evidence for Pre- and Postmineralization Diatremes in

- the Tujuh Bukit District , Southeast Java , Indonesia *’, pp. 163–192.
- Heald, P., Foley, N. K. and Hayba, D. O. (1987) ‘Comparative anatomy of volcanic-hosted epithermal deposits: acid- sulfate and adularia- sericite types.’, *Economic Geology*, 82(1), pp. 1–26.
- Hedenquist, J. W., Arribas, A. and Reynolds, T. J. (1998) ‘Evolution of an intrusion-centered hydrothermal system: far southeast-Lepanto porphyry and epithermal Cu-Au deposits, Philippines’, *Economic Geology*, 93(4), pp. 373–404.
- Hedenuist, N. C. W. and J. W. (1995) ‘Epithermal Gold Deposits : Styles , Characteristics and Exploration’, *SEG Newsletter*, (23), pp. 9–13.
- van Hinsberg, V. Berlo.K, Van Bergen.M. (2010) ‘Extreme alteration by hyperacidic brines at Kawah Ijen volcano, East Java, Indonesia: I. Textural and mineralogical imprint’, *Journal of Volcanology and Geothermal Research*. Elsevier B.V., 198(1–2), pp. 253–263.
- Jeffery, W. H. (2000) ‘Exploration-for-Epithermal-Gold-Deposits.pdf’, *SEG Review*. Ontario, pp. 245–277.
- Jing, C. Bokun,Y. Runsheng,W. and Feng.T. (2014) ‘Regional-scale mineral mapping using ASTER VNIR/SWIR data and validation of reflectance and mineral map products using airborne hyperspectral CASI/SASI data’, *International Journal of Applied Earth Observation and Geoinformation*. Elsevier B.V., 33(1), pp. 127–141.
- Kanlinowsk I, A. and Oliver, S. (2004) ‘ASTER Mineral Index Processing Manual. Remote Sensing Application Geoscience Australia.’, (October).
- L.Hellman, P. (2012) ‘Tujuh Bukit Project Report on Mineral Resource’.

- Loughlin, W. P. (1991) 'Principal Component Analysis for Alteration Mapping *', (April 1985).
- Mahanta, P. and Maiti, S. (2018a) 'Regional scale demarcation of alteration zone using ASTER imageries in South Purulia Shear Zone, East India: Implication for mineral exploration in vegetated regions', *Ore Geology Reviews*. Elsevier B.V.
- Mahanta, P. and Maiti, S. (2018b) 'Regional scale demarcation of alteration zone using ASTER imageries in South Purulia Shear Zone , East India : Implication for mineral exploration in vegetated regions', *Ore Geology Reviews*. Elsevier, 102(July), pp. 846–861.
- Manuel, R. Brito.M and Chichorro.M. (2017) 'Remote Sensing for Mineral Exploration in Central Portugal', *Minerals*, 7(10), p. 184. doi: 10.3390/min7100184.
- Myaing, Y. Y., Idrus, A. and Titisari, A. D. (2018) 'Fluid Inclusion Study of The Tumpangpitu High Sulfidation Epithermal Gold Deposit in Banyuwangi District, East Java, Indonesia', *Journal of Geoscience, Engineering, Environment, and Technology*, 3(1), p. 8.
- Nishiyama, T. (2005) 'The roles of Asia and Chile in the world copper market', *Resources Policy*, 30(2), pp. 131–139.
- Pour, A. B. and Hashim, M. (2012) 'The application of ASTER remote sensing data to porphyry copper and epithermal gold deposits', *Ore Geology Reviews*. Elsevier B.V., 44, pp. 1–9.
- Pour, A. B. and Hashim, M. (2015) 'Hydrothermal alteration mapping from Landsat-8 data, Sar Cheshmeh copper mining district, south-eastern

Islamic Republic of Iran’, *Journal of Taibah University for Science*.
Taibah University, 9(2), pp. 155–166.

Sabins, F. F. (1999) ‘Remote sensing for mineral exploration’.

San, B., Sumer, O. and Gurcay, B. (2004) ‘Comparison of band rationing and spectral indices methods for detecting alunite and kaolinite minerals using Aster data in Biga region, Turkey’, *Proceedings ISPRS*. Available at: <http://cartesia.org/geodoc/isprs2004/comm7/papers/15.pdf>.

Scher, S., Williams-Jones, A. E. and Williams-Jones, G. (2013) ‘Fumarolic activity, acid-sulfate alteration, and high sulfidation epithermal precious metal mineralization in the crater of Kawah Ijen volcano, Java, Indonesia’, *Economic Geology*, 108(5), pp. 1099–1118.

Setijadji, L. D., Kanjino, S., Imai, A. and Watanabe, K. (2006) ‘Cenozoic island arc magmatism in Java Island (Sunda Arc, Indonesia): Clues on relationships between geodynamics of volcanic centers and ore mineralization’, *Resource Geology*, 56(3), pp. 267–292.

Setijadji, L. D. and Maryono, A. (2012) ‘Geology and Arc Magmatism of the Eastern Sunda Arc, Indonesia’, (November).

Sillitoe, R. H. (1987) ‘Styles of High-Sulphidation Gold, Silver and Copper Mineralisation in Porphyry and Epithermal Environments SOME GENERAL FEATURES OF HS SYSTEMS’.

Steven M., D. J. and Clever, J. G. P. . (2006) ‘Remote sensing image analysis-including the spatial domain’, *The Photogrammetric Record*, 21(113), pp. 84–85. doi: 10.1111/j.1477-9730.2006.00359_4.x.

Thompson, A. J. B. and Thompson, A. J. H. (1996) ‘Atlas of Alteration. A field and

petrographic guide to hydrothermal alteration minerals’, p. 119.

- Di Tommaso, I. and Rubinstein, N. (2007) ‘Hydrothermal alteration mapping using ASTER data in the Infiernillo porphyry deposit, Argentina’, *Ore Geology Reviews*, 32(1–2), pp. 275–290.
- Wang, L. Zhong, J. and Pirajno, F. (2019) ‘A review of intermediate sulfidation epithermal deposits and subclassification’, *Ore Geology Reviews*. Elsevier, 107(19), pp. 434–456.
- Zhong, J., Pirajno, F. and Chen, Y. J. (2017) ‘Epithermal deposits in South China: Geology, geochemistry, geochronology and tectonic setting’, *Gondwana Research*. Elsevier B.V., 42, pp. 193–219.