

DAFTAR PUSTAKA

- Allin, K. H., & Nordestgaard, B. G. (2011). Elevated C-reactive protein in the diagnosis, prognosis, and cause of cancer. *Critical Reviews in Clinical Laboratory Sciences*, 48(4), 155–170.
<https://doi.org/10.3109/10408363.2011.599831>
- April, M. D., Aguirre, J., Tannenbaum, L. I., Moore, T., Pingree, A., Thaxton, R. E., ... Lantry, J. H. (2017). Sepsis Clinical Criteria in Emergency Department Patients Admitted to an Intensive Care Unit: An External Validation Study of Quick Sequential Organ Failure Assessment. *The Journal of Emergency Medicine*, 52(5), 622–631.
<https://doi.org/10.1016/j.jemermed.2016.10.012>
- Arif, S. K., Suyata, M. P., Gaus, S., & Ahmad, M. R. (2017). Procalcitonin and C-Reactive Protein as a Predictor of Organ Dysfunction and Outcome of Sepsis and Septic Shock Patients in Intensive Care Unit. *Global Journal of Health Science*, 9(12), 169. <https://doi.org/10.5539/gjhs.v9n12p169>
- Bali, R., Sharma, P., Ghanghas, P., Gupta, N., Tiwari, J. D., Singh, A., ... Goyal, D. (2017). To Compare the Efficacy of C-Reactive Protein and Total Leucocyte Count as Markers for Monitoring the Course of Odontogenic Space Infections. *Journal of Maxillofacial and Oral Surgery*, 16(3), 322–327. <https://doi.org/10.1007/s12663-016-0978-3>
- Candy, Heber B. Sapan, Laurens T. B. Kalesaran, & Ferry Kalitouw. (2017). Besaran Neutrofil dan Kadar C-reactive Protein sebagai Faktor Prognostik Multi Organ Failure pada Pasien Multi-trauma. *Jurnal Biomedik (JBM)*, Volume 9, 184–190.
- Clyne, B., & Olshaker, J. S. (1999). The C-reactive protein. *The Journal of Emergency Medicine*, 17(6), 1019–1025.
- Darmawan, I. (2008). *Update on SEPSIS* (Pertama). Jakarta: Farmedia.
- Dombrovskiy, V. Y., Martin, A. A., Sunderram, J., & Paz, H. L. (2007). Rapid increase in hospitalization and mortality rates for severe sepsis in the United States: A trend analysis from 1993 to 2003*: *Critical Care Medicine*, 35(5), 1244–1250.
<https://doi.org/10.1097/01.CCM.0000261890.41311.E9>
- Gogos, C. A., Giali, S., Paliogianni, F., Dimitracopoulos, G., Bassaris, H. P., & Vagenakis, A. G. (2001). Interleukin-6 and C-reactive protein as early markers of sepsis in patients with diabetic ketoacidosis or hyperosmosis. *Diabetologia*, 44(8), 1011–1014. <https://doi.org/10.1007/s001250100592>
- Gonzalez, M. A., & Ochoa, C. D. (2018). Multiorgan System Failure in Sepsis. Dalam G. Ortiz-Ruiz & C. Dueñas-Castell (Ed.), *Sepsis* (hlm. 67–71). New York, NY: Springer New York. https://doi.org/10.1007/978-1-4939-7334-7_6

- Henny Rosita Iskandar, Dharma Mulyo, Antonius Pudjiadi, A. Pratiwi, & Y. Suryatin. (2010). Sensitifitas dan Spesifisitas Pemeriksaan Procalcitonin, C-Reactive Protein (CRP), dan Hitung Leukosit untuk Memprediksi Infeksi Bakterial pada Sindrom Syok Dengue di Pediatric Intensive Care Unit. *Sari Pediatri*, *Vol. 12, No. 4, Desember 2010*, 233–240.
- Jeschke, M. G., Finnerty, C. C., Kulp, G. A., Kraft, R., & Herndon, D. N. (2013). Can we use C-reactive protein levels to predict severe infection or sepsis in severely burned patients? *International Journal of Burns and Trauma*, *3*(3), 137–143.
- Komiya, K., Ishii, H., Teramoto, S., Takahashi, O., Yamamoto, H., Oka, H., ... Kadota, J. (2012). Plasma C-reactive protein levels are associated with mortality in elderly with acute lung injury. *Journal of Critical Care*, *27*(5), 524.e1-524.e6. <https://doi.org/10.1016/j.jcrc.2011.10.005>
- Lanziotti, V. S., Póvoa, P., Soares, M., Silva, J. R. L. e, Barbosa, A. P., & Salluh, J. I. F. (2016). Use of biomarkers in pediatric sepsis: literature review. *Revista Brasileira de Terapia Intensiva*, *28*(4). <https://doi.org/10.5935/0103-507X.20160080>
- Lobo, S. M. A., Lobo, F. R. M., Bota, D. P., Lopes-Ferreira, F., Soliman, H. M., Meélot, C., & Vincent, J.-L. (2003). C-Reactive Protein Levels Correlate With Mortality and Organ Failure in Critically Ill Patients. *Chest*, *123*(6), 2043–2049. <https://doi.org/10.1378/chest.123.6.2043>
- Luzzani, A., Polati, E., Dorizzi, R., Rungtatscher, A., Pavan, R., & Merlini, A. (2003). Comparison of procalcitonin and C-reactive protein as markers of sepsis: *Critical Care Medicine*, *31*(6), 1737–1741. <https://doi.org/10.1097/01.CCM.0000063440.19188.ED>
- M Dilip Kumar, M R Renuka Devi, & P Sai Kumar. (2012). High Sensitive C - reactive protein – A Risk Marker For Coronary Artery Disease. *IOSR Journal of Dental and Medical Sciences (JDMS)*, *2*(Nov.-Dec. 2012), 28–32.
- Ma, L., Liu, X., Lu, Z., Wu, L., Tang, Y., Luo, X., ... Xie, Y. (2017). Assessment of high-sensitivity C-reactive protein tests for the diagnosis of hepatocellular carcinoma in patients with hepatitis B-associated liver cirrhosis. *Oncology Letters*. <https://doi.org/10.3892/ol.2017.5890>
- Marshall, J. C., & Reinhart, K. (2009). Biomarkers of sepsis: *Critical Care Medicine*, *37*(7), 2290–2298. <https://doi.org/10.1097/CCM.0b013e3181a02afc>
- Meisner, M., Tschaikowsky, K., Palmaers, T., & Schmidt, J. (1999). Comparison of procalcitonin (PCT) and C-reactive protein (CRP) plasma concentrations at different SOFA scores during the course of sepsis and MODS. *Critical Care*, *3*(1), 45. <https://doi.org/10.1186/cc306>
- Miguel-Bayarri, V., Casanoves-Laparra, E. B., Pallás-Beneyto, L., Sancho-Chinesta, S., Martín-Osorio, L. F., Tormo-Calandín, C., & Bautista-Rentero, D. (2012). Prognostic value of the biomarkers procalcitonin, interleukin-6 and C-reactive protein in severe sepsis. *Medicina Intensiva (English Edition)*, *36*(8), 556–562. <https://doi.org/10.1016/j.medine.2012.10.008>

- Minasyan, H. (2017). Sepsis and septic shock: Pathogenesis and treatment perspectives. *Journal of Critical Care*, *40*, 229–242. <https://doi.org/10.1016/j.jcrc.2017.04.015>
- Pedro Póvoa, Luís Coelho, Eduardo Almeida, Antero Fernandes, Rui Mealha, Pedro Moreira, & Henrique Sabino. (2006). Early identification of intensive care unit-acquired infections with daily monitoring of C-reactive protein: a prospective observational study. <https://doi.org/10.1186/cc4892>
- Pierrakos, C., & Vincent, J.-L. (2010). Sepsis biomarkers: a review. *Critical Care*, *14*(1), R15. <https://doi.org/10.1186/cc8872>
- Póvoa, P. (2002). C-reactive protein: a valuable marker of sepsis. *Intensive Care Medicine*, *28*(3), 235–243. <https://doi.org/10.1007/s00134-002-1209-6>
- Ryu, J.-A., Yang, J. H., Lee, D., Park, C.-M., Suh, G. Y., Jeon, K., ... Chung, C. R. (2015). Clinical Usefulness of Procalcitonin and C-Reactive Protein as Outcome Predictors in Critically Ill Patients with Severe Sepsis and Septic Shock. *PLOS ONE*, *10*(9), e0138150. <https://doi.org/10.1371/journal.pone.0138150>
- Schmit, X., & Vincent, J. L. (2008). The Time Course of Blood C-reactive Protein Concentrations in Relation to the Response to Initial Antimicrobial Therapy in Patients with Sepsis. *Infection*, *36*(3), 213–219. <https://doi.org/10.1007/s15010-007-7077-9>
- Simon, L., Gauvin, F., Amre, D. K., Saint-Louis, P., & Lacroix, J. (2004). Serum Procalcitonin and C-Reactive Protein Levels as Markers of Bacterial Infection: A Systematic Review and Meta-analysis. *Clinical Infectious Diseases*, *39*(2), 206–217. <https://doi.org/10.1086/421997>
- Singer, M., Deutschman, C. S., Seymour, C. W., Shankar-Hari, M., Annane, D., Bauer, M., ... Angus, D. C. (2016). The Third International Consensus Definitions for Sepsis and Septic Shock (Sepsis-3). *JAMA*, *315*(8), 801. <https://doi.org/10.1001/jama.2016.0287>