

DAFTAR PUSTAKA

1. Vialle R, Levassor N, Rillardon L (2005) Radiographic analysis of the sagittal alignment and balance of the spine in asymptomatic subjects. *J Bone Joint Surg Am* 87, 260–267. [PubMed]
2. Lee CS, Chung SS, Kang KC, Park SJ, Shin SK (2011) Normal patterns of sagittal alignment of the spine in young adults radiological analysis in a Korean population. *Spine* 36, E1648–E1654. [PubMed]
3. Pratali RR, Luz CO, Barsotti CG, Santos FE, Oliveira CS (2014) Analysis of sagittal balance and spinopelvic parameters in a Brazilian population sample. *Coluna/Columna* 13, 108–111.
4. Vrtovec T, Janssen M, Likar B, et al. (2012) A review of methods for evaluating the quantitative parameters of sagittal pelvic alignment. *Spine J* 12, 433–446. [PubMed]
5. Itio E (1991) Roentgenographic analysis of posture in spinal osteoporotics. *Spine* 16, 750–756. [PubMed]
6. Kim WJ, Kang JW, Yeom JS, et al. (2003) A comparative analysis of sagittal spinal balance in 100 asymptomatic young and older aged volunteers. *J Korean Soc Spine Surg* 10, 327–334.
7. Tomaž Vrtovec, Franjo Pernuš, and Boštjan Likar. Determination of axial vertebral rotation in MR images: comparison of four manual and a computerized method. *European Spine Journal*, 19 (5) : 774–781, 2010



8. Tomaž Vrtovec, Boštjan Likar, and Franjo Pernuš. Determination of 3D location and rotation of lumbar vertebrae in CT images by symmetry-based auto-registration. SPIE Medical Imaging 2007, Image Processing, 17-22 Feb, San Diego, CA, USA, J.P.W. Pluim, J.M. Reinhardt (Eds.), Proceedings of SPIE, vol. 6512 : 65121Q, 2007.
9. Tomaž Vrtovec. Modality-independent determination of vertebral position and rotation in 3D. 4th International Workshop on Medical Imaging and Augmented Reality – MIAR 2008, 1-2 Aug, Tokyo, Japan, T. Dohi, I. Sakuma, H. Liao (Eds.), Springer, Lecture Notes in Computer Science (LNCS), vol. 5128 : 89–97, 2008
10. Nicolai Bodguk, Clinical Anatomy of the Lumbar Spine and Sacrum, Fourth Edition, Elsevier, 2005
11. Nojiri K, Matsumoto M, Chiba K, Toyama Y. Morphometric analysis of the thoracic and lumbar spine in Japanese on the use of pediculus screws. Surg Radiol Anat (2005) 27: 123–128
12. Boulay et al., Sagittal alignment of spine and pelvis regulated by pelvic incidence: standard values and prediction of lordosis. Eur Spine J. 2006 Apr;15(4):415-22. Epub 2005 Sep 23.
13. Boulay et al. Pelvic Incidence: A Predictive Factor for Three-Dimensional Acetabular Orientation—A Preliminary Study. Anat Res Int. 2014; 2014: 594650
14. Lee CS, Chung SS, Kang KC, Park SJ, Shin SK (2011) Normal patterns of sagittal alignment of the spine in young adults radiological analysis in a Korean population. Spine 36, E1648–E1654.



15. Pratali RR, Luz CO, Barsotti CG, Santos FE, Oliveira CS (2014) Analysis of sagittal balance and spinopelvic parameters in a Brazilian population sample. *Coluna/Columna* 13, 108–111.
16. Vrtovec T, Janssen M, Likar B, et al. (2012) A review of methods for evaluating the quantitative parameters of sagittal pelvic alignment. *Spine J* 12, 433–446
17. Vargas Romero, B. Zarate. (2013) The impact of body mass index and central obesity on the spino-pelvic parameters: a correlation study. *Europe Spine Journal*
18. Tao Wang,Hui Wang, (2016) The characteristics of spino-pelvic sagittal parameters and obesity factors for adolescents with lumbar disc herniation. *Int J Clin Exp Med*;14321-14328