

DAFTAR PUSTAKA

- Baghchehsaraee, B., (2009), *Batch and Continuous Biohydrogen Production Using Mixed Microbial Culture*, Dissertation, Engineering Science, Department of Chemical and Biochemical Engineering, The University of Western Ontario, Canada.
- Bhakov, Z. K., Korazbekova, K. U., and Lakhanova, K. M. (2014). *The kinetics of biogas production from codigestion of cattle manure*. Pak. J. Biol. Sci. 17, pp 1023–1029.
- Cappai G, Gioannis GD, Friargiu M, Massi E, Muntoni A, Poletini A, Pomi R, Spiga D. (2014). ‘An experimental study on fermentative H₂ production from food waste as affected by pH’. *Waste Manag*;34:1510–9.
- Chen, C.C., Lin, C.Y., Lin, M.C. (2002) ‘Acid-base enrichment enhances anaerobic hydrogen production process’. *Appl. Microbiol. Biotechnol.* 58, 224–228.
- Chen, W. H., Chen, S. Y., Khanal, S. K. and Sung, S., (2006) ‘Kinetic Study of Biological Hydrogen Production by Anaerobic Fermentation’, *Int. J. Hydrogen Energy*, 31 (15), 2170-8.
- Chong, M. L., Sabaratnam, V., Shirai, Y. and Hassan, M. A., (2009) ‘Biohydrogen Production from Biomass and Industrial Wastes by Dark Fermentation’, *Int. J. Hydrog. Energy*, 34, 3277-3287.
- Chu CF, Li YY, Xu XQ, Ebie Y, Inamorie Y, Kong HN. (2008) ‘A pH- and temperaturephased two-stage process for hydrogen, methane production from food waste’. *Int J Hydrog Energy*;33:4739–46.
- Collet, C., Adler, N., Schwitzgu`ebel, J. P. and P`eringer, P., (2004) ‘Hydrogen Production by *Clostridium thermolacticum* During Continuous Fermentation of Lactose’, *Int. J. Hydrogen*, 29, 1479-85.
- Das, D. and Veziroglu, T. N., (2001), ‘Hydrogen Production by Biological Processes: A Survey of Literature’, *Int. J. Hydrogen Energy*, 26, 13-28.
- Das, D. T.N. Veziroglu, (2008) ‘Advances in biological hydrogen production processes’, *Int. J. Hydrogen Energy* 33, pp 6046-6057.

- Demirbas, A., (2009), 'Biohydrogen for Future Engine Fuel Demands', *Springer Dordrecht Heidelberg, New York*.
- Demirel B, Scherer P. (2011) 'Trace element requirements of agricultural biogas digesters during biological conversion of renewable biomass to methane'. *Biomass Bioenergy*;35:992–8.
- Dennis, A. and Burke, P. E., (2001) 'Dairy Waste Anaerobic Digestion Handbook', *Environmental Energy Company*, Olympia, WA
- Dinesh,G. Kumaravel, Chauhan. Rohit, Chakma. Sankar, (2018) 'Influence and strategies for enhanced biohydrogen production from food waste'. *Renewable and Sustainable Reviews* 92, pp 807-822.
- Dong, L., Zhenhong, Y., Yongming, S., Xiaoying, K., Yu, Z., (2009) 'Hydrogen production characteristics of the organic fraction of municipal solid wastes by anaerobic mixed culture fermentation'. *Int. J. Hydrogen Energy* 34, 812–820.
- T. Dung, B. Sen, C. Chen, G. Kumar, C. (2014). 'LinFood waste to bioenergy via anaerobic processes'. *Energy Procedia*, 61 , pp. 307-31
- Fang, H.H.P., Liu, H., (2002) 'Effect of pH on hydrogen production from glucose by a mixed culture'. *Bioresour. Technol.*82, 87–93
- FAO, (2011). Global food losses and food waste.
URL:<http://www.fao.org/docrep/014/mb060e/mb060e00.pdf>. Accessed on 5/15/2017
- Fuqing Xu, Yangyang Li, Mary Wicks, Yebo Li, and Harold Keener. (2018). 'Anaerobic Digestion of Food Waste for Bioenergy Production'. *Elsevier*, pp 530-537
- Ganesh Dattatraya Saratale¹, Rijuta Ganesh Saratale², J. Rajesh Banu³, Jo-Shu Chan. (2019). 'Biohydrogen Production from Renewable Biomass Resources'. *Biohydrogen*.pp 247-277.
- Ghimire, A., Frunzo, L., Pirozzi, F., Trably, E., Escudie, R., Lens, Piet N. L. and Esposito, G., (2015) 'A Review on Dark Fermentative Biohydrogen Production from Organic Biomass: Process Parameters and Use of By-Products', *Applied Energy*, 144, 73-95.

- Hawkes, F. R., Hussy, I., Kyazze, G., Dinsdale, R. and Hawkes, D. L., (2007) ‘Continuous Dark Fermentative Hydrogen Production by Mesophilic Microfolora: Principles and Progress’, *Int. J. Hydrogen Energy*, 32 (2), 172-84
- Heyndrickx, M., De Vos, P., Vancanneyt, M., De Ley, J., (1991) ‘The fermentation of glycerol by *Clostridium butyricum* LMG 1212 t2 and 1213tl and *C. pasteurianum* LMG 3285’. *Appl. Microbiol. Biotechnol.* 34, 637–642.
- Hwang, M. H., Jang, N. J., Hyun, S. H. and Kim, I. S., (2004) ‘Anaerobic Biohydrogen Production from Ethanol Fermentation: The Role of pH’, *J. Biotechnol.* 111 (3), 297-309.
- Kapdan, I. K. and Kargi, F., (2006) ‘Biohydrogen Production from Waste Materials’, *Enzyme Microb. Technol.*, 38 (5), 569-582
- Karadag D, Puhakka JA. (2010) ‘Enhancement of anaerobic hydrogen production by iron and nickel’. *Int J Hydrog Energy*;35:8554–60.
- M.A. Khan, H.H. Ngo, W.S. Guo, Y. Liu, L.D. Nghiem, F.I. Hai, L.J. Deng, J. Wang, Y. Wu, (2016) ‘Optimization of process parameters for production of volatile fatty acid, biohydrogen and methane from anaerobic digestion’, *Bioresour. Technol.* 219. 738-748.
- Keskin, T., H. Nalakath., Abubackar., K. Arslan. (2019). “Biohydrogen Production From Solid Waste’. *Biohydrogen* pp 312-346.
- Kharisma, A. D., (2014), ‘Pengaruh Hidrogen Peroksida Terhadap Produksi Hidrogen dari Limbah Buah Melon (*Cucumis Melo L.*) oleh Mikroba Digester Biogas’, Tesis di Jurusan Bioteknologi Sekolah Pascasarjana Universitas Gadjah Mada, Yogyakarta.
- Kim, J. K., Oh, B. R., Chun, Y. N. and Kim, S. W., (2006) ‘Effects of Temperature and Hydraulic Retention Time on Anaerobic Digestion of Food Waste’, *J. Biosci. Bioeng.*, 102, 328-32.
- Kim SH, Han SK, Shin HS. (2008) ‘Optimization of continuous hydrogen fermentation of food waste as a function of solids retention time independent of hydraulic retention time’. *Process Biochem* ;43:213–8.

- Kim DH, Kim SH, Kim HW, Kim MS, Shin HS. (2011) 'Sewage sludge addition to food waste synergistically enhances hydrogen fermentation performance'. *Bioresour Technol*;102:8501–6
- Koutrouli, E.C., Gavala, H.N., Skiadas, I.V., Lyberatos, G. (2006) 'Mesophilic biohydrogen production from olive pulp'. *Proc. Saf. Environ. Prot.* 84, 285–289.
- Lay JJ, Fan KS, Hwang JI, Chang JI, Hsu PC. (2005) 'Factors affecting hydrogen production from food wastes by clostridium-rich composts'. *J Environ Eng* ;131:595–602.
- Li, C., Fang, H.H.P., (2007) 'Fermentative hydrogen production from wastewater and solid wastes by mixed cultures'. *Crit. Rev. Environ. Sci. Technol.* 37, 1–39.
- Li, M., Zhao, Y., Guo, Q., Qian, X., Niu, D., (2008) 'Bio-hydrogen production from food waste and sewage sludge in the presence of aged refuse excavated from refuse landfill'. *Renew. Energy* 33, 2573–2579.
- Logan, B. E., (2004) 'Extracting Hydrogen and Electricity from Renewable Resources', *Environmental Sci. Tech.*, 38 (9)pp. 160-7.
- M. Rao, S.Singh, (2004), 'Bioenergy conversion studies of organic fraction of MSW : Kinetics studies and gas yield-organic loading relationships for process optimisation', *Bioresource Technology* , 92 (2) , 173-185
- Mandal T, Mandal NK. (1998) 'Biomethanation of some waste materials with pure metallic magnesium catalyst: improved biogas yields'. *Energy Convers Manag*;39:1177–9
- Mariakakis, I., Bischoff, P., Krampe, J., Meyer, C. and Steinmetz, H., (2011) 'Effect of Organic Loading Rate and Solids Retention Time on Microbial Population During Biohydrogen Production by Dark Fermentation in Large Lab-scale', *Int. J. Hydrogen Energy*, 36 (17), 10690-700
- Mu, Y., Zheng, X. J., Yu, H. Q. and Zhu, R. F., (2006) 'Biological Hydrogen Production by Anaerobic Sludge at Various Temperatures' *Int. J. Hydrogen Energy*, 31 (6), 780-5.

- Okamoto, M., Miyahara, T., Mizuno, O., Noike, T., (2000) 'Biological hydrogen potential of materials characteristic of the organic fraction of municipal solid wastes'. *Water Sci. Technol.* 41, 25–32
- Ozeler, D., Yetis, U., Demirer, G.N., (2006) 'Life cycle assesment of municipal solid waste management methods: Ankara case study'. *Environ. Int.* 32, 405–411.
- Prabhudessai, V., (2013) 'Anaerobic Digestion of Food Waste in a Horizontal Plug Flow Reactor', Thesis in Department of Biological Sciences Birla Institute of Technology and Science Pilani (Rajasthan), India.
- Posmanik, R., Labatut, R., Kim, A., Usack, J., Tester, J., Angenent, L., (2017) 'Coupling hydrothermal liquefaction and anaerobic digestion for energy valorization from model biomass feedstocks'. *Bioresour. Technol.* 233, pp 134–143.
- Ren, N. Q., Chen, Z. B., Wang, X. J., Hu, D. X. and Wang, A. J., (2005) 'Optimized Operational Parameters of a Pilot Scale Membranes Bioreactor for High Strength Organic Wastewater Treatment', *Int. Biodeterior Biodegradation*, 56, 216-23.
- Rasit, N., Idris, A., Harun, R., Ghani, W. A. W. A. K. (2015) 'Effects of lipid inhibition on biogas production of anaerobic digestion from oily effluents and sludges: An overview', *Renewable and Sustainable Energy Reviews*. Elsevier, 45, pp. 351–358.
- G.D. Saratale, S.D. Kshirsagar, R.G. Saratale, S.P. Govindwar, M.K. Oh, (2015) 'Fermentative hydrogen production using sorghum husk as a biomass feedstock and process optimization, *Biotechnol*'. *Bioproc. Eng.* 20 733-743.
- Şengor SS, Barua S, Gikas P, Ginn TR, Peyton B, Sani RK, Spycher NF. (2009) 'Influence of heavy metals on microbial growth kinetics including lag time: mathematical modeling and experimental verification'. *Environ Toxicol Chem*;28:2020–9
- Sheets, J.P., Yang, L., Ge, X., Wang, Z., Li, Y., (2015) 'Beyond land application: Emerging technologies for the treatment and reuse of anaerobically digested agricultural and food waste'. *Waste Manag.* 44, pp 94–115.

- Siddiqui Z, Horan NJ, Salter M. (2011) 'Energy optimization from co-digested waste using a two-phase process to generate hydrogen and methane'. *Int J Hydrog Energy*;36:4792–9.
- Trchounian K, Poladyan A, Trchounian A. (2017) 'Enhancement of Escherichia coli bacterial biomass and hydrogen production by some heavy metal ions and their mixtures during glycerol vs glucose fermentation at a relatively wide range of pH'. *Int J Hydrog Energy*;42:6590–7.
- Tugba Keskin, Haris Nalakath Abubackar, Kubra Arslan, Nuri Azbar. (2019) 'Biohydrogen Production From Solid Wastes'. *Biohydrogen*. pp 321-346.
- Valdez-Vazquez, I., Rios-Leal, E., Esparza-Garcia, F., Cecchi, F., Poggi-Varaldo, H.M. (2005) 'Semi-continuous solid substrate anaerobic reactors for H₂ production from organic waste: Mesophilic versus thermophilic regime'. *Int. J. Hydrog. Energy* 30, pp 1383–1391.
- Van Ginkel, S. W., Sung, S. and Lay, J. J., (2001) 'Biohydrogen Production as a Function of pH and Substrate Concentration', *Environmental Science and Technology*, 35 (24), pp 4726-4730.
- Wang J, Wan W. (2008) 'Effect of Fe²⁺ concentration on fermentative hydrogen production by mixed cultures'. *Int J Hydrog Energy*;33:1215–20.
- Wang, J., Wan, W. (2009) 'Factors influencing fermentative hydrogen production: a review'. *Int. J. Hydrog. Energy* 34(2), pp 799–811.
- Zhang, Q., Hu, J. and Lee, D. J. (2016) 'Biogas from anaerobic digestion processes: Research updates', *Renewable Energy*. Elsevier Ltd, 98, pp. 108–119.
- Zhong, J. D.K. Stevens, C.L. Hansen, (2015) 'Optimization of anaerobic hydrogen and methane production from dairy processing waste using a two-stage digestion in induced bed reactors (IBR)', *Int. J. Hydrogen Energy* 40 pp 15470-15476