

## DAFTAR PUSTAKA

- Abergel R.P., Pizzurro D., Meeker C.A, Lask G, Matsuoka LY, Minor RR, Chu ML, Uitto J. 1995. Biochemical composition of the connective tissue in keloids and analysis of collagen metabolism in keloid fibroblast cultures. *J Invest Dermatol*. 84:384-390.
- Abreu JG, Ketpura NI, Reversade N, De Robertis EM. 2002. Connective tissue growth factor (CTGF) modulates cell signaling by BMP and TGF-beta. *Nat Cell Biol*; 4 : 599-604.
- Adams BB., Gloster HM. Surgical pearl : excision with suprakeloidal flap and radiation therapy for keloid. *J Am Acad Dermatol* 2002; 47:307-9.
- Akaishi S, gawa R, Hyakusoku H. 2008. Kleoid and hypertropic scar : neurogenic inflammation hypotheses. *Med Hypotheses*;71 : 32-38.
- Akasaka Y, Fujita K, Ishikawa Y, Asuwa N, Inuzuka K, Ishihara M, Ito M, Masuda T, Akishima Y, Zhang L, Ito K, Isii T 2001. Detection of apoptosis in keloids and a comparative study on apoptosis between keloids, hypertropic scars, normalhealed flat scars, and dermatofibroma. *Wound Reair Regen*; 9 : 501-506.
- Ali S., Saeed S. 2011. Comparison of The Efficacy of Intralesional Triamcinolone Acetonide and 5-Fluorouracil Tatooning for The Treatment of Keloids. *Dermatol Surg*. 38 : 104-109.
- Al-Attar A., Mess S., Thomassesn J.M., Kauffman C.L., Davison S.P. 2006. Keloid pathogenesis and treatment. *Plast Reconstr Surg*. 117:286-300.
- Babin RW., Ceilly RI. Combined modalities in the management of hypertropics scars and keloids. *J Otolaryngol* 1979; 8 : 457-60.
- Bayat A., McGrouther D.A., Ferguson M.W.J. 2003 Skin scarring. *Br Med J*. 326:88-92
- Berman B., Villa A.M., Ramirez C.C. 2005. 1997. Novel opportunities in the treatment and prevention of scarring. *J Cutan Med Surg*. 32-36.

- Bettinger D. A., Yager D. R., Dielgelmann R.F., Cohen I.K., 1996. The effect of TGF-beta on keloid fibroblast proliferation and collagen synthesis. *Plast Reconstr Surg.* 98 : 827-833.
- Boyce DE, Ciampolini J, Ruge F, Murison MS, Harding KG. 2001. Inflammatory-cell subpopulations in keloid scars. *Br J Plast Surg*; 54 : 511-566.
- Butler P.D., Longkaker M.T., Yang G.P. 2008. Current Progress in Keloid Research and Treatment. *J Am Coll Surg.* 206:731-741.
- Calderon M., Lawrence W.T., Banes A. 1996. Increased Proliferation in Keloid Fibroblast Wounded *in vitro*. *J Surg Res.* 61:343-347.
- Carrol LA., Hanasono MM., Mikulec AA., Kita M., Koch RJ., Triamcinolon stimulates bTGF production and inhibits TGF- $\beta$  production by human dermal fibroblast dermal surgery 2002; 28:704-9.
- Chipev CC, Simman R, Hatch G, Katz AE, Siegel DM, Simon M. 2000. Myofibroblast phenotype and apoptosis in keloid and palmar fibroblast *in vitro*. *Cell Death Differ*; 7 : 166-176.
- Clark RAF. Cutaneous tissue repair : Basic biologic consideration 1. *J Am.Acad.Dermatol.*1985; 13 : 701-25
- Cohen JK., Diegelmann RF. 1977. The biology of keloid and hypertrophic scar and the influence of corticosteroid. *Clinical Plastic Surgery*; : 297-310.
- Dunkin CS, Pleat JM, Gillespie PH, Tyler MP, Roberts AH, MCGrouther DA. 2007. Scarring occurs at a critical depth of skin injury : precisemeasurement in a graduated dermal scratch in human volunteers. *Plast Reconstr Surg.* 119 : 1722-1732.
- English RS, Shenefelt PD. 1999. Keloids and Hipertropic scars. *Dermatol Surg.* 25 ; 631-638.
- Fitzpatrick TB, Freedberg IM. Eisen AZ., Wolff K., Austen KF., Goldsmith LA., Katz SI., eds. *Dermatology in general medicine.*, edisi ke 6 New york : Mc Graw Hill; 2003:236-45
- Fluorouracil. December 2003. Didapat dari : *file:///A:/fluorouracil(51—21-8)index from Fluoride Action Network Pesticide Project.htm.*

- Freshney RI. Primary culture. Dalam : Culture of animal cells, A manual of basic technique, edisi ke-4. United States : Wiley-Liss Inc;2000.
- Friedman D.W., Boyd C.D., Mackenzie J.W., Norton P., Olson R.M., Deak S.B. 1993. Regulation of collagen gene expression in keloids and hypertrophic scars. *J Surg Res.* 55:214-222.
- From L, Assaad D, Neoplasms, pseudoneoplasms and hiperplasias of the dermis, Dalam : Fitzpatrick TB, Freedberg IM, Eisen AZ, Wolff K., Austen KF, Goledsmith LA, Katz SI, eds. *Dermatology in general medicine*, edisi ke -6 New York : Mc Graw Hill; 2003 : 1163-5.
- Funayama E, Chodon T, Oyama A, Sugihara T. 2003. Keratinocytes promote proloferation and inhibit apoptosis of the underlying fibroblast : an important role in the pathogenesis of keloid. *J Invest Dermatol.* 121 : 1326-1331.
- Gilchrest BA Relationship between actinic damage and cronologic aging in keratinocyte cultures of human skin. *J Invest Dermatol* 1979;72(5):219-23.
- Gurtner GC, Werner S, Barrandon Y, Longaker MT. 2008. Wound repair and regeneration. *Nature*; 453 ; 314-321.
- Harting M., Hicks J.M., Levy M. 2008. Dermal Hypertrophies. In : Wolffk, Goldsmith L.A., Katz S.I., Gilchrest B.A., Paller A.S., Leffel D.J., editor. *Fitzpatrick's Dermatology in General Medicine*, 7th ed. Mc Graw Hill Companies, Inc. 556-558.
- Ikawa Y., Ng P.S., Endo K., Kondo M., Chujo S.,Ishida W.,Shirasaki F., Fujimoyo M., Takehara K. 2008. Neutralizing monoclonal antibody to human connective tissue growth factor ameliorates transforming growth factor beta induced mouse fibrosis. *J Cell Physiol.* 216(3) : 680-687.
- Junqueira LC, Carneiro J. *Basic histology*, edisi ke-3. Calofornia: Lanngge Medicala Publication; 1982:89-94.
- Kelly A.P. 1991. *Keloids and Hipertrophic scars. 1st ed.*. New York; McGraaw-Hill.Inc.
- Ketchum LD., Cohen IK., Master FW. Hypertrophic scars and keloids : a collective review *Paltic Reconstruction Surgery.* 1974; 53: 140-54

- Khalifa E.S., Adil A.N., Mahmood R.A. 2014. Debulking of Keloid Combined with Intralesional Injection of 5-Fluorouracil and Triamcinolone Versus Intralesional Injection of 5-Fluorouracil and Triamcinolone. *Journal of Cosmetics, Dermatological Sciences and Applications*. 4:85-91.
- Koh HK., Bhawan L., Peled Z., 5-Fluorouracil inhibits keloid fibroblast proliferasi and keloid fibroblast populated collagen lattice contraction. Dipublikasikan 15 januari 2002. Didapat dari : <http://www.journalofburns.com>.
- Luo S., Benathan M., Raffoul W., Panizzon R. G., and EgloffD. V. 2001. Abnormal balance between proliferation and apoptotic cell death in fibroblast derived from keloid lesions. *Plast Reconstr Surg*. 107 : 87-96.
- Levinson H, Liu W, Peled Z., 5-Fluorouracil inhibits keloid fibroblast proliferation and keloid fibroblast populated collagen lattice contraction. Dipublikasikan 15 januari 2002. Didapat dari : <http://www.journalofburns.com>.
- Machie RM., 1992. Soft Tissue Tumor. Dalam : Champion R.H., Burthorn J.L., Ebling F.J.G. eds. Textbook of dermatology, edisi ke-5. London : Blackwell Scientific Publ; : 2073-100.
- Marneros AG, Norris JE, Olsen BR, Reichenberger E. 2001. Clinical genetics of familial keloids. *Arch Dermatol*; 137 : 1429-1434.
- Maustafa MF, Abdel-Fattah MA, Abdel-Fattah DC. 1975. Presumptive evidence of the effect of pregnancy estrogens on keloid growth. Case report. *Plast Reconstr Surg*; 56 : 450 – 453.
- McCoy B.J., Cohen I.K. 1982. Collagenase in keloid biopsies and fibroblasts. *Connect Tissue Res*. 9:181-5.
- O’Kane S. 2002. Wound remodelling and scarring. *J Wound Care*; 11 : 296-299.
- Pierce G. F., Tarpley J. E., Yanagihara D., Mustoe T. A., Fox G. M., Thomason, A. 1992. Platelet-derived growth factor (BB homodimer), transforming growth factor-beta 1, and basic fibroblast growth factor in dermal wound healing; Neovessel and matrix formation and cessation of repair. *Am J Pathol*. 140: 1375-1388.

- Peltonen J., Hsiao L.L, Jaakkola S., Sollberg S., Aumailley M., Timpi R. 1991. Activation of Collagen Gene Expression in Keloid : Co-Localization of Type I and VI Collagen and Transforming Growth Factor – 1 mRNA. *J Invest Dermatol.* 97: 240-248.
- Perdanakusuma D.S. dan Noer M.S. 2006. Penanganan parut hipertrofik dan keloid. *Airlangga University Press.*
- Pollack SU. 2004. Management of keloids. Dalam : Wheeland RG. Cutaneous surgery. Philadelphia : W.B Saunders Co ; : 688-698.
- Raynolds JEF. Corticosteroids. Dalam : Martindale the extra pharmacopoeia, edisi ke-29. London : The Pharmaceutical Press; 1989:901-2
- Robles D.T., Berg D. 2007. Abnormal wound healing : keloids. *Clin Dermatol.* 25: 26-32.
- Russell S. B., Tupin K. M., Rodriguez-Eaton S., Russell J. D., Trupin J. S. 1998. Reduced Growth-factor requirement of keloid-derived fibroblast may account for growth. *Proc Natl Acad Sci U. S. A.* 85:587-591.
- Sandulache VC, Parekh A, Li-Korotky H, Dohar JE, Hebda PA. 2007. Prostaglandin E2 inhibition of keloid fibroblast migration, contraction, and transforming growth factor (TGF) – beta1-induced collagen synthesis. *Wound Repair Regen.* 15 : 122-133
- Subramanian R, White CJ, Sternbergh WC 3<sup>rd</sup>, Ferguson DL, Gilchrist IC. 2003. Nonhealing wound resuting from a foreign-body reaction to a radial arterial sheath. *Catheter Cardiovasc Interv;* 59 : 205-206.
- Shaffer JJ, Taylor SC., Boledn FC., 2002 Keloid scar : A review with a critical look at therapeutic option. *J Am Acad Dermatol.*; 46:S63-97).
- Shah M, Foreman DM, ferguson MW. 1995. Neutralisation of TGF-beta 1 and TGF-beta 2 or exogenous addition of TGF-beta 3 to cutaneous rat wound reduces scarring. *J Cell Sci;* 108 (Part 3) : 985-1002.
- Shejbal D., Bedekovic V., Lvkic., Kalogjera L., Aleric., Drvis P. 2004. Strategies in the treatment of keloid and hypertrophic scarr. *Acta Clinica Croatia.* 43: 417-422.

- Shi C.M, Cheng T.M., Su Y.P. 2004. Effect of dermal multipotent cell transplantation on skin wound healing. *J Surg Res.* 121(1):13-19.
- Smith JW, Bellinger CG. Keloid and hypertrophic scar. Dalam : Grabb WC, Smith JW eds. *Pastic surgery a concise guide to clinical practise*, edisi ke-2. Bston : Little Brown & Co ; 200 : 740-50
- Uitto J, Eisen AZ. Collagen. Dalam : Fitzpatrick TB., Freedberg IM., Eisen AZ., Wolff K., Austen KF., Goledsmith LA., Katz SI. Eds. *Dermatology in general medicine*, edisi ke 6 New york : Mc Graw Hill; 2003 : 165-78)
- Ulrich D., Ulrich F., Unglaub F., Platkowski A., Palhua N. 2010. Matrix metalloproteinases and tissue inhibitors of metalloproteinases in patient with different types of scars and keloids. *J Plast Reconst Aesthet Surg.* 63:1015-1021.
- Xia W, Phan TT, Lim IJ, Longaker MT, Yang GP. 2004. Complex epithelial-mesenchymal interaction modulate transforming growth factor-beta expression in keloid-derived cells. *Wound Repair Regen*; 12 : 546-556.
- Yang GP., Lim IJ., Phan TT., Lorenz HP., Longaker MT. 2003. From scarless fetal wounds to keloid : molecular studies in wound healing. *Wound repair Regen.* 11 : 411-418.
- Younai S., Nichter L. S., Wellisz T., Reinisch J., Nimni M. E., Tuan T. L. 1994. Modulation of collagen synthesus by transforming growth factor-beta in keloid and hypertrophic scar fibroblast. *Ann Plast Surg.* 33; 148-151
- Zachariae H. 1998. Delayed wound jealing and keloid formation following argon laser treatment or dermabrasion during isotretinoin treatment. *Br J Dermatol*; 118 : 703-706.