

## DAFTAR PUSTAKA

- Altar, C. a *et al.* (2015) 'Clinical validity: Combinatorial pharmacogenomics predicts antidepressant responses and healthcare utilizations better than single gene phenotypes.', *The pharmacogenomics journal*. Nature Publishing Group, 15(August 2014), pp. 1–9. doi: 10.1038/tpj.2014.85.
- American Diabetes Association (2014) 'Diagnosis and Classification of Diabetes Mellitus', *Diabetes Care*, 37(Suplemen 1), pp. 581–590. doi: 10.2337/dc14-S081.
- Baequny, A., Hartono, M. and Harnany, A. S. (2015) 'Efek Pemberian Susu Kedelai Terhadap Kadar Gula Darah Penderita Diabetes Mellitus Tipe 2', *Jurnal Informasi Kesehatan Indonesia (Jiki)*, 1, pp. 89–96.
- Beinborn, M. *et al.* (2005) 'A human glucagon-like peptide-1 receptor polymorphism results in reduced agonist responsiveness', *Regulatory Peptides*, 130(1-2), pp. 1–6. doi: 10.1016/j.regpep.2005.05.001.
- Bowen, J., Noakes, M. and Clifton, P. M. (2006) 'Appetite regulatory hormone responses to various dietary proteins differ by body mass index status despite similar reductions in ad Libitum energy intake', *J Clin Endocrinol Metab*, 91(8), pp. 2913–2919. doi: 10.1210/jc.2006-0609.
- Cahyono, A. D. (2011) 'Manfaat susu kedelai sebagai terapi penurunan kadar glukosa darah pada klien diabetes mellitus', *jurnal AKP*, 2(4), pp. 28–37.
- Chang, C. *et al.* (2013) 'Herbal Therapies for Type 2 Diabetes Mellitus: Chemistry, Biology, and Pote...: EBSCOhost', *Evidence-Based Complementary and Alternative Medicine*, 2013, pp. 1–33. doi: <http://dx.doi.org/10.1155/2013/378657>.
- Chang, J. H. *et al.* (2008) 'Effects of soybean supplementation on blood glucose, plasma lipid levels, and erythrocyte antioxidant enzyme activity in type 2 diabetes mellitus patients.', *Nutrition research and practice*, 2(3), pp. 152–7. doi: 10.4162/nrp.2008.2.3.152.
- Cheng, A. Y. Y. and Fantus, I. G. (2005) 'Oral antihyperglycemic therapy for type 2 diabetes mellitus.', *CMAJ*, 172(2), pp. 213–226. doi: 10.1503/cmaj.1031414.
- Debbiea, S. *et al.* (2014) 'Pharmacovigilance of herbal medicines', *Journal of Ethnopharmacology*, 46(10), pp. 19–22. doi: 10.3233/JRS-150643.
- Delima *et al.* (2012) 'Gambaran Praktik Penggunaan Jamu Oleh Dokter di Enam Provinsi Di Indonesia', *Bul. Penelit. Kesehat*, 40(3), pp. 109 – 122. Available at: <https://www.mysciencework.com/publication/read/1835049/gambaran-praktik-penggunaan-jamu-oleh-dokter-di-enam-provinsi-di-indonesia#page-null>.
- Ding, M. *et al.* (2016) 'Consumption of soy foods and isoflavones and risk of type 2 diabetes: A pooled analysis of three US cohorts', *EJCN*, 70(12), pp. 1381–1387. doi: 10.1038/ejcn.2016.117.
- Douglas, S. M., Lasley, T. R. and Leidy, H. J. (2015) 'Consuming Beef vs. Soy Protein Has Little Effect on Appetite, Satiety, and Food Intake in Healthy Adults', *Journal of Nutrition*, 145(5), pp. 1010–1016. doi: 10.3945/jn.114.206987.
- Herman, M. J., Supardi, S. and Handayani, R. S. (2013) 'Policy on Herbal Traditional Medicines Therapy in Three Provinces in Indonesia', *Bul. Penelit. Kesehat*, 41(2), pp. 111–119.
- International Diabetes Federation (IDF) (2015) *IDF Diabetes Atlas*. Seventh Ed, International Diabetes Federation. Seventh Ed. doi: 10.1289/image.ehp.v119.i03.

- Jakubowicz, D. and Froy, O. (2013) 'Biochemical and metabolic mechanisms by which dietary whey protein may combat obesity and Type 2 diabetes', *J Nutr Biochem*. Elsevier Inc., 24(1), pp. 1–5. doi: 10.1016/j.jnutbio.2012.07.008.
- Kaput, J. and Dawson, K. (2007) 'Complexity of type 2 diabetes mellitus data sets emerging from nutrigenomic research: A case for dimensionality reduction?', *Mutation Research*, 622(1-2), pp. 19–32. doi: 10.1016/j.mrfmmm.2007.02.033.
- Khardori, R. (2016) 'Type 2 Diabetes Mellitus', *Comparative and General Pharmacology*, pp. 1–21. Available at: <http://emedicine.medscape.com/article/117853-overview#a5>.
- Kwak, J. H. *et al.* (2010) 'Black soy peptide supplementation improves glucose control in subjects with prediabetes and newly diagnosed type 2 diabetes mellitus.', *J Med Food*, 13(6), pp. 1307–1312. doi: 10.1089/jmf.2010.1075.
- Laguer, A. and Paulsen, S. (2003) 'Name of Herb/Supplement Soy', pp. 4–7. Available at: [www.ucdenver.edu/academics/colleges/pharmacy/currentstudents/OnCampusPharmDStudents/ExperientialProgram/Documents/nutr\\_monographs/Monograph-soy.pdf](http://www.ucdenver.edu/academics/colleges/pharmacy/currentstudents/OnCampusPharmDStudents/ExperientialProgram/Documents/nutr_monographs/Monograph-soy.pdf).
- Li, W. *et al.* (2018) 'Soy and the risk of type 2 diabetes mellitus: A systematic review and meta-analysis of observational studies', *Diabetes Res. Clin. Pract.* Elsevier B.V., 137, pp. 190–199. doi: 10.1016/j.diabres.2018.01.010.
- Liu, D. *et al.* (2006) 'Genistein acutely stimulates insulin secretion in pancreatic ??-cells through a cAMP-dependent protein kinase pathway', *Diabetes*, 55(4), pp. 1043–1050. doi: 10.2337/diabetes.55.04.06.db05-1089.
- Man, S. C. *et al.* (2012) 'Systematic review on the efficacy and safety of herbal medicines for vascular dementia.', *Evidence-based complementary and alternative medicine : eCAM*, 2012(Article ID 426215), pp. 1–22. doi: 10.1155/2012/426215.
- Mansour, A. *et al.* (2013) 'Nutrients related to GLP1 secretory responses', *Nutrition*. Elsevier Inc., 29(6), pp. 813–820. doi: 10.1016/j.nut.2012.11.015.
- Moghaddam, A. S. *et al.* (2014) 'The Effects of Soy Bean Flour Enriched Bread Intake on Anthropometric Indices and Blood Pressure in Type 2 Diabetic Women: A Crossover Randomized Controlled Clinical Trial', *Int J Endocrinol*, 5(12), pp. 1529–1536. doi: 10.1155/2014/240760.
- Nguyen, C. T. *et al.* (2017) 'Soyfood and isoflavone intake and risk of type 2 diabetes in Vietnamese adults', *EJCN*, 71(10), pp. 1186–1192. doi: 10.1038/ejcn.2017.76.
- Niggemann, B. and Grüber, C. (2003) 'Side-effects of complementary and alternative medicine.', *Allergy*, 58(8), pp. 707–716. doi: 10.1034/j.1398-9995.2003.00219.x.
- Olokoba, A.B., Obatero, O.A., Olokoba, L. B. (2012) 'Type 2 diabetes mellitus: A review of current trends', *Oman Medical Journal*, 27(4), pp. 269–273. doi: 10.5001/omj.2012.68.
- PERKENI (2015) *Konsensus Pengelolaan dan Pencegahan Diabetes Melitus Tipe 2 di Indonesia 2015*. Juli 2015. PB. PERKENI. Available at: <https://pbperkeni.or.id/wp-content/uploads/2019/01/4.-Konsensus-Pengelolaan-dan-Pencegahan-Diabetes-melitus-tipe-2-di-Indonesia-PERKENI-2015.pdf> (Accessed: 18 March 2019).

- Petznick, A. (2011) 'Insulin management of type 2 diabetes mellitus', *American Family Physician*, 84(2), pp. 183–190. doi: 10.1016/j.mcna.2004.04.002.
- Punthakee, Z., Goldenberg, R. and Katz, P. (2018) 'Definition, Classification and Diagnosis of Diabetes, Prediabetes and Metabolic Syndrome', *Can J Diabetes*, 42, pp. 10–15. doi: 10.1016/j.jcjd.2017.10.003.
- RISKESDAS (2013) *Riset Kesehatan Dasar*.
- Sathananthan, A. *et al.* (2010) 'Common Genetic Variation in GLP1R and Insulin Secretion in Response to Exogenous GLP-1 in Nondiabetic Subjects', *Diabetes Care*, 33(9), pp. 2074–2076. doi: 10.2337/dc10-0200.
- Sathyapalan, T. *et al.* (2017) 'Effect of soy in men with type 2 diabetes mellitus and subclinical hypogonadism: A randomized controlled study', *J Clin Endocrinol Metab*, 102(2), pp. 425–433. doi: 10.1210/jc.2016-2875.
- Savelyeva, A. V. *et al.* (2017) 'Variety of RNAs in Peripheral Blood Cells, Plasma, and Plasma Fractions', *Biomed Res. Int.*, 2017. doi: 10.1155/2017/7404912.
- Sheikh, H. I. *et al.* (2010) 'Glucagon-like peptide-1 receptor gene polymorphism (Leu260Phe) is associated with morning cortisol in preschoolers', *Prog Neuropsychopharmacol Biol Psychiatry*. Elsevier Inc., 34(6), pp. 410–412. doi: 10.1016/j.pnpbp.2010.05.007.
- Sinaga, E. and Wirawanni, Y. (2013) 'Pengaruh pemberian susu kedelai terhadap kadar glukosa darah puasa pada wanita prediabet', *Journal of Nutrition College*, 2(1), pp. 312–321. Available at: <http://ejournal-s1.undip.ac.id/index.php/jnc>.
- Sun, L. *et al.* (2016) 'Soya milk exerts different effects on plasma amino acid responses and incretin hormone secretion compared with cows' milk in healthy, young men', *British Journal of Nutrition*, 116(07), pp. 1216–1221. doi: 10.1017/s0007114516003214.
- Talaei, M. (2015) 'Role of phytoestrogens in prevention and management of type 2 diabetes', *World J Diabetes*, 6(2), pp. 271–283. doi: 10.4239/wjd.v6.i2.271.
- Tokuyama, Y. *et al.* (2004) 'Five missense mutations in glucagon-like peptide 1 receptor gene in Japanese population', *Diabetes Research and Clinical Practice*, 66(1), pp. 63–69. doi: 10.1016/j.diabres.2004.02.004.
- Tulk, B. *et al.* (2014) 'Identification of a bioactive peptide from Glycine max that stimulates the release of cholecystokinin and glucagon-like peptide-1 from enteroendocrine cells (372.1)', *Faseb J*, 28(1\_Supplement), p. 372.1–. Available at: [http://www.fasebj.org/cgi/content/long/28/1\\_Supplement/372.1](http://www.fasebj.org/cgi/content/long/28/1_Supplement/372.1).
- Wang, H. P. *et al.* (2007) 'From pharmacovigilance to pharmacovigilance planning - The system building for safe medication', *J Food Drug Anal*, 15(4), pp. 377–386.
- Williams, J. a (2014) 'Glp-1', *The pancreapedia*, 1(67), pp. 1–14. doi: 10.3998/panc.2014.7.
- Yang, B. *et al.* (2011) 'Systematic review and meta-analysis of soy products consumption in patients with type 2 diabetes mellitus', *Asia Pacific Journal of Clinical Nutrition*, 20(4), pp. 593–602.
- Zhang, X. M., Zhang, Y. B. and Chi, M. H. (2016) 'Soy protein supplementation reduces clinical indices in type 2 diabetes and metabolic syndrome', *Yonsei Medical J*, 57(3), pp. 681–689. doi: 10.3349/ymj.2016.57.3.681.