

DAFTAR PUSTAKA

- AAP COMMITTEE ON FETUS AND NEWBORN and SECTION ON ANESTHESIOLOGY AND PAIN MEDICINE (2016) 'Prevention and Management of Procedural Pain in the Neonate : An Update', *Pediatrics*, 137(2), p. e20154271. doi: 10.1542/peds.2015-4271.
- Akcan, E. and Polat, S. (2016) 'Comparative Effect of the smells of Amniotic Fluid, Breast milk, and Lavender on Newborns ' Pain During Heel Lance', *Breastfeeding Medicine*, 11(6). doi: 10.1089/bfm.2015.0174.
- Almutary, H., Bonner, A. and Douglas, C. (2015) 'Arabic translation , adaptation and modification of the Dialysis Symptom Index for chronic kidney disease stages ... Arabic translation , adaptation and modification of the dialysis symptom index for chronic kidney disease stages four and five', *BMC Nephrology*, 16(36). doi: 10.1186/s12882-015-0036-2.
- Anand, K. J. S. and Hickey, P. . (1987) 'Pain and its effects in the human neonate and fetus', *The New England Journal of Medicine*, 317(21), pp. 1321–1329.
- Aoyama, S. *et al.* (2010) 'Maternal breast milk odour induces frontal lobe activation in neonates : A NIRS study', *Early Human Development*. Elsevier Ireland Ltd, 86, pp. 541–545. doi: 10.1016/j.earlhumdev.2010.07.003.
- Badan kependudukan dan Keluarga Berencana Nasional, B. *et al.* (2013) 'Survei Demografi dan Kesehatan Indonesia 2012'.
- Badiee, Z., Asghari, M. and Mohammadzadeh, M. (2013) 'The Calming Effect of Maternal Breast Milk Odor on Premature Infants', *Pediatrics and Neonatology*. Elsevier Taiwan LLC, pp. 322–325. doi: 10.1016/j.pedneo.2013.04.004.
- Badr, L. K. *et al.* (2010) 'Determinants of Premature Infant Pain', *Pediatric nursing*, 36(3), pp. 129–136.
- Badr, L. K. (no date) 'Pain Interventions in Premature Infants : What Is Conclusive Evidence and What Is Not', *Newborn and Infant Nursing Reviews*. Elsevier Inc., 12(3), pp. 141–153. doi: 10.1053/j.nainr.2012.06.007.
- Barett, T., Kent, S. and Voudouris, N. (1999) 'Does melatonin modulate beta-endorphine, corticosterone, and pain threshold', *Life Science*, 66(6), pp. 467–476.
- Belliemi, C. V. *et al.* (2002) 'Effect of Multisensory Stimulation on Analgesia in Term Neonates : A Randomized Controlled Trial', *Pediatric research*, 51(4), pp. 460–463.

- Bellieni, C. V. *et al.* (2012) 'Sensorial saturation for infants ' pain', *Journal of maternal-Fetal and Neonatal Medicine*, pp. 1–3. doi: 10.3109/14767058.2012.663548.
- Bellieni, C. V., Tei, M. and Buonocore, G. (2016) 'Contextualized pain management in newborns', *Journal of pediatric and neonatal individualized medicine*, 5(2), pp. 1–7. doi: 10.7363/050209.
- Bellieni, C. V *et al.* (2007) 'Sensorial Saturation for Neonatal Analgesia', *Clinical Journal of Pain*, 23(3), pp. 219–221.
- Bembich, S. *et al.* (2013) 'Functional Neuroimaging of Breastfeeding Analgesia by Multichannel Near-Infrared Spectroscopy', *Neonatology*, 104(4), pp. 255–9. doi: 10.1159/000353386.
- Blass, E. M., Hoffmeyer, L. B. and Uni-, C. (1991) 'Sucrose as an Analgesic for Newborn Infants', *Pediatric research*, 87(2).
- Bouza, H. (2009) 'The impact of pain in the immature brain.', *The journal of maternal-fetal & neonatal medicine : the official journal of the European Association of Perinatal Medicine, the Federation of Asia and Oceania Perinatal Societies, the International Society of Perinatal Obstetricians*, 22(9), pp. 722–732. doi: 10.3109/14767050902926962.
- Breivik, H. *et al.* (2008) 'Assessment of pain', *British Journal of Anaesthesia*, 101(1), pp. 17–24. doi: 10.1093/bja/aen103.
- Brislin, R. W. (1986) 'The wording and translation of research instruments', in *Field methods in cross-cultural research*. Newbury Park, pp. 137–164.
- Brummelte, S. *et al.* (2012) 'Procedural pain and brain development in premature newborns', *Annals of Neurology*, 71(3), pp. 385–396. doi: 10.1002/ana.22267.Procedural.
- Bueno, M. *et al.* (2012) 'Breast Milk and Glucose for Pain Relief in Preterm Infants: A Noninferiority Randomized Controlled Trial', *Pediatrics*, 129(4). doi: 10.1542/peds.2011-2024.
- Carbajal, R. *et al.* (1999) 'Randomised trial of analgesic effects of sucrose, glucose, and pacifiers in term neonates', *British Medical Journal*, 319(7222), pp. 1393–1397.
- Carbajal, R. *et al.* (2003) 'Analgesic effect of breast feeding in term neonates: Randomised Controlled Trial', *British Medical Journal*, 326(7379), pp. 13–15.
- Carbajal, R. *et al.* (2008) 'Epidemiology and Treatment of Painful Procedures in Neonates in Intensive Care Units', *JAMA: The Journal of the American Medical Association*, 300(1).
- Carbajal, R. (2008) 'Nonpharmacological treatment of neonatal pain', in *Neonatal Pain: Suffering, Pain and Risk of Brain Damage in the Fetus and Newborn*, pp. 83–97. doi: 10.1007/978-88-470-0732-1_11.

- Catelin, C. *et al.* (2005) 'Clinical, Physiologic, and Biologic Impact of Environmental and Behavioral Interventions in Neonates During a Routine Nursing Procedure', *The Journal of Pain*, 6(12), pp. 791–797. doi: 10.1016/j.jpain.2005.07.010.
- Chang, Y. *et al.* (2014) 'Sample Size Determination for a Three-Arm Equivalence Trial of Normally Sample Size Determination for a Three-Arm Equivalence Trial of Normally Distributed Responses', *Journal of Biopharmaceutical Statistics*, 24(6), pp. 1190–1202. doi: 10.1080/10543406.2014.948552.
- Cignacco, E. *et al.* (2008) 'Routine procedures in NICUs: Factors influencing pain assessment and ranking by pain intensity', *Swiss Medical Weekly*, 138(33–34), pp. 484–491. doi: 2008/33/smw-12147.
- Codipietro, L., Ceccarelli, M. and Ponzone, A. (2008) 'Breastfeeding or Oral Sucrose Solution in Term Neonates Receiving Heel Lance: A Randomized', *Pediatrics*, 122, pp. 716–721. doi: 10.1542/peds.2008-0221.
- Courtois, E. *et al.* (2016) 'The burden of venipuncture pain in neonatal intensive care units: EIPPAIN 2, a prospective observational study', *International Journal of Nursing Studies*, 57, pp. 48–59. doi: 10.1016/j.ijnurstu.2016.01.014.
- Fitzgerald, M. (2005) 'The development of nociceptive circuits', *Nature Review Neuroscience*, 6(7), pp. 507–520. doi: 10.1038/nrn1701.
- Fitzgerald, M. and Beggs, S. (2001) 'The Neurobiology of Pain: Developmental Aspects', *The Neuroscientist*, 7(3), pp. 246–257.
- Gibbins, S. M. D. (2001) *Efficacy and safety of sucrose for procedural pain relief in preterm and term neonates*. University of Toronto.
- Gibbins, S. and Stevens, B. (2001) 'Mechanisms of sucrose and non-nutritive sucking in procedural pain management in infants', *Pediatric Pain Management*, 6(1), pp. 33–37.
- Gitto, E. *et al.* (2012) 'Stress response and procedural pain in the preterm newborn: The role of pharmacological and non-pharmacological treatments', *European Journal of Pediatrics*, 171(6), pp. 927–933. doi: 10.1007/s00431-011-1655-7.
- Golianu, B. *et al.* (2007) 'Non-Pharmacological Techniques for Pain Management in Neonates', *Seminars in Perinatology*. Elsevier Inc., 31(5), pp. 318–322. doi: 10.1053/j.semperi.2007.07.007.
- Graham, Y. P. *et al.* (1999) 'The effects of neonatal stress on brain development: Implications for psychopathology', *Development and Psychopathology*, 11, pp. 545–565.
- Grunau, R. E. *et al.* (2001) 'Demographic and therapeutic determinants of pain reactivity in Very Low Birth Weight neonates at 32 weeks'

- postconceptional age', *Pediatrics*, 107(1), pp. 105–112.
- Hall, R. W. and Anand, K. J. S. (2005) 'Physiology of Pain and Stress in the Newborn', *Neo Review*, 6(2).
- Hatfield, L. A. (2014) 'Neonatal pain: What's age got to do with it?', *Surgical Neurology International*, 5(Suppl 13), pp. S479–S489. doi: 10.4103/2152-7806.144630.
- Heine, W. (1999) 'The significance of tryptophane in infant nutrition', in Huether, G. et al. (eds) *Tryptophane, serotonin, and melatonin, basic aspects and application*. 1st edn. New York: Springer science+business media New York, p. 705. doi: 10.1007/978-1-4615-4709-9.
- Herrington, C. J. (2007) *Reducing pain of heelstick in premature infants with gentle human touch*. Wayne State University Detroit, Michigan.
- Hoffmann, T. C. et al. (2014) 'Better reporting of interventions: template for intervention description and replication (TIDieR) checklist and guide', *BMJ*, 1687(March), pp. 1–13. doi: 10.1136/bmj.g1687.
- Johnston, C. C., Fernandes, A. M. and Campbell-yeo, M. (2011) 'Pain in neonates is different', *Pain International Association for the Study of Pain*, 152(3), pp. S65–S73. doi: 10.1016/j.pain.2010.10.008.
- Jones, P. S. et al. (2001) 'An Adaptation of Brislin's Translation Model', *Nursing research*, 50(5), pp. 300–304. doi: 10.1097/00006199-200109000-00008.
- Kementrian Kesehatan RI (2015) *PROFIL KESEHATAN INDONESIA 2014*.
- Kunisch, J. A. (2014) 'Reducing stress in the neonatal intensive care unit: An occupational therapy approach to preterm infant massage.', *Reducing Stress in the Neonatal Intensive Care Unit: An Occupational Therapy Approach to Preterm Infant Massage*, pp. 137 p – 137 p 1p. Available at: <http://search.ebscohost.com/login.aspx?direct=true&db=rzh&AN=109763233&site=ehost-live>.
- Lee, J. S. et al. (2016) 'Pain Response to Procedural Pain in Premature Infants Pain Response to Procedural Pain in Premature Infants', *J Korean Acad Child Health Nurs*, 16(4), pp. 352–359. doi: 10.4094/jkachn.2010.16.4.352.
- Li, J. and Baccei, M. L. (2016) 'Neonatal Tissue Damage Promotes Spike Timing-Dependent Synaptic Long-Term Potentiation in Adult Spinal Projection Neurons', *The Journal of neuroscience*, 36(19), pp. 5405–5416. doi: 10.1523/JNEUROSCI.3547-15.2016.
- Lima, J. De, Anaesthetist, P. and Carmo, K. B. (2010) 'Best Practice & Research Clinical Anaesthesiology Practical pain management in the neonate', *Best Practice & Research Clinical Anaesthesiology*. Elsevier Ltd, 24(3), pp. 291–307. doi: 10.1016/j.bpa.2010.04.001.

- Loeser, J. D. and Ronald Melzack (1999) 'Pain: an overview', *Pain*, 353(9164), pp. 1607–1609.
- Maastrup, R. *et al.* (2012) 'Breastfeeding support in neonatal intensive care: A national survey', *Journal of Human Lactation*, 28(3), pp. 370–379. doi: 10.1177/0890334412440846.
- Meliala, L. (2004) *Terapi rasional nyeri Tinjauan khusus nyeri neuropatik*. Yogyakarta: Aditya media.
- Mooney-Leber, S. M. and Brummelte, S. (2017) 'Neonatal pain and reduced maternal care: Early-life stressors interacting to impact brain and behavioral development', *Neuroscience*. IBRO, 342, pp. 21–36. doi: 10.1016/j.neuroscience.2016.05.001.
- Mörelus, E., He, H.-G. and Shorey, S. (2016) 'Salivary Cortisol Reactivity in Preterm Infants in Neonatal Intensive Care: An Integrative Review', *International Journal of Environmental Research and Public Health*, 13(3), p. 337. doi: 10.3390/ijerph13030337.
- Do Nascimento, M. B. R. and Issler, H. (2004) 'Breastfeeding in premature infants: In-hospital clinical management', *Jornal de Pediatria*, 80(5 SUPPL. 5), pp. 163–172.
- Neshat, H., Jembreili, M. and Seyyedrasouli, A. (2016) 'Effects of Breast Milk and Vanilla Odors on Premature Neonate ' s Heart Rate and Blood Oxygen Saturation During and After Venipuncture', *Pediatrics and Neonatology*. Elsevier Taiwan LLC, 57(3), pp. 225–231. doi: 10.1016/j.pedneo.2015.09.004.
- Provenzi, L. *et al.* (2015) 'Pain-related stress during the Neonatal Intensive Care Unit stay and SLC6A4 methylation in very preterm infants.', *Frontiers in behavioral neuroscience*, 9(April), p. 99. doi: 10.3389/fnbeh.2015.00099.
- Ramenghi, L. A., Evans, D. J. and Levene, M. I. (1999) "“Sucrose analgesia”: Absorptive mechanism or taste perception?", *Archives of Disease in Childhood: Fetal and Neonatal Edition*, 80(2), pp. F146–F147.
- Riddell, P. *et al.* (2013) 'Non-pharmacological management of infant and young child procedural pain', *Cochrane Database Syst Rev*, (10), p. CD006275. doi: 10.1002/14651858.CD006275.pub2.
- Riset Kesehatan Dasar (2013) 'Penyajian Pokok-Pokok Hasil Riset Kesehatan Dasar 2013'.
- Rocha, E. M. and Prkachin, K. M. (2007) 'Temperament and pain reactivity predict health behavior seven years later', *Journal of Pediatric Psychology*, 32(4), pp. 393–399. doi: 10.1093/jpepsy/jsl036.
- Rodrigues, A. C. and Guinsburg, R. (2013) 'Pain evaluation after a non-nociceptive stimulus in preterm infants during the first 28 days of life', *Early Human Development*. Elsevier Ireland Ltd, 89, pp. 75–79. doi: 10.1016/j.earlhumdev.2012.08.002.

- Rosali, L. *et al.* (2015) 'Efficacy of Expressed Breast Milk in Reducing Pain During ROP Screening--a Randomized Controlled Trial', *Journal of Tropical Pediatrics*, 61(2), pp. 135–138. doi: 10.1093/tropej/fmu073.
- Schiller, C. (1999) *Clinical utility of two neonatal pain assessment measures*. University of Toronto.
- Schulz, K. F. *et al.* (2010) 'CONSORT 2010 Statement : updated guidelines for reporting parallel group randomised trials', *Trials*, 11(32), pp. 1–8.
- Schwaller, F. and Fitzgerald, M. (2014) 'The consequences of pain in early life: Injury-induced plasticity in developing pain pathways', *European Journal of Neuroscience*, 39(3), pp. 344–352. doi: 10.1111/ejn.12414.
- Senkowski, D., Höfle, M. and Engel, A. K. (2014) 'Crossmodal shaping of pain: A multisensory approach to nociception', *Trends in Cognitive Sciences*, 18(6), pp. 319–327. doi: 10.1016/j.tics.2014.03.005.
- Shah, P. S. *et al.* (2012) 'Breastfeeding or breast milk for procedural pain in neonates.', *Cochrane database of systematic reviews (Online)*, 3(12), p. CD004950. doi: 10.1002/14651858.CD004950.pub3.
- Simons, S. H. P. *et al.* (2003) 'Do We Still Hurt Newborn Babies ? A prospective study of procedural pain and analgesia in neonates', *Arch Pediatr Adolesc Med*, 157, pp. 5–9.
- Simonse, A. E. and Mulder, P. G. H. (2012) 'Analgesic Effect of Breast Milk Versus Sucrose for Analgesia During Heel Lance in Late Preterm Infants', *Pediatrics*, 129(4), pp. 657–663. doi: 10.1542/peds.2011-2173.
- Slater, R. *et al.* (2010) 'Oral sucrose as an analgesic drug for procedural pain in newborn infants : a randomised controlled trial', *The Lancet*. Elsevier Ltd, 376(9748), pp. 1225–1232. doi: 10.1016/S0140-6736(10)61303-7.
- Smith, E. M. W. (2011) 'Procedural pain management in neonates , infants and children', *Reviews in Pain*, 5(3), pp. 4–12.
- Stevens, B. *et al.* (2016) 'Sucrose for analgesia in newborn infants undergoing painful procedures (Review)', *Cochrane Database of Systematic Reviews*, (7). doi: 10.1002/14651858.CD001069.pub5. www.cochranelibrary.com.
- Stevens, B. J. *et al.* (2014) 'The Premature Infant Pain Profile-Revised (PIPP-R)', *Clinical Journal of Pain*, 30(3), pp. 238–243.
- Tomczak, M. and Tomczak, E. (2014) 'The need to report effect size estimates revisited. An overview of some recommended measures of effect size', *Trends in Sport Sciences*, 1(21), pp. 19–25. doi: 10.7326/0003-4819-159-3-201308060-00005.
- Treede, R. D. (2018) 'The International Association for the Study of Pain definition of pain: As valid in 2018 as in 1979, but in need of regularly updated footnotes', *Pain Reports*, 3(2), pp. 3–5. doi:

10.1097/PR9.0000000000000643.

- Trivedi, D. (2015) 'Cochrane Review Summary : Massage for promoting mental and physical health in typically developing infants under the age of six months', *Primary Health Care Research & Development*, 16, pp. 3–4. doi: 10.1017/S1463423614000462.
- Twycross, A., Drowden, S. and Stinson, J. (2014) *Managing pain*. Wiley Blackwell.
- Twycross, A., Stephanie, D. and Stinson, J. (2014) *Managing Pain in Children, A Clinical Guide for Nurses and Healthcare professionals*. 2nd edn. Wiley Blackwell.
- Valeri, B. O. *et al.* (2014) 'Pain reactivity in preterm neonates : examining the', *European Journal of Pain*, 18, pp. 1431–1439. doi: 10.1002/ejp.508.
- Verriotis, M. *et al.* (2016) 'The development of the nociceptive brain', *Neuroscience*. IBRO, 338, pp. 207–219. doi: 10.1016/j.neuroscience.2016.07.026.
- Vinall, J. and Grunau, R. E. (2014) 'Impact of repeated procedural pain-related stress in infants born very preterm.', *Pediatric research*, 75(5), pp. 584–7. doi: 10.1038/pr.2014.16.
- WHO (2018) *Process of translation and adaptation of instruments*.
- Williams, A. C. D. C. and Craig, K. D. (2016) 'Updating the definition of pain', *Pain*, 157, pp. 2420–2423. doi: 10.1097/j.pain.0000000000000613.
- Witt, N. *et al.* (2016) 'A Guide to Pain Assessment and Management in the Neonate', *Current Emergency and Hospital Medicine Reports*. Springer US, 4(1), pp. 1–10. doi: 10.1007/s40138-016-0089-y.
- Zeraati, H. *et al.* (2017) 'Effect of Multisensory Stimulation on Pain of Eye Examination in Preterm Infants', *Anesthesiology and Pain Medicine*, 7(1), pp. 1–7. doi: 10.5812/aapm.42561.
- Zhong, B. (2009) 'How to Calculate Sample Size in Randomized Controlled Trial?', *Journal of Thoracic Disease*, 1(1), pp. 51–54.
- Zwicker, J. G. *et al.* (2015) 'Score for neonatal acute physiology-II predict corticospinal tract development in premature newborns', *Pediatric neurology*, 48(2), pp. 123–129. doi: 10.1016/j.pediatrneurol.2012.10.016.SNAP-II.