

## DAFTAR PUSTAKA

- Abe M, Hamano T, Wada A. Effect of dialyzer membrane materials on survival in chronic hemodialysis patients: results from the annual survey of the Japanese Nationwide Dialysis Registry. *Plos One* 2017; 12: 018442
- Abravkar PN, Avikar JS, Shilpa B. Study of serum uric acid and C- reactive protein levels in patients with renal disease., *International Journal Biology Medical Research*. 2013;4 (1): 2758-61.
- Black, J.M, Hawks, J.H. 2005. *Medical Surgical Nursing Clinical Management for Possitive Outcome 7th edition*. Philadelphia : W.B Saunders Company.
- Borazan A, Hasan Ustun, Yucel Ustundag, Selim Aydemir, Taner Bayraktaroglu, Mehmet Sert, Ahmet Yilmaz, 2004. The effects of Peritoneal Dialysis and Hemodialysis on serum tumor necrosis factor Alpha, Interleukin -6, Interleukin -10 and C-reactive protein Levels. *Mediator of Inflammation*, 13 (3) : 201-204.
- Borazan A, Selim Aydemir, Mehmet Sert, Ahmet Yilmaz, 2004. The effects of Hemodialysis and Peritoneal Dialysis on serum homocysreine and Creactive protein Levels. *Mediator of Inflammation*, 13 (5/6) : 361-364.
- Cheung AK, Nathan WL, Greene T, Agodoa L, Bailey J, Beck G.2003. Effects of high flux hemodialysis on clinical outcomes: Results of the HEMO study. *J Am Soc Nephrol*;14(12):3251-63.
- Chu PL, Chiu YL, Lin JW, Chen SI, Wu KD, 2008: Effects of low- and high-flux dialyzers on oxidative stress and insulin resistance. *Blood Purif*, 26: 213–220.
- C.K. Wong, C.C. Szeto, M.H.M. Chan, C.B. Leung, P.K.T. Li, C.W.K. Lam, 2007. Elevation of Protein Inflammatory Cytokines, C-Reactive Protein and Cardiac Troponin T in Chronic Renal Failure Patients on Dialysis. *Immunological Investigations*, 36 : 47-57.
- Coresh J, Byrd-Holt D, Astor BC, 2005. Chronic kidney disease awareness, prevalence, and trends among U.S. adults, 1999 to 2000. *J AmSoc Nephrol*;16:180-8
- Cronin, R.E., Henrich, W.L. 2010. Kt/V and The Adequacy of Hemodialysis. <http://www.ncbi.nlm.nih.gov/pubmed/9807323>
- Daugirdas, J.T., Blake, P.G., Ing, T.S. 2007. Handbook of Dialysys 4th edition. Philadelphia: Lippincott.
- Dati F, Myron Johnson A, Whicher JT 2001. The Existing Interim Consensus Reference Ranges and the Future Approach. *Clin Chem Lab Med*; 39(11):1134-1136.
- Descamps, Latscha B., Goldford, B., Nguyen, A. T., 1991. Esthablishing the relationship between complement activation and stimulation of fagocyte oxitative metabolism in hemodialysed patients: a randomized prospectivestudy. *Nephron*, 59:279-285.
- Dewi, I.G.A.P.A. 2010. Hubungan antara Quick of Blood/Qb dengan Adekuasi Hemodialisis di Ruang HD BRSU Daerah Tabanan Bali. Tesis. Tidak dipublikasikan.

- DU-Clos, T.W., 2000. Function of C-Reactive Protein, *Annual Medicine* 32 : 274-278.
- Eknoyan, G., Levin, N.W., Eschbach, J.W. 2000. KDOQI Clinical Practice Guidelines. <http://www.asnjournals.org>
- El-Wakil HS, Abou-Zeid AA, El-Gohary IE, Abou El-Seoud NA, Abou El-Yazeed MA, El-Leithy MA., 2011. Effect of high flux versus low flux hemodialysis on serum beta-2 microglobulin, advanced oxidation protein products and protein carbonyl. *Alexandria JMed.*;47(1):37–42.
- Emad A.,M, Magdy Elsayed Mohamed, Haytham Sabry Abdelhamid, Mohammed Al-Arabi Mohammed., 2017. Effects of High Flux versus Low Flux on Serum C-Reactive ProteinA as an Inflammatory Biomarker in Hemodialysis Patients., *The Egyptian Journal of Hospital Medicine* Vol.67 (2), Page 645-655
- Erlinda, 2014. Hubungan Kadar Asam Urat Serum Dengan Kadar High Sensitivity C-Reactive Protein (HSCRP) Pasien Hemodialisis Rumah Sakit Umum Daerah DR. ZAINOEL ABIDIN BANDA ACEH. Tesis. Tidak Dipublikasikan.
- Filiopoulos V, Hadjiyannakos D, Metaxaki P., 2008. Inflammation and oxidative stress in patients on hemodiafiltration. *American Journal of Nephrology*. 28(6):949– 957.
- Fink, J.C. 2001. Spesifik faktor on Varians in Dialysis Adequacy. *Journal of the American Society of Nephrology*, 12 : 164-169.
- Fluchart, J.C., 2003. Atherosclerosis: an Inflammatory Disease?, International Task Force for Prevention of Coronary Heart disease. Task Force Symposium.
- Gatot, D. 2003. Rasio Reduksi Ureum dalam Dialisis. <http://library.usu.ac.id/download/fk/penyakitdalam-dairot%20gatot.pdf>.
- Hadim Akoglu., Fatih Dede., Serhan Piskinpasa., Mesude Y. Falay., Ali Riza Odabas., 2013. Impact of Low- or High-Flux Haemodialysis and Online Haemodiafiltration on Inflammatory Markers and Lipid Profile in Chronic Haemodialysis Patients, Ankara Numune Research and Education Hospital, Ankara , Turkey, *Blood Purif* 2013;35:258–264.
- Hak AE., Stehouwer C, Bots Moche., Polderman K, Schalkwijk C, Westendorp I et al., 1999, Associations of C Reactive Protein With Measure of Obesity, Insulin Resistance and SubClinical Atherosclerosis in Healthy, Midlle Aged Woman, *Atherosclerosis Thrombosis Vascular Biol.*, 19: 1986-91.
- Hastono, P.S. 2007. Analisa Data Kesehatan. Fakultas Kesehatan Masyarakat Universitas Indonesia. Buku tidak dipublikasikan.
- Henrie Met al.(2008).In vitro assessment of dialysis membranes as an endotoxin transfer barrier: geometry, morphology and permeability. *Artif Organs*, 32, pp. 701–710
- Hirschfield, G.M., Pepys,M.B., 2003. C-Reactive Protein and Cardiovascular Disease: New Insight From an Old Molecule, *Quarterly Journal of Medicine*. 96:793-807.
- Imelda Fitri, Endang Susalit, M Bonar M Marbun, Cleopas Martin Rumende, 2017., Gambaran Klinis dan Kualitas Hidup Pasien Penyakit Ginjal Tahap

- Akhir yang Menjalani Hemodialisis Dua Kali Dibandingkan Tiga Kali Seminggu., *Jurnal Penyakit Dalam.*, Vol.4., no.3., September 2017.
- Indonesian renal registry. 2014. 7<sup>th</sup> Report Of Indonesian Renal Registry.
- Jindal, K., Chan, C.T. 2006. Hemodialysis adequacy in Adult. *Journal of The American Society of Nephrology*. 2006: 17:4-7.
- Johanna, C.K., Jeannette, G., Van Manem, Friedo, W.D., Dirk, R.D.W., Elisabeth, W.B., Raymond, T.K., 2004. Effect of increase in C-reactive Protein level during a Hemodialysis session on Mortality. *J Am Soc Nephrol* 15 : 2916-2922.
- Johnson RJ, Herrera-Acosta J, Schreiner GF, Rodriguez-Iturbe B. 2002. Subtle acquired renal injury as a mechanism of salt-sensitive hypertension. *N Engl J Med*. 346:913–923.
- K/DOQI. 2006. Clinical practice guidelines for hemodialysis adequacy. Maret 2, 2010. <http://www.kidney.org/Professionals/kdoqi>.
- Karkar A, Abdelrahman M, Locatelli F., 2015. A randomized trial on health-related patient satisfaction level with high-efficiency online hemodiafiltration versus high-flux dialysis. *Blood Purif*; 40: 84-91.
- Koenig W, Lowel H, Baumert J, Meisinger C, 2004. C-reactive protein modulates risk prediction based on the Framingham Score: implications for future risk assessment: results from a large cohort study in southern Germany. *Circulation, American Heart Association.*;109:1349-53.
- Li, F.K., Lai, K.N. 2000. Current Issues in the Chronic Renal failure : *Dialysis. Havas Media*.
- Marion Haubitz, Reinhard Brukhorst, Eike Wrenger, Peter Froese, Matthias Schulze, Karl-Martin Koch., 1996. Chronic Induction of C-Reactive Protein by Hemodialysis, but not by Peritoneal dialysis therapy. *Journal of The International Society for Peritoneal Dialysis*. 16 : 158-162.
- Memoli B. Postiglione L, Cianciaruso B., 2000. Role of different dialysis membran in the release of interleukin- 6 – Soluble receptor in uremic patiens. *International Society of Nephrology*. 58(1): 417-24.
- Merello Godino JI, Rentero R, Orlandini G, Marcelli D, Ronco C., 2002.:Results from EuCliD (European Clinical Dialysis Database):Impact of shifting treatment modality.*Int J Artif Organs*25: 1049–1060.
- Munshi R, Ahmad S., 2014. Comparison of urea clearance in low efficiency low flux vs. high efficiency high- flux dialyzer membrane with reduced blood and dialysate flow: An in vitro analysis. *Hemodial Int.*;18:172-4. doi: 10.1111/hdi.12054.
- National Kidney Foundation. 2001. Guidelines for Hemodialysis adequacy. <http://www.kidney.org/Professionals/kdoqi/>
- Nguyen, D. Van-Hong., and McLaughlin M,A., 2002, Coronary Artery Disease in Women: A Review of Emerging Cardiovascular Risk factor, *The Mount Sinai Journal of Medicine*, Vol 69 no 5: 338-47.
- Owen, W.F. 2000. Hemodialysis Adequacy. <http://www.oxfordjournals.org>.
- Pearson T, Blair S, Daniels S, Stephen R Daniels., Robert H Eckel., Joan M Fair., Stephen P. Portmann., et al., 2002. AHA guidelines for primary prevention

of cardiovascular disease and stroke: 2002 update: Consensus panel guide to comprehensive risk reduction for adult patients without coronary or other atherosclerotic vascular diseases. *Circulation. American Heart Association.* 106:388–391.

Pernefri. 2003. Konsensus Dialisis Perhimpunan Nefrologi Indonesia. Jakarta.

Podrid, P., 2004. C-Reactive Protein in Cardiovascular Disease-I, *in*: B.D Rose (editor), UptoDate 12.2 edition, Up to Date, Wellesley, M.A.

Porth, M.C., Matfin, G., 2009. Pathophysiology Concepts of altered Health States. 8th ed. Philadelphia : Lippincott Williams & Wilkins.

Pourfarzianu, V., Ghanbarpour, F., Nemati, E., Taheri, S., Einollahi, B. 2008. Laboratory variables and Treatment Adequacy in Hemodialysis Patient in Iran. *Saudi Journal of Nursing Scholarship Volume42.Issue 3. Pages 242249.*

Rashid, H., Haq R.ul, Rehman-ur., 2011. Comparison of C- Reactive protein levels with Delivered Dose of Kt/V in Patients with End- stage Renal Disease on Maintenance Hemodialysis. *Saudi J Kidney Dis Transpl* 2015;26(4):692-696.

Roesli,R.2005.Bila ginjal Aus <http://www.solusi.kesehatan.com/penyakitginjal/bila-ginjal-aus.html>.

Romão JE Jr, Haiashi AR, Elias RM, Luders C, Ferraboli R, Castro M C., et al.,2006. Positive acute-phase inflammatory markers indifferent stages of chronic kidney disease. *Am JNephrol*;26:59-66.

Santoso. 2010. Gagal Ginjal Kronik. <http://antiloans.org>

Schneider A, Drechsler C, Krane V, Krieter DH, Scharnagl H, Schneider MP, et al.,2012. The effect of high-flux hemodialysis on hemoglobin concentrations in patients with CKD: results of the MINOXIS study. *Clin J Am Soc Nephrol.*;7:52-9. doi: 10.1371/journal.pone.0128079.

Septiwi. C., 2010. Hubungan antara adekuasi hemodialisis dengan kualitas hidup pasien hemodialisis di unit hemodialisis RS. Prof. Dr. Margono Soekarjo Purwokerto. Tesis. Tidak dipublikasikan.

Simatupang, T. 2006. Gangguan cardiovascular pada penderita Ginjal. [http://litbang.depkes.go.id/aktual/kliping/ginjal 250406.htm](http://litbang.depkes.go.id/aktual/kliping/ginjal%20406.htm)

Sudigdo S. 2014. *Dasar- dasar Metodologi Penelitian klinis* . Edisi ke-5 . Sagung Seto hal 352- 387.

Suhardjono. 2014. Gagal Ginjal Kronik dalam *Buku ajar ilmu penyakit dalam*. edisi ke-4. Jilid I. Jakarta : Perhimpunan Dokter Spesialis Penyakit Dalam Indonesia.

Suharjono., 2010. Penderita gagal Ginjal di Indonesia. [222.ikcc.or.id/content.php?c=1&id=275](http://222.ikcc.or.id/content.php?c=1&id=275)

Suwitra, Ketut. 2014. Gagal Ginjal Kronik dalam *Buku ajar ilmu penyakit dalam*. edisi ke-4. Jilid I. Jakarta : Perhimpunan Dokter Spesialis Penyakit Dalam Indonesia.

Tielemans C, Husson C, schurmans T., 1996. Effect of Ultrapure and non- sterile dialysate on the inflammatory respon during in vitro hemodialysis. *International Society of Nephrology.*; 49 (1):236-43.

- Tsirpanlis, G., 2005. The pattern of inflamatio and a potential new clinical meaning and usefullness of C-reactive protein in end stage renal failure patients. *Kidney blood Press Res* 28 : 55-61.
- Vincenzo Panichi, M.D., Massimiliano Migliori, M.D., Stefano De Pietro, M.D., Daniele Taccola, M.D., Maria Rita Matelli Ph.D., Luca Giovannini, M.D., *et al.*, 2001. C-Reactive Protein in Patient With Chronic Renal disease. *Renal Failure*, London: Informa Healthcare, 23 (3&4) : 551-562.
- Wasse. 2007. Association of initial hemodialysis vascular Acces with Quality of Life. <http://cjasn.asnjournals.org>.
- Weihong W., Zhengcong D., Longqian Li., Jie Li., Xueqin Jin.,2016. Association of hyper-sensitive C-reactive protein with arterial stiffness and endothelial function in patients with hyperlipidemia., Xinjiang China; School of Medicine, Jiangnan University, Wuhan 430056, Hubei, China., *Int J Clin Exp Med.*, ;9(12):23416-2342.
- Weinhold B, Bader A, Valeria POLI, Rütehr U., 1997. Interleukin-6 is necessary, but not sufficient, for induction of the human C-reactive protein gene in vivo. *Biochem J* 325(3):617–21. doi:10.1042/bj3250617.
- Zadeh, K., & Kople, J.D., 2006. Inflammation in Renal Insufficiency. Up to Date 14.3.
- Zietz., B., Herfarth, H., Paul, G., 2003, Adiponectin Represents as Independent Cardiovascular Risk Factor Predicting Serum HDL Cholesterol Level in Type 2 Diabetes, *FEBS Lett*: 545: 103-4.