

## DAFTAR PUSTAKA

- Altman RP, Butchmiller TL. The jaundiced infant: biliary atresia. In: Grossfeld JL, O'Neill Jr JA, Fonkalsrud EW, Coran AG, editors. *Pediatric Surgery* 6<sup>th</sup> ed. Mosby Inc.; 2006.p.1603-19.
- Aulbach AD, Amuzie CJ. Biomarkers of Hepatobiliary Injury. In: *A Comprehensive Guide to Toxicology in Nonclinical Drug Development*. Second Ed. Elsevier Inc.; 2017.p.452-454.
- Balistreri WF, Grand R, Hoofnagle JH, *et al.* Biliary Atresia: Current Concepts And Research Directions. Summary Of A Symposium. *Hepatology* 1996;23:1682–92.
- Cabrera-Abreu JC, Green A. Gamma-glutamyltransferase: value of its measurement in paediatrics. *Ann Clin Biochem* 2002;39:22-5.
- Chardot C. Treatment for biliary atresia in 2003. *J Pediatr Gastroenterol Nutr* 2003;37:407–8.
- Chung PH, Wong KK, Tam PK. Predictors for failure after Kasai operation. *J Pediatr Surg* 2015 50:293–296.
- Chuslip S, Sookpotarom P, Tepmalai K, Rajatapiti P, Chongsrisawat V, Poovorawan Y, Vejchapipat. Prognostic values of serum bilirubin at 7th day post- Kasai for survival with native livers in patients with biliary atresia. *Pediatr surg Int* 2016 DOI 10.1007/s00383-016-3951-9.
- Desmet V. The cholangiopathies. In: Suchy FJ, ed. *Liver Disease in Children*. Philadelphia: Lippincott, Williams & Wilkins; 2001:39-62.
- El-Guindi MA, Sira MM, Sira AM, *et al.* Design and validation of a diagnostic score for biliary atresia. *J Hepatol* 2014;61:116-23.
- Goda T, Kawahara H, Kubota A *et al.* The most reliable early predictors of outcome in patients with biliary atresia after Kasai's operation. *J Pediatr Surg* 2013 48:2373–2377.
- Gunadi, Gunawan TA, Widiyanto G, Yuanita A, Mulyani NS, Makhmudi A. Liver transplan score for prediction of biliary atresia patients survival following Kasai procedure. *BMC Res Notes*, 2018, 11: 381.
- Fischler B, Haglund B, Hjern A. A population-based study on the incidence and possible pre- and perinatal etiologic risk factors of biliary atresia. *J Pediatr* 2002;141:217-22.
- Hartley JL, Davenport M, Kelly DA. Biliary atresia. *Lancet* 2009 374(9702):1704–1713 [CrossRefPubMedGoogle Scholar](#).
- Hadzic N (2009). Biliary atresia. *Acta Medica Academica* 2009;38:92-103.
- Haafiz AB. Liver fibrosis in biliary atresia. *Expert Rev Gastroenterol Hepatol* 2010;4:335–43.
- Hand NJ, Horner AM, Master ZR, *et al.* MicroRNA profiling identifies miR-29 as a regulator of disease-associated pathways in experimental biliary atresia. *J Pediatr Gastroenterol Nutr* 2012;54:186–92.
- Ian D. D'Agata, MD and William F. Balistreri, MD. Evaluation of Liver Disease in the Pediatric Patient. *Pediatrics in Review*, 1999. Vol.20 No.11

- Ihn, K., Ho, I. G., Chang, E. Y., & Han, S. J. (2018). Correlation between gamma-glutamyl transpeptidase activity and outcomes after Kasai portoenterostomy for biliary atresia. *Journal of Pediatric Surgery*, 53(3), 461–467. doi:10.1016/j.jpedsurg.2017.10.001
- Keplinger KM, Bloomston M. Anatomy and embryology of biliary tract. *Surgical Clinics* 94 (2), 2014; 203-217.
- Khalil BA, Thamara M, Perera PR, Mirza DF. Clinical practice: management of biliary atresia. *Eur J Pediatr* 2010;169:395-402.
- Khanna K, Bhatnagar V, Agarwala S, Srinivas M, Gupta SD. Ratio of Preoperative and Postoperative Serum Bilirubin Levels Predicts Early Outcome Following Biliary Atresia Surgery. *J Indian Assoc of Pediatr Surg*, 2018 vol 23.Issue 2
- Koga, M Wada, H Nakamura, et al. Factors influencing jaundice-free survival with the native liver in post-portoenterostomy biliary atresia patients: Results from a single institution. *J Pediatr Surg* 2013;48:2368-2372.
- Liu CS, Chin TW, Wei CF. Value of gamma-glutamyl transpeptidase for early diagnosis of biliary atresia. *Zhonghua Yi Xue Za Zhi (Taipei)* 1998;61:716-20.
- Meyers RL, Book LS, O'Gorman MA, et al. High-dose steroids, ursodeoxycholic acid, and chronic intravenous antibiotics improve bile flow after Kasai procedure in infants with biliary atresia. *J Pediatr Surg* 2003;38:406–11.
- Ohi R. Biliary atresia. A surgical perspective. *Clin Liver Dis* 2000;4:779–804.
- Ogawa T, Enomoto M, Fujii H, et al. MicroRNA-221/222 upregulation indicates the activation of stellate cells and the progression of liver fibrosis. *Gut* 2012;61:1600–9.
- Pakarinen MP, Rintala RJ. Surgery of biliary atresia. *Scand J Surg* 2011;100:49–53.
- Qiao, G., Li, L., Cheng, W., Zhang, Z., Ge, J., & Wang, C. Conditional probability of survival in patients with biliary atresia after Kasai portoenterostomy: a Chinese population-based study. *Journal of pediatric surgery*, 2015: 50(8): 1310-1315
- Rodeck B, Becker AC, Gratz KF, Petersen C. Early predictors of success of Kasai operation in children with biliary atresia. *Eur J Pediatr Surg* 2007 17:308–312
- Schreiber RA, Barker CC, Roberts EA et al. Biliary atresia:the Canadian experience. *J Pediatr* 2007 151:659–665.
- Schwarz SM, Cuffari C, Windle ML, Guandalini S, Vargas JH, Pediatric Biliary Atresia Workup.Medscape, update Sep, 2017.
- Shen W, Chen G, Dong R, Zhao R, Zheng S. MicroRNA-21/PTEN/Akt axis in the fibrogenesis of biliary atresia. *J Pediatr Surg*. 2014;49(12):1738-41.
- Shneider BL, Brown MB, Haber B et al. A multicenter study of the outcome of biliary atresia in the United States, 1997 to 2000. *J Pediatr* 2006 148:467–474.
- Sinha CK, Davenport M. Biliary atresia. *J Indian Assoc Pediatr Surg*;2008 13(Issue 2):4956
- Suzuki T., Hashimoto T., Hussein MH., Hara F., Hibi M., Kato T., Biliary Atresia type I cyst and choledochal cyst: can we differentiated or not ? *J Hepatobiliary Pancreat Sci* (2013) 20:465-470.DOI 10.1007/s00534-013-0605-3.
- Sokol RJ, Mack C, Narkewicz MR, Karrer FM. Pathogenesis and outcome of biliary atresia: current concepts. *J Pediatr Gastroenterol Nutr* 2003;37:4–21.
- Wei Lai M, Hwei Chang M, Chien Hsu S, Chi Shu H, Tau su C, Lian Kao C, Yun Lee C. Differential diagnosis of extrahepatic atresia from neonatal

hepatitis: a prospective study. [J Pediatr Gastroenterol Nutr.](#) 1994 Feb;18(2):121-7.

Yan K.J. Nonparametric Predictive Inference with Right-Censored Data. 2002. Thesis Ph.D. University of Durham.

Zhou L., Wang W., Shan Q., Liu B., Zheng Y, et al. Optimizing the US Diagnosis of Biliary Atresia with a Modified Triangular Cord Thickness and Gallbladder Classification, May 6 2015 <https://doi.org/10.1148/radiol.20151422309>