

DAFTAR PUSTAKA

- Ahmed, M. 2018. Acute cholangitis - an update. *World J Gastrointestinal Pathophysiol*, 9(1): 1–7.
- Buzzetti, E., Parikh, P.M., Gerussi, A. & Tsochatzis, E. 2017. Gender differences in liver disease and the drug-dose gender gap. *Pharmacological Res*, 120: 97–108.
- Chiu, C.Y., Chen, P.H., Chan, C.F., Chang, M.H. & Wu, T.C. 2013. Biliary atresia in preterm infants in Taiwan: A nationwide survey. *J Pediatr*, 163(1): 1–5.
- Davenport, M. 2006. Biliary atresia: outcome and management. *Indian j ped*, 73(9 PG-825–8): 825–828.
- Dimitrova, N., Zamudio, J.R., Jong, R.M., Soukup, D., Resnick, R., Sarma, K., *et al.* 2011. The anatomic pattern of biliary atresia identified at time of Kasai hepatportoenterostomy and early postoperative clearance of Jaundice are significant predictors of transplant-free survival. *Ann Surg*, 32(7): 736–740.
- Ernest, L., Saing, H. & Pau, K. 2003. Cholangitis after hepatic portoenterostomy for biliary atresia : a multivariate analysis of risk factors. *J Pediatr*, 142(566): 71.
- Ghazy, R.M., Adawy, N.M., Khedr, M.A. & Tahoun, M.M. 2018. Biliary atresia recent insight. *Egyptian Pediatric Association Gazette*, 66(1): 1–8.
- Goda, T., Kawahara, H., Kubota, A., Hirano, K., Umeda, S., Tani, G., *et al.* 2013. The most reliable early predictors of outcome in patients with biliary atresia after Kasai’s operation. *J Pediatr Surg*, 48(12): 2373–2377.
- Gustomi S, Makhmudi A, Gunadi, D.A. 2017. Faktor Prognostik Terjadinya Kolangitis Pasca Prosedur Kasai di RSUP Dr. Sardjito Yogyakarta. Fakultas

Kedokteran Universitas Gadjah Mada.

- Guy, J. & Peters, M.G. 2013. Liver disease in women: The influence of gender on epidemiology, natural history, and patient outcomes. *Clin Gastroenterol and Hepatol*, 9(10): 633–639.
- Jiang, H., Gao, P., Chen, H., Zhong, Z., Shu, M., Zhang, Z., *et al.* 2018. The prognostic value of CD8+ and CD45RO+ T cells infiltration and beclin1 expression levels for early postoperative cholangitis of Biliary atresia patients after Kasai operation. *J Korean Med Sci*, 33(30): 1–12.
- Lampela, H. 2013. Biliary atresia treatment results and native liver function. Children’s Hospital Institute of Clinical Medicine, University of Helsinki and National Graduate School of Clinical Investigation Helsinki.
- Li, D., Chen, X., Fu, K., Yang, J. & Feng, J. 2017. Preoperative nutritional status and its impact on cholangitis after Kasai portoenterostomy in biliary atresia patients. *Pediatr Surg Int*, 33(8): 901–906.
- Lünzmann, K. & Schweizer, P. 1999. The influence of cholangitis on the prognosis of extrahepatic biliary atresia. *Eur J Pediatr Surg*, 9(1): 19–23.
- Luo, Y. & Zheng, S. 2008. Current concept about postoperative cholangitis in biliary atresia. *World J Pediatr*, 4(1): 14–19.
- Maheranny, M. 2013. Faktor-faktor yang mempengaruhi angka keberhasilan portoenterostomi Kasai. Fakultas Kedokteran. Universitas Indonesia.
- Mahmud, M.N., Thambidot, C.R., Zakaria, Z., Mohan, N., Sheila, M., Cpaeds, M.M., *et al.* 2002. Jaundice clearance and cholangitis in the first year following portoenterostomy for biliary atresia. *Med J Malaysia*, 57(1): 92–96.
- Moghtaderi, M., Gorji, M., Farahmand, F., Fallahi, G. & Ashjal, B. 2016. Long-term outcome of children with biliary atresia after Kasai surgery in Iranian

infants. *Pediatrics & Therapeutics*, 06(01): 2–5.

Mosler, P. 2011. Diagnosis and management of acute cholangitis. *Curr Gastroenterol Rep*, 13(2): 166–172.

Namaseyam, D. & Nallusamy, M. 2017. Factors influencing outcome after hepatic portoenterostomy among extrahepatic bile duct atresia patients in hospital Sultanah Bahiyah, Alor Setar. *Med J Malaysia*, 72(6): 329–332.

Nio, M., Ohi, R., Miyano, T., Saeki, M., Shiraki, K. & Tanaka, K. 2003. Five- and 10-year survival rates after surgery for biliary atresia: A report from the Japanese Biliary Atresia Registry. *J Pediatr Surg*, 38(7): 997–1000.

Sanghai, S.R., Shah, I., Bhatnagar, S. & Murthy, A. 2009. Incidence and prognostic factors associated with biliary atresia in western India. *Annals of hepatology*, 8(2): 120–2.

Scott. E., L., Janet., P., Michael., G., Andrew.A., K. & Michael.D., K. 2000. Bile duct size does not predict success of portoenterostomy for biliary atresia. *J Pediatr Surg*, 35(6): 1006–1007.

Tainaka, T., Kaneko, K., Seo, T., Ono, Y., Ogura, Y., Wada, H., *et al.* 2006. Hepatolithiasis after hepatic portoenterostomy for biliary atresia. *J Pediatr Surg*: 808–811.

Wang, K.S. 2015. Newborn screening for biliary atresia. *Pediatrics*, 136(6): e1663–e1669.

Webb, N.L., Jiwane, A., Ooi, C.Y., Nightingale, S., Adams, S.E. & Krishnan, U. 2017. Clinical significance of liver histology on outcomes in biliary atresia. *J Paediatr Child Health*, 53(3): 252–256.

Wong, K.K.Y., Fan, A.H., Lan, L.C.L., Lin, S.C.L. & Tam, P.K.H. 2004. Effective antibiotic regime for postoperative acute cholangitis in biliary atresia - An evolving scene. *J Pediatric Surg*, 39(12): 1800–1802.

Wu, E., Chen, H., Ni, Y., Lee, P., Hsu, H., Lai, H., *et al.* 2002. Bacterial cholangitis in patients with biliary atresia: impact on short-term outcome. *Pediatr Surg Int*, 17(5–6): 390–395.