

DAFTAR PUSTAKA

- Achmad. 2012. *Jamur*. Agriflo. Bogor.
- Adamopoulos, S., Voulgaridis, E., and Passialis, C. 2005. Variation of certain chemical properties within the stemwood of black locust (*Robinia pseudoacacia* L.). *Holz als roh-und Werkstoff*. 63(5): 327-333.
- Alonso, M., Rozados, M.J., Vega, J.A., Perez-Gorostiaga, P., Cuinas, P., Fonturbel, M.T., and Fernandez, C. 2002. Biochemical responses of *Pinus pinaster* trees to fire-induced trunk girdling and crown scorch: secondary metabolites and pigments as needle chemical indicators. *Journal of Chemical Ecology*. 28(4): 687-700.
- Anda, R.R., Koch, G., Richter, H.G., Talavera, F.J.F., Guzmán, J.A.S., and Satyanarayana, K.G. 2019. Formation of heartwood, chemical composition of extractives and natural durability of plantation-grown teak wood from Mexico. *Holzforchung*. 73(6): 547-557.
- ASTM. 1989. Annual book of ASTM standards. *Section Four Construction Volume 04.09 wood*. Philadelphia.
- ASTM. 2002. Annual book of ASTM standards. *Section Four Construction Volume 04.10 Wood*. Baltimore.
- Ayadi, N., Lejeune, F., Charrier, F., Charrier, B., and Merlin, A. 2003. Color stability of heat-treated wood during artificial weathering. *Holz als Roh-und Werkstoff*. 61(3): 221-226.
- Baeza, J. 2001. Chemical characterization of wood and its components. *Wood and Cellulosic Chemistry*. Second edition. Hon, D.N.S., and Shiraishi, N. Clemson. South Carolina. Clemson University, Japan. Kyoto University.
- Baly, E. C. 1937. Photosynthesis of carbohydrates in vitro. *Nature*. cxi 930.
- Basri, E., and Wahyudi, I. 2013. Sifat dasar kayu Jati Plus Perhutani dari berbagai umur dan kaitannya dengan sifat dan kualitas pengeringan. *Jurnal Penelitian Hasil Hutan*. 31(2): 93-102.
- Basri, E., Yuniarti, K., Wahyudi, I., and Damayanti, R. 2015. Effects of girdling on wood properties and drying characteristics of *Acacia mangium*. *Journal of Tropical Forest Science*. 27(4): 498-505.
- Bhat, K.M., Thulasidas P.K., Florence, E.J.M., and Jayaraman, K. 2005. Wood durability of home-garden teak against brown-rot and white-rot fungi. *Trees*. 19: 654-660.
- Bhat, K.M., and Florence, E.J.M. 2003. Natural decay resistance of juvenile teak wood grown in high input plantations. *Holzforchung*. 57: 453-455.
- Caron, A., Altaner, C.M., Gardiner, B., and Jarvis, M.C. 2013. Distribution of extractives in Sitka spruce (*Picea sitchensis*) grown in the northern UK. *European Journal of Wood and Wood Products*. 71(6): 697-704.

- Chirinos, R., Betalleluz-Pallardel, I., Huamán, A., Arbizu, C., Pedreschi, R., and Campos, D. 2009. HPLC-DAD characterisation of phenolic compounds from Andean oca (*Oxalis tuberosa* Mol.) tubers and their contribution to the antioxidant capacity. *Food Chemistry*. 113(4): 1243-1251.
- Cordes, J.W.H. 1992. Hutan jati di Jawa. Perum Perhutani Unit II. Malang Jawa Timur.
- Datta, S.K., and Kumar, A. 1987. Histochemical studies of the transition from sapwood to heartwood in *Tectona grandis*. *IAWA Journal*. 8(4): 363-368.
- Deslauriers, A., Giovannelli, A., Rossi, S., Castro, G., Fragnelli, G., and Traversi, L. 2009. Intra-annual cambial activity and carbon availability in stem of poplar. *Tree physiology*. 29(10): 1223-1235.
- Fahey, T.D., Snellgrove, T.A., and Plank, M.E. 1986. Changes in product recovery between live and dead Lodgepole pine: a compendium. USDA, Forest Service, *Pacific Northwest Research Station*, Portland, OR. Pp. 1–25.
- Febrianto, F., Syafii, W., dan Barata, A. 2000. Keawetan alami kayu jati (*Tectona grandis* L.f.) pada berbagai kelas umur. *Jurnal Teknologi Hasil Hutan*. 13(2): 25-33.
- Fengel, D., and Wegener. 1989. Wood chemistry, ultrastructure, reactions. New York.
- Fernandes, F.H.A., and Salgado, H.R.N. 2016. Gallic acid: review of the methods of determination and quantification. *Critical Reviews in Analytical Chemistry*. 46(3): 257-265.
- Gao, H., Shupe, T.F., Hse, C.Y., and Eberhardt, T.L. 2006. Antioxidant activity of extracts from the bark of *Chamaecyparis lawsoniana* (A. Murray) Parl. *Holzforschung*. 60(4): 459-462.
- Gierlinger, N., Jacques, D., Grabner, M., Wimmer, R., Schwanninger, M., Rozenberg, P., and Pâques, L.E. 2004. Colour of larch heartwood and relationships to extractives and brown-rot decay resistance. *Trees*. 18(1): 102-108.
- Giovannelli, A., Emiliani, G., Traversi, M.L., Deslauriers, A., and Rossi, S. 2011. Sampling cambial region and mature xylem for non-structural carbohydrates and starch analyses. *Dendrochronologia*. 29(3): 177-182.
- Gominho, J., Figueira, J., and Rodrigues, J.C. 2001. Within-tree variation of heartwood, extractives and wood density in the eucalypt hybrid urograndis. *Wood and Fiber Science*. 33(1): 3-8.
- Gominho, J., Lourenço, A., Miranda, I., and Pereira, H. 2015. Radial and axial variation of heartwood properties and extractives in mature trees of *Eucalyptus globulus*. *BioResources*. 10(1): 721-731.
- Hanes, C. S. 1937. The action of amylases in relation to the structure of starch. Parts I-VIII. *New Phyt.* xxxvi.
- Hauch, S., and Magel, E. 1998. Extractable activities and protein content of sucrose-phosphate synthase, sucrose synthase and neutral invertase in trunk tissues

- of *Robinia pseudoacacia* L. are related to cambial wood production and heartwood formation. *Planta*. 207(2): 266-274.
- Haupt, M., Leithoff, H., Meier, D., Puls, J., Richter, H.G., and Faix, O. 2003. Heartwood extractives and natural durability of plantation-grown teakwood (*Tectona grandis* L.)—a case study. *Holz als Roh-und Werkstoff*. 61(6): 473-474.
- Hillis, W.E. 1987. Heartwood and tree exudates. Springer-Verlag Berlin Heidelberg. Germany.
- Huang, D., Ou, B., and Prior, R.L. 2005. The chemistry behind antioxidant capacity assays. *Journal of Agricultural and Food Chemistry*. 53(6): 1841-1856.
- Humphreys, F.R., and Kelly, J. 1961. The determination of starch in wood. *Anal. Chim. Acta*. 24: 66-70. Dalam Lambert, M. Methods for chemical analysis. *Forestry Commission of N.S.W.* Technical paper no. 25.
- Irawati, D., Azwar, N.R., Syafii, W., dan Artika, I.M. 2009. Pemanfaatan serbuk kayu untuk produksi etanol dengan perlakuan pendahuluan delignifikasi menggunakan jamur *Phanerochaete chrysosporium*. *Jurnal Ilmu Kehutanan*. 3(1): 13-22.
- Irawati, R.H., dan Purnomo, H. 2012. Pelangi di tanah kartini: kisah aktor mebel jepara bertahan dan melangkah ke depan. CIFOR, Bogor, Indonesia.
- Kelsey, R.G., Hennon, P.E., Huso, M., and Karchesy, J.J. 2005. Changes in heartwood chemistry of dead yellow-cedar trees that remain standing for 80 years or more in southeast Alaska. *Journal of Chemical Ecology*. 31(11): 2653-2670.
- Koch, P. 1996. Lodgepole pine in North America Madison, Wisconsin, *Forest Products Society*.
- Khoddami, A., Wilkes, M.A., and Roberts, T.H. 2013. Techniques for analysis of plant phenolic compounds. *Molecules*. 18(2): 2328-2375.
- Koyani, R.D., and Rajput, K.S. 2014. Light microscopic analysis of *Tectona grandis* Lf wood inoculated with *Irpex lacteus* and *Phanerochaete chrysosporium*. *European Journal of Wood and Wood Products*. 72(2): 157-164.
- Labosky J.P., Baldwin, R.C., and Zhang, J. 1990. Chemical composition of gypsy moth-killed red oak. *Wood and Fiber Science*. 22(2): 217-226.
- Lukmandaru, G. 2011. Variability in the natural termite resistance of plantation teak wood and its relations with wood extractive content and color properties. *Journal of Forestry Research*. 8(1): 17-31.
- Lukmandaru, G. 2013. Antifungal activities of certain components of teak wood extractives. *Jurnal Ilmu dan Teknologi Kayu Tropis*. 11(1): 11-18.
- Lukmandaru, G. and Ogiyama, K., 2005a. Bioactive compounds from ethyl acetate extract of teakwood (*Tectona grandis* L.f.), *Proceedings of the 6th International Wood Science Symposium LIPI-JSPS Core*, Bali, Indonesia, August 29–31, 2005, pp. 346–350.

- Lukmandaru, G. and Ogiyama, K., 2005b. Bioactive extract from teakwood (*Tectona grandis* L.f.), *Proceedings of International Symposium on Wood Science and Technology*, Volume II: poster presentations. Yokohama, Japan, November 27–30, 2005, pp. 413–414.
- Lukmandaru, G and Takahashi, K. 2008. Variation in the natural termite resistance of teak (*Tectona grandis* L.f.) as a function of tree age. *Annals of Forest Science*. 65(7): 708 p1 - p8.
- Lukmandaru, G and Takahashi, K. 2009. Radial distribution of quinones in plantation teak (*Tectona grandis* L.f.). *Annals of Forest Science*. 66(6): 605 p1 – p9.
- Lukmandaru, G., Sayudha, I.G.N.D., Gustomo, L.S., dan Prasetyo, V.E. 2011. Pengukuran kadar ekstraktif dan sifat warna kayu Acacia mangium dari lima provenans. *Prosiding Seminar Nasional MAPEKI XIII*, Bali. Hlm: 372-380.
- Lukmandaru, G., Wargono, P., Mohammad, A.R., dan Prasetyo, V.E. 2018. Studi mutu kayu jati di hutan rakyat Gunungkidul. VII. Ketahanan terhadap rayap tanah. *Jurnal Ilmu Kehutanan*, 12(1), 22-39.
- Martawijaya, A., Kartasudjana, I., Kadir, K., dan Amongprawira, S. 2005. Atlas Kayu Indonesia Jilid I. Balai Penelitian Hasil Hutan. Badan Litbang Kehutanan. Bogor. 42-47.
- Michelot, A., Simard, S., Rathgeber, C., Dufrêne, E., and Damesin, C. 2012. Comparing the intra-annual wood formation of three European species (*Fagus sylvatica*, *Quercus petraea* and *Pinus sylvestris*) as related to leaf phenology and non-structural carbohydrate dynamics. *Tree physiology*. 32(8): 1033-1045.
- Mihara, R., Barry, K.M., Mohammed, C.L., and Mitsunaga, T. 2005. Comparison of antifungal and antioxidant activities of *Acacia mangium* and *A. auriculiformis* heartwood extracts. *Journal of Chemical Ecology*. 31(4): 789-804.
- Miranda, I., Sousa, V., and Pereira, H. 2011. Wood properties of teak (*Tectona grandis*) from a mature unmanaged stand in East Timor. *Journal of Wood Science*. 57(3): 171-178.
- Moya, R., and Berrocal, A. 2010. Wood colour variation in sapwood and heartwood of young trees of *Tectona grandis* and its relationship with plantation characteristics, site, and decay resistance. *Annals of Forest Science*. 67(1): 109.
- Mukhortova, L.V., Kirdyanov, A.V., Myglan, V.S., and Guggenberger, G. 2009. Wood transformation in dead-standing trees in the forest-tundra of central Siberia. *Biology Bulletin*. 36(1): 58-65.
- Neverova, N.A., Levchuk, A.A., Ostroukhova, L.A., Medvedeva, E.N., Onuchina, N.A., and Babkin, V.A. 2013. Distribution of extractive substances in wood of the Siberian larch (*Larix sibirica* Ledeb.). *Russian Journal of Bioorganic Chemistry*. 39(7): 712-719.

- Niamké F.B., Amusant N., Charpentier J.P., Chaix, G., Baissac, Y., Boutahar, N., Adima, A.A., Coulibaly, S.K., and Allemand, C.J. 2011. Relationships between biochemical attributes (non-structural carbohydrates and phenolics) and natural durability against fungi in dry teak wood (*Tectona grandis* L. f.). *Annals of Forest Science*. 68: 201–211.
- Niamke, F.B., Amusant, N., Kokutse, A., Chaix, G., Charpentier, J.P., Adima, A.A., and Jay-Allemand, C. 2010. Radial distribution of non-structural carbohydrates in Malaysian teak. *International Journal of Biological and Chemical Sciences*. 4(3): 710-720.
- Niamke, F.B., Adima, A.A., Seraphin, K.C., Amusant, N., and Jay-Allemand, C. 2018. Heartwood formation process in teak (*Tectona grandis* L.f): fate of non-structural carbohydrates and characterization of forsythoside B. *International Journal of Biological and Chemical Sciences*. 12(3): 1102-1112.
- Nobuchi, T., Okada, N., Nishida, M., Siripatanadilok, S., Veenin, T., Tobing, T. L., and Sahri, M.H. 2005. Some characteristics of wood formation in Teak (*Tectona grandis*) with special reference to water conditions. *Quality Timber Products of Teak from Sustainable Forest Management*. India. 495-499.
- Noel, A.R.A. 1970. The girdled tree. *The botanical review*. 36(2): 162.
- Oberhuber, W., Swidrak, I., Pirkebner, D., and Gruber, A. 2011. Temporal dynamics of nonstructural carbohydrates and xylem growth in *Pinus sylvestris* exposed to drought. *Canadian Journal of Forest Research*. 41(8): 1590-1597.
- Oliveira, L.S., Santana, A.L., Maranhão, C.A., Rita de Cássia, M., de Lima, V.L.A.G., Da Silva, S.I., Nascimento, M.S., and Bieber, L. 2010. Natural resistance of five woods to *Phanerochaete chrysosporium* degradation. *International Biodeterioration & Biodegradation*. 64(8): 711-715.
- Paloheimo, L. 1930. Zur Verwendbarkeit des Jodkolorimetrischen Prinzips bei StSrkebestimmungen. *Biochem. Zeitschr. ccrrii*. 150.
- Pandey, K.K., and Pitman, A.J. 2003. FTIR studies of the changes in wood chemistry following decay by brown-rot and white-rot fungi. *International Biodeterioration & Biodegradation*. 52(3): 151-160.
- Parkin, E.A. 1938. The depletion of starch from timber in relation to attack by *Lyctus beetles*: II. A preliminary experiment upon the effect of girdling standing oak trees. *Forestry: An International Journal of Forest Research*. 12(1): 30-37.
- Paul, E.A. 1992. Organic Matter Decompositionn. *Encyclopedia of Microbiology*, Vol.3. Academic Press. Inc.
- Perhutani. 2011. Monitoring dan evaluasi pengembangan jati plus perhutani (JPP) di KPH Kendal. Kendal: Perum Perhutani KPH Kendal.
- Perhutani. 2014. www.puslitbangperhutani.com
- Perhutani. 2014. Suara Rimba. <http://www.bumn.go.id/perhutani/halaman/144>

- Perhutani. 2016. Statistik Perum Perhutani 2011-2015. Perum Perhutani. Jakarta.
- Pujiono, S. 2014. Produksi bibit jati unggul dari klon dan budidaya. IPB Press.
- Phillips, E.W.J. 1938. The depletion of starch from timber in relation to attack by *Lyctus Beetles*: I. Starch, with special reference to its occurrence in timber. *Forestry: An International Journal of Forest Research*. 12(1): 15-29.
- Polato, R., Laming, P.B., and Sierra-Alvarez, R. 2005. Assessment some wood characteristics of teak of brazilian origin. Quality timber products of teak from sustainable forest management. *Proceedings of the International Conference on Quality Timber Products of Teak Sustainable Forest Management*. Peechi, India (pp. 2-5).
- Pometti, C.L., Palanti, S., Pizzo, B., Charpentier, J.P., Boizot, N., Resio, C., and Saidman, B.O. 2010. Durability of five native Argentine wood species of the genera *Prosopis* and *Acacia* decayed by rot fungi and its relationship with extractive content. *Biodegradation*. 21(5): 753-760.
- Pramasari, D.A., Wahyuni, I., Adi, D.S., Amin, Y., Darmawan, T., and Dwianto, W. 2014. Effect of age on chemical component of platinum teak wood—a fast growing teak wood from LIPI. In *The 6th International Symposium of Indonesian Wood Research Society* (p. 211).
- Prawirohatmodjo, S. 2004. Kimia kayu. Universitas Gadjah Mada. Yogyakarta.
- Purwanta, S., Sumantoro, P., Setyaningrum, H.D., dan Saparinto, C. 2015. Budidaya dan bisnis kayu jati. Penebar Swadaya, Jakarta.
- Retnaningrum, N.I. 2012. Studi keawetan alami terhadap rayap dihubungkan dengan komposisi ekstraktif kayu jati dari hutan rakyat Kulon Progo. Skripsi. Fakultas Kehutanan UGM. Yogyakarta (Tidak diterbitkan).
- Rini, D.S. dan Marsoem, S.N. 2013. Pengaruh lama peneerasan terhadap sifat fisika kayu jati (*Tectona grandis* Lf) hutan rakyat (Master thesis, Universitas Gadjah Mada).
- Rizanti, D.E., Darmawan, W., George, B., Merlin, A., Dumarcay, S., Chapuis, H., and Syafii, W. 2018. Comparison of teak wood properties according to forest management: short versus long rotation. *Annals of Forest Science*. 75(2): 39.
- Rowell, R.M., Pettersen, R., Han, J.S., Rowell, J.S., and Tshabalala, M.A. 2005. Cell wall chemistry. Handbook of wood chemistry and wood composites, 2.
- Shamoun, S.F., and Levi, M.P. 1985. A chemical and microscopic study of decayed earlywood and latewood of loblolly pine killed by the Southern pine beetle. *Wood and Fiber Science*. 17(1): 22-28.
- Shrimpton, D.M. 1973. Extractives associated with wound response of Lodgepole pine attacked by the mountain pine beetle and associated with microorganisms. *Canadian J. Botany*. 51: 527–533.
- Simard, S., Giovannelli, A., Treydte, K., Traversi, M.L., King, G.M., Frank, D., and Fonti, P. 2013. Intra-annual dynamics of non-structural carbohydrates in the

cambium of mature conifer trees reflects radial growth demands. *Tree Physiology*. 33(9): 913-923.

- Sirmah, P., Iaych, K., Poaty, B., Dumarçay, S., and Gerardin, P. 2009. Effect of extractives on durability of *Prosopis juliflora* heartwood. *International Research Group on Wood Protection*. Doc. No.: IRG/WP, 09-30518.
- Soenarno, S., dan Idris, M.M. 1990. Pengaruh lama teresan dan diameter tegakan jati terhadap persentase kerusakan kayu. *Jurnal Penelitian Hasil Hutan*. 7(3).
- Spoehr, H.A., and Milner, H.W. 1935. Leaf starch: its isolation and some of its properties. *J. Biol. Chem. ad.* (3): 679.
- Srinivasan, C., Dsouza, T.M., Boominathan, K., and Reddy, C.A. 1995. Demonstration of laccase in the white rot basidiomycete *Phanerochaete chrysosporium* BKM-F1767. *Appl. Environ. Microbiol.* 61(12): 4274-4277.
- Sumarni, G., dan Muslich, M. 2008. Kelas awet jati cepat tumbuh dan jati konvensional pada berbagai umur pohon. *Jurnal Penelitian Hasil Hutan*. 26(4): 342-351.
- Sumthong, P., Damveld, R.A., Choi, Y.H., Arentshorst, M., Ram, A.F., van den Hondel, C.A., and Verpoorte, R. 2006. Activity of quinones from teak (*Tectona grandis*) on fungal cell wall stress. *Planta Medica*. 72(10): 943-944.
- Taylor, A., and Cooper, P. 2002. The effect of stem girdling on wood quality. *Wood and Fiber Science*. 34(2): 212-220.
- Thulasidas, P.K., and Bhat, K.M. 2007. Chemical extractive compounds determining the brown-rot decay resistance of teak wood. *Holz als Roh-und Werkstoff*. 65(2): 121-124.
- Umezawa, T. 2001. Chemistry of extractives. *Wood and Cellulosic Chemistry*. Second edition. Hon, D.N.S., and Shiraishi, N. Clemson. South Carolina. Clemson University, Japan. Kyoto University.
- Walter, M., Boul H.L., Chong R., and Ford C. 2002. Biodegradation of PCP in contaminated soil by NewZealand white-rot fungi. Submitted to *Aust. J. Soil Res.*
- Wangard, F.F. 1966. Resistance of wood to chemical degradation. *Forest Product Journal*. 16(2): 53-64.
- Willför, S., and Holmbom, B. 2004. Isolation and characterisation of water soluble polysaccharides from Norway spruce and Scots pine. *Wood Science and Technology*. 38(3): 173-179.
- Willför, S., Sundberg, A., Pranovich, A., and Holmbom, B. 2005. Polysaccharides in some industrially important hardwood species. *Wood Science and Technology*. 39(8): 601-617.
- Windeisen, E., Klassen, A., and Wegener, G. 2003. On the chemical characterisation of plantation teakwood from Panama. *Holz als roh-und werkstoff*. 61(6): 416-418.

- Windeisen, E., Wegener, G., Lesnino, G., and Schumacher, P. 2002. Investigation of the correlation between extractives content and natural durability in 20 cultivated larch trees. *European Journal of Wood and Wood Products*. 60(5): 373-374.
- Wissam, Z., Ghada, B., Wassim, A., and Warid, K. 2012. Effective extraction of polyphenols and proanthocyanidins from pomegranate's peel. *International Journal of Pharmacy and Pharmaceutical Sciences*. 4(3): 675-682.
- Woo, K.L., Watson, P., and Mansfield, S.D. 2005. The effects of mountain pine beetle attack on lodgepole pine wood morphology and chemistry: Implications for wood and fiber quality. *Wood and Fiber Science*. 37(1): 112-126.
- Yunanta, R.R.K., Lukmandaru, G., dan Fernandes, A. 2014. Sifat kimia dari kayu *Shorea retusa*, *Shorea macroptera*, dan *Shorea macrophylla*. *Jurnal Penelitian Ekosistem Dipterokarpa*. 8(1): 15-24.