

## DAFTAR PUSTAKA

- Adedapo, A.A., Mogbojuri, O.M., Emikpe, B.O. (2009). Safety evaluations of the aqueous extract of the leaves of *Moringa oleifera* in rats. *Journal of Medicinal Plants Research*, 3(8): 586–591.
- Anief, M. (1990). *Perjalanan dan Nasib Obat Dalam Badan*. (pp.5-10). Yogyakarta: UGM Press.
- Awodele, O., Oreagba, I. A., Odoma, S., da Silva, J. A., Osunkalu, V.O. (2012). Toxicological evaluation of the aqueous leaf extract of *Moringa oleifera* lam. (*moringaceae*). *Journal of Ethnopharmacology*. 139(2): 330-336.
- Bandelow, B., Zohar, J., Hollander, E., Kasper, S., Mo"ller, H.J. (2002). Guidelines for the pharmacological treatment of anxiety, obsessive-compulsive and posttraumatic disorders. *World J Biol Psychiatry*. 3(4):171-199
- Bhat, S.K., Joy, A.E., (2014). Antianxiety effect of ethanolic extract of leaves of *Moringa oleifera* in swiss albino mice. *Arch Med Health Science*, 2(1):5-7.
- Blumstein, L.K., Crawley, J.N. (1983). Further characterisation of a simple, automated exploratory model for the anxiolytic effects of benzodiazepines. *Pharmacol. Biochem. Behav*, 18(1):37 – 40.
- Borsini, F., Podhorna, J., Marazziti, D. (2002). Do animal models of anxiety predict anxiolytic-like effects of antidepressants?, *Psychopharmacology*, 163(2):121–41.
- Bystritsky, A., Khalsa, S.S., Cameron, M.E., & Schiffman, J. (2013). Current diagnosis and treatment of anxiety disorders. *P & T : a peer-reviewed journal for formulary management*, 38(1), 30–57.
- Calabrese, E.J., (2008). An assessment of anxiolytic drug screening tests: hormetic dose responses predominate. *Crit Rev Toxicol*, 38(6):489-542
- Craske, M.G., Rauch, S.L., Ursano, R., Prenoveau, J., Pine, D.S., Zinbarg, R.E. (2009). What is an anxiety disorder?. *Depress Anxiety*, 26(12):1066-1085.
- Craske, M.G., Stein, M.B. (2016). Anxiety. *Lancet*, 388(10063):3048-3059
- Crawley, J.N., Davis, L.G. (1982). Base line exploratory activity predicts anxiolytics responsiveness to diazepam in five mouse strains. *Brain Res. Bull*, 8(6): 609 – 612.

- Crawley, J.N. (2007). Emotional behaviors: animal models of psychiatric diseases. In Crawley, J.N., *What's Wrong With My Mouse? Behavioral phenotyping of transgenic and knockout mice*. 2<sup>nd</sup> ed. (pp. 226-265). New York : John Wiley and Sons.
- Cryan, J.F., Sweeney, F.F. (2011). The age of anxiety: role of animal models of anxiolytic action in drug discovery. *Br J Pharmacol*, 164(4): 1129–1161.
- Fedotova, J, Kubatka, P, Büsselberg, D, Shleikin, A.G., Caprnda, M, Dragasek, J, et al. (2017). Therapeutical strategies for anxiety and anxiety-like disorders using plant-derived natural compounds and plant extracts. *Biomed Pharmacother.*, 95:437-446
- File, S.E., Lippa, A.S., Beer, B., Lippa, M.T. (2004). Animal tests of anxiety. *Curr Protoc Neurosci*, 26(1):1-22.
- Ganatra, T.H., Joshi, U.H., Bhalodia, P.N., Desai, T.R., Tirgar, P.R. (2012). A panoramic view on pharmacognostic, pharmacological, nutritional, therapeutic and prophylactic values of *Moringa oleifera* lam. *International Res Journal of Pharmacy*, 3(6):1-7.
- Hakim, Bey. (2010). *All things Moringa*. Diakses dari: <http://www.allthingsmoringa.com>.
- Hamid, H.A., Ramli, A.N., Yusoff, M.M. (2017). Indole Alkaloids from Plants as Potential Leads for Antidepressant Drugs: A Mini Review. *Frontiers Pharmacol.*, 8:1-7
- Handley, S.L., Mithani, S. (1984). Effects of  $\alpha$ -adrenoreceptor agonists and antagonists in a maze-exploration model of 'fear'-motivated behavior. *Naunyn-Schmeideberg's Archives of Pharmacology*, 327(1): 1–5.
- Hascoe't, M., Bourin, M. (1998). A new approach to the light/dark procedure in mice. *Pharmacol Biochem Behav*, 60(3):645 – 653.
- Ingale, S.P., Gandhi, F.P. (2016). Effect of aqueous extract of *moringa oleifera* leaves on pharmacological models of epilepsy and anxiety in mice. *International Journal of Epilepsy*, 3(1):12-19
- Jäger, A.K., Sabay, L. (2011). Flavonoids and the CNS. *Molecules*, 16(2):1471–1485.
- Karim, N., Khan, I., Khan, H., Ayub, B., Abdel-Halim, H., Gavande, N. (2018). Anxiolytic potential of natural flavonoids. *SM J Steroids Horm.*, 1(1):1001.

- Kessler, R.C., Ruscio, A.M., Shear, K., Wittchen, H.U. (2010). Epidemiology of anxiety disorders. *Curr Top Behav Neurosci*, 2: 21–35.
- Khawaja, T.M., Tahira, M., Ikram, U.H. (2010). *Moringa oleifera*: a natural gift – a review. *Journal of Pharmacy Science Res*, 2(11):775–781.
- Klingberg, G., Broberg, A.G. (2007). Dental fear/anxiety and dental behaviour management problems in children and adolescents: a review of prevalence and concomitant psychological factors. *Int J Paediatr Dent.*, 17(6):391-406.
- Komada, M., Takao, K., Miyakawa, T. (2008). Elevated plus maze for mice. *J Vis Exp*, 22:1088.
- Leo, L.M., Pamplona, F.A. (2014). Elevated plus maze test to assess anxiety-like behavior in the mouse. *Bio-protocol*, 4(16): 1211.
- Lister, R.G. (1987). The use of a plus-maze to measure anxiety in the mouse. *Psychopharmacology*, 92(2):180–185.
- Mishra, G. (2011). Traditional use, phytochemistry, and pharmacological properties of *Moringa oleifera* plants : An Overview. *Scholars Research library, Der pharmacia letter*, 3(2);141-164
- Nevid, J.S, Spencer, R.A, Greene, B. (2005). *Psikologi Abnormal Jilid 1*. (pp.163-165). Jakarta: Erlangga.
- Newton, T., Asimakopoulou. K., Daly, B., Scambler, S., Scott, S. (2012). The management of dental anxiety: time for a sense of proportion?. *Br Dent J.*, 213(6):271-274.
- Nuss, P. (2015). Anxiety disorders and GABA neurotransmission: a disturbance of modulation. *Neuropsychiatr Dis Treat*, 11:165–175.
- Orret, E.O., Hertz, M.B. (2012). Anxiety Control in the Dental Patient. *Dental Clinic N Am*, 56(1): 1–16.
- Onaivi, E.S., Martin, B.R. (1989). Neuropharmacological and physiological validation of a computer-controlled two compartment black and white box for the assessment of anxiety. *Prog. Neuro-psychopharmacol. Biol. Psychiatry*, 13(6): 963 – 976.
- Oosterink, F.M., de Jongh, A., Hoogstraten, J. (2009). Prevalence of Dental Fear and Phobia Relative to Other Fear and Phobia Subtypes. *European Journal of Oral Science*, 117(2):135–43.

- Paikra, B.K., Dhongade, H., Gidwani, B. (2017). Phytochemistry and pharmacology of *moringa oleifera* lam. *Journal of pharmacopuncture*, 20(3):194-200.
- Pellow, S., Chopin, P., File, S.E., Briley, M. (1985). Validation of open: closed arm entries in an elevated plus-maze as a measure of anxiety in the rat. *J Neurosci Methods*, 14(3): 149–167.
- Pusat Penelitian dan Pengembangan Peternakan. (2017). Penggunaan dan Penanganan Hewan Coba Rodensia dalam Penelitian. Diakses dari : <https://peternakan.litbang.pertanian.go.id/fullteks/booklet/juknis-rodensia-2107/isi-juknis-rodensia-2017.pdf>.
- Riss, J., Cloyd, J., Gates, J. (2008). Benzodiazepines in epilepsy: pharmacology and pharmacokinetics. *Acta Neurol Scand*, 118(2):69–86.
- Rupprecht, R., Eser, D., Zwanzger, P., Möller, H. (2006). GABAA receptors as targets for novel anxiolytic drugs. *World J Biol Psychiatry*, 7(4): 231–237.
- Setiawati A., Zunilda S.B., Suyatna F.D. (1995). Pengantar farmakologi. Dalam Ganiswara S.G. : *Farmakologi dan Terapi*. (pp. 1-14). Jakarta: Bagian Farmakologi Fakultas Kedokteran Universitas Indonesia.
- Sharma, K., Kumar, N., Raj, K., Niazi, J., Gupta, V. (2011). Anti-anxiety activity of *Eriobotrya japonica* leaf extracts. *Res Journal of Pharmacy Biology Chemistry Science*, 2:255-259.
- Singh, B.N., Singh, B.R., Singh, R.L., Prakash D., Dhakarey, R., Upadhyay, G., Singh, H.B. (2009). Oxidative DNA damage protective activity, antioxidant and anti-quorum sensing potentials of *Moringa oleifera*. *Food Chem Toxicol*, 47(6): 1109- 1116
- Tolistiawaty, I, Widjaja, J, Sumolang, P.P., Octaviani, F. (2014). Gambaran Kesehatan pada Mencit (*Mus musculus*) di Instalasi Hewan Coba. *Jurnal Vektor Penyakit*, 8(1): 27 – 32.
- Wei, X.Y., Yang, J.Y., Wang, J.H., Wu, C.F. (2007). Anxiolytic Effect of Saponins from *Panax Quinquefolium* in Mice. *J Ethnopharmacol*, 111(3):613–618.