

DAFTAR PUSTAKA

- Ahyong S.T., Lai J.C.Y., Sharkey D., Colgan D.J., & Ng P.K.L. (2007) Phylogenetics of the brachyuran crabs (Crustacea: Decapoda): the status of Podotremata based on small subunit nuclear ribosomal RNA. *Mol Phylogenet Evol.* 45:576–586.
- Arif, J., Nugroho, H., and Budiagung, S. (2015). Geoarkeologi Teras Purba Bengawan Solo di Sekitar Kabupaten Bojonegoro, Jawa Timur. *Purbawidya.* 4(2): 71-82
- Arifhidayat (talk) - Karya sendiri (Original text: I (Arifhidayat (talk)) created this work entirely by myself.), Domain Publik, <https://commons.wikimedia.org/w/index.php?curid=10513929>
- Ashkenazi, S., Motro, U., Goren-Inbar, N., Biton, R. & Rabinovich, R. (2005) New morphometric parameters for assessment of body size in the fossil freshwater crab assemblage from the Acheulian site of Gesher Benot Ya'aqov, Israel. *Journal of Archaeological Science.* 32:675-689.
- Barianto, D.H., Faza, A.F., Novian, M.I., & Husein, S. 2015. Stratigrafi Karbonat Formasi Seloredjo Anggota Dander di Sungai Banyurejo Kecamatan Bubulan Kabupaten Bojonegoro, Jawa Timur, Indonesia. *Proceeding, Seminar Nasional kebumihan ke- 8*, p. 249-254.
- Bird, M. I., Taylor, D., & Hunt, C. (2005) Palaeoenvironments of insular Southeast Asia during the Last Glacial Period: a savanna corridor in Sundaland? *Quaternary Science Reviews.* 24: 2228-2242.
- Bouteaux, A., Moigne, Anne-Marie., Semah, F., & Jacob, T. (2007). Les assemblages fauniques associées aux sites à Homo erectus du d'ome de Sangiran (Pléistocène moyen, Java, Indonésie). *C.R. Palevol.* 6: 169-179.
- Brosing A., Richter S., & Scholtz G. (2007) Phylogenetic analysis of the Brachyura (Crustacea, Decapoda) based on characters of the foregut with establishment of a new taxon. *J Zool Syst Evol Res.* 45: 20–32.
- Carpenter, K.E., & Niem, V. H. (1998) *The Living Marine Resources of The Western Central Pacific, Vol 2: Cephalopods, crustaceans, holothurians and sharks.* FAO: Roma.
- Collins, J.S.H., Lee, C., & Noad, J. (2003) Miocene and Pleistocene Crab (Crustacea, Decapoda) from Sabah and Sarawak. *Journal of Systematic Palaentology.* 1(3): 187-226.
- Corsini-Foka, M., Kondylatos, G., & Pancucci-Papadopoulou, M.A. (2013) A new alien crab for the Mediterranean Sea: *Xanthias lamarckii* (H. Milne Edwards, 1834) (Crustacea: Decapoda: Brachyura: Xanthidae). *Mediterranean Marine Science.* 14(2): 295-297.
- Davie, P. J. F. (1987) A New Species and New Records of *Portunus* (Decapoda: Portunidae) from Northern Australia. *Memoirs of the Queensland Museum.* 25(1): 227-231.
- Davie, P. J.F. (1993) A New Species of Sesarmine Crab (Brachyura: Grapsidae) form Japan and Taiwan, previously known as *Sesarma erythodactyla* Hess, 1865. *Crustacean Research.* 22: 65-74.
- Dennel, R., (2010) Out of Africa I: Current Problems and Future Prospects. In: J.G. Fleagle *et al.*, (eds.) *Vertebrate Paleobiology and*

- Palaeoanthropology*. New York, Springer Science and Business Media, pp. 247-273.
- Fae, M. (1996) Lithostratigraphy and fossil hominids of the Sangiran-Krikilan area, Java (Yogyakarta, Indonesia). *Memorie di Scienze Geologiche*, 48: 148-153.
- Fair use, <https://id.wikipedia.org/w/index.php?curid=2140650> diakses tanggal 5 September 2019
- Fenwick. (2019) *Pilumnus hirtellus*. http://www.aphotomarine.com/crab_hairy_pilumnus_hirtellus.html diakses tanggal 15 Agustus 2019
- Gilpin, D. (2006) *Lobsters, Crabs, and Other Crustaceans*. Minnesota, Capstone, p. 14
- Hutomo, M. & Moosa, M. K. (2005) A fossil freshwater crab from the Pliocene Tatrot Formation (Siwalik Group) in Northern India (Crustacea, Brachyura, Potamidae). *Indian Journal of Marine Sciences*, 34(1), 88-97
- ITIS. (2018) *Brachyura*. https://www.itis.gov/servlet/SingleRpt/SingleRpt?search_topic=TSN&search_value=98276#null (diakses tanggal 24 Juli 2018).
- Kato, H. & Koizumi, A. (2001) A new fossil geryonid crab (Crustacea: Decapoda: Brachyura) from the late Pliocene to early Pleistocene of Japan. *Proceedings of The Biological Society of Washington*, 114(1): 172-177.
- Klaus, S., Bohme, M., Schneider, S., Prieto, J., & Phetsomphou, B. (2011) Evidence of The Earliest Freshwater Decapod From Southeast Asia (Crustacea: Decapoda: Brachyura). *The Raffles Bulletin of Zoology*. 59(11): 47-51.
- Kumazawa, S. (1994) Quaternary Geology and Hydrogeology of the Madiun Basin, Indonesia. *Journal of Geosciences*. 37(8): 213-242.
- Lee, S.Y. (2015) Ecology of Brachyura. In: Castro, P., Davie, P.J.F., Guinot, D., Schram, F.R and von Faupel Klein, J.C. *The Crustacea vol. 9*. Boston, Brill, pp. 469-451.
- Lemaitre, R., Campos, N.H., Maestre, E.A.V., & Windsor, A.M. (2013) Discovery of an alien crab, *Scylla serrata* (Forsskal, 1755) (Crustacea: Decapoda: Portunidae), from the Caribbean coast of Colombia. *BioInvasion Records*. 2(4): 311-315.
- Moraes, S. A. S.N., Alencar, C. E. R. D., Thomsen, E., & Freire, F.A.M. (2015) New Record of the hairy crab *Pilumnus dasypodus* (Decapoda, Brachyura, Pilumnidae) in Northeastern Brazil. *The Journal of biodiversity data*. 11(2): 1-5.
- Morris, R.H., Abbott, D.P., & Haderlie, E.C. (1980) *Intertidal Invertebrates of California*. California, Stanford University Press, pp. 592-630.
- Ng, P.K.L. (1998) Crabs. In: Carpenter, K.E. & Niem, V.H. (eds.) *The Living Marine Resources of The Western Central Pacific Vol.2*. Rome, Food and Agriculture Organization of United Nations, pp. 1045-1083.
- Ng, P.K.L. (2004) Crustacea: Decapoda, Brachyura. In: Yule, C.M. and Yong, H.S. (eds.) *Freshwater Invertebrates of the Malaysian Region*. Kuala Lumpur, Academy of Sciences Malaysia, pp. 311-336.
- Ng, P.K.L., & Tan, L.W.H. (1984) The Indo-Pacific Pilumnidae I. Description of Four New Species of the Genus *Pilumnus* Leach, 1815, and Definition of a

- New Genus, *Bathypilumnus*. *Journal of The Singapore National Academy of Science*. 13: 1-7.
- Ng, P.K.L., & Cranbrook, E. (2014) Fossil Brachyuran Crabs from The Jambusan Caves (Bau, Sarawak), Collected by A.H. Everett in 1878-1879. *Scripta Geol.* 147: 289-308
- Pringgoprawiro, H. (1983) *Biostratigrafi dan Paleogeografi Cekungan Jawa Timur Utara, Suatu Pendekatan Baru*. Desertasi Doktor, Bandung: Institut Teknologi Bandung.
- Schultz, G. (2014) Evolution of Crabs – history and deconstruction of a prime example of convergence. *Zoology.*, 83(2): 87-105.
- Schweitzer, C.E., Feldmann, R.M., & Bonadio, C. (2009) A new family of brachyura (Crustacea: Decapoda: Goneplacoidea) from the eocene of Java, Indonesia. *Scripta Geol.* 138: 1-10.
- Semah, F. (1984) Palynology and Javanese Pithecanthropus palaeoenvironment. *Courier Forchungen Institut Senckenberg*. 69: 237-243.
- Semah, F., Saleki, H., & Falgueres, C. (2000). Did Early Man reach Java during the Late Pliocene? *Journal of Archaeological Science*. 27: 763-769.
- Semah, Anne-Marie, & Semah, F. (2012). The rain forest in Java through the Quaternary and its relationships with humans (adaptation, exploitation and impact on the forest). *Quaternary International*. 249:120-128
- Semah, Anne-Marie, Semah, F., Moigne, Anne-Marie, Ingicco, T., Purnomo, A., Simanjuntak, T., & Widiyanto, H. (2016) The palaeoenvironment context of the Palaeolithic of Java: A brief review. *Quaternary International*. 416: 38-45.
- Shih, His-Te, Komai, T., & Liu, Min-Yun. (2013) A new species of fiddler crab from the Ogasawara (Bonin) Islands, Japan, separated from the widely-distributed sister species *Uca (Paraleptuca) crassipes* (White, 1847) (Crustacea: Decapoda: Brachyura: Ocypodidae). *Zootaxa*. 3746 (1): 175-193.
- Simanjuntak, T. Prasetyo, B., & Handini, R. (2001). *Sangiran: Man, Culture, and Environment in Pleistocene Times*. Yayasan Obor Indonesia: Jakarta
- Susilohadi, Late tertiary and quaternary geology of the East Java Basin, Indonesia, Doctor of Philosophy thesis, School of Geosciences, University of Wollongong, 1995. <http://ro.uow.edu.au/theses/1973>
- Tsang, L.M., Schubart, C.D., Ahyong, S.T., Lai, J.C.Y., Au, E.Y.C., Chan, Tim-Yan, Ng, P.K.L., & Chu, K.H. (2014) Evolutionary history of true crabs (Crustacea: Decapoda: Brachyura) and the origin of freshwater crabs. *Mol. Biol. Evol.*, 31(5): 1173-1187.
- Valinassab, T., Fatemi, S.M.R.F., & Ghotbeddin, N. 2012. First Record of *Galene bispinosa* Herbst, 1783 [Decapoda; Brachyura; Galenidae] from the Gulf of Oman, Iran. *World Applied Science Journal* 17(4): 439-441.
- Voris, H.K. (2000) Maps of Pleistocene sea levels in Southeast Asia: shorelines, river systems and time durations. *Jurnal of Biogeography* 27: 1153-1167.
- Watanabe, N., & Kadar, D. (1985) *Quaternary Geology of The Hominid Fossil Bearing Formations in Java*. Jakarta: Geological Research and Development Centre, pp. 11-44, 63-68.
- Whitten, T., Soeriaatmadja R., & Afiff S.A. (1997) *The Ecology of Java and Bali* (Ecology of Indonesia Series, Volume 2). Portland, Timber Press.



- Yudha, D.A., Ramadhani, R., Suriyanto, R.A., & Novian, M.I. (2002) The diversity of sharks fossils in Plio-Pleistocene of Java, Indonesia. *AIP Conference Proceedings*. doi: 10.1063/1.5050109
- Yudha, D.S., & Suriyanto, R.A. (2016) Marine-mollusc fossils (Mollusca: Bivalvia and Gastropoda) and their paleoenvironmental interpretation from the Situs manusia purba Sangiran area. *AIP Conference Proceedings*. 1744.
- Yudha, D. S., & Rahmayani, D.N.A. (2017) Keanekaragaman Fosil Kerang (Bivalvia dan Gastropoda) di Situs Banjarejo, Kecamatan Gabus, Kabupaten Grobogan. *Jurnal Sangiran*. 6: 40-51