

DAFTAR PUSTAKA

- Adab, P., Pallan, M., & Whincup, P. H. (2018). Is BMI the best measure of obesity? *British Medical Journal*, 360(k1274), 1–2.
- Aguirre, T., Koehler, A., & Tovar, A. (2015). Relationships among hypertension, waist circumference, and body composition in a rural Mexican-American population. *Journal of Family Medicine & Community Health*, 2(7), 1057–1061.
- Ahmad, N., Adam, S. I. M., Nawi, A. M., Hassan, M. R., & Ghazi, H. F. (2016). Abdominal obesity indicators: waist circumference or waist-to-hip ratio in Malaysian adults population. *International Journal of Preventive Medicine*, 7, 82–86.
- Batsis, J. A., Mackenzie, T. A., Bartels, S. J., Sahakyan, K. R., Somers, V. K., & Jimenez, F. L.-. (2016). Diagnostic accuracy of body mass index to identify obesity in older adults: NHANES 1999–2004. *International Journal of Obesity*, 40(5), 761–767.
- Beg, M., Sharma, V., Akhtar, N., Gupta, A., & Mohd, J. (2011). Role of Antioxidants in Hypertension. *Indian Academy of Clinical Medicine Journal*, 12(2), 122-7.
- Bouchi, R., Asakawa, M., Ohara, N., Nakano, Y., Takwuchi, T., Murakami, M., Sasahara, Y., Numasawa, M., Minami, I., Izumiyama, H., Hasimoto, K., Yoshimoto, T., & Ogawa, Y. (2016). Indirect measure of visceral adiposity ‘A Body Shape Index’ (ABSI) is associated with arterial stiffness in patients with type 2 diabetes. *BMJ Open Diabetes Research and Care*, 4, 1-7.
- Browning, L. M., Hsieh, S. D., & Ashwell, M. (2010). A systematic review of waist-to height ratio as a screening tool for the prediction of cardiovascular disease and diabetes: 0.5 could be a suitable global boundary value. *Nutrition Research Reviews*, 23, 247–269.
- Buford, T. W. (2016). Hypertension and Aging. *Ageing Research Reviews*, 2, 96–111.
- Caminha, T. C. S., Ferreira, H. S., Costa, N. S., Nakano, R. P., Carvalho, R. E. S., Xavier, A. F. S., & Asuncao, M. L. (2017). Waist-to-height ratio is the best anthropometric predictor of hypertension: A population-based study with women from a state of northeast of Brazil. *Medicine*, 96(2), 1-8.
- CDC. (2015). *Body mass index: considerations for practitioners 2015*.
- Cheung, Y. B. (2014). “A body shape index” in middle-age and older Indonesian population: scaling exponents and association with incident hypertension. *PLoS ONE*, 9(1), 2–6.

- Chobanian, A. V., Backris, G., Black, H., Cushman, W., Green, L., & Izzo, J. (2003). The seventh report of the joint national committee on prevention, detection, evaluation, and treatment of high blood pressure. *The Journal of the American Medical Association*, 289, 1–86.
- Choi, J. R., Koh, S. B., & Choi, E. (2018). Waist-to-height ratio index for predicting incidences of hypertension: the ARIRANG study. *BMC Public Health*, 18, 767 - 773.
- Chung, W., Park, C. G., & Ryu, O. (2016). Association of a new measure of obesity with hypertension and health-related quality of life. *PLOS One*, 11(5), 1-10.
- Dahlan, S. (2010). *Penelitian diagnostik: dasar-dasar teoretis dan aplikasi dengan program spss dan stata*. Jakarta: Epidemiologi Indonesia.
- Dahlan, S. (2016). *Statistik untuk kedokteran dan kesehatan: deskriptif, bivariat, dan multivariat*. Jakarta: Epidemiologi Indonesia.
- Donmezi, S., Gokalp, O., Dogan, M., Vural, H., & Yigit, B. (2014). Effects of short-term hyperglycemia on the vasoconstriction of the aorta. *Turkish Journal of Medical Sciences*, 44, 941–945.
- Dua, S., Bhuker, M., Sharma, P., Dhall, M., & Kapoor, S. (2014). Body mass index relates to blood pressure among adults. *North American Journal of Medical Sciences*, 6(2), 89–95.
- Faulkner, J. L. & Belin de Chantemele, E. J. (2018). Sex differences in mechanisms of hypertension associated with obesity. *Hypertension*, 71(1), 15–21.
- Feng, R.-N., Zhao, C., Wang, C., Niu, Y.-C., Li, K., Guo, F.-C., Li, S.-T., Sun, C.-H., & Li, Y. (2012). BMI is strongly associated with hypertension, and waist circumference is strongly associated with type 2 diabetes and dyslipidemia, in Northern Chinese adults. *Journal of Epidemiology*, 22(4), 317–323.
- Frayon, S., Cavaloc, Y., Wattelez, G., Cherrier, S., Lerrant, Y., Ashwell, M., & Galy, O. (2018). Potential for waist-to-height ratio to detect overfat adolescents from a Pacific Island, even those within the normal bmi range. *Obesity Research and Clinical Practice*, 12(4), 351–357.
- Halperin, R., Sesso, H., Ma, J., Buring, J., Stampfer, M., & Gaziano, J. (2006). Dyslipidemia and the risk of incident hypertension in men. *Hypertension*, 47, 45–50.
- Hastuti, J., Kagawa, M., Byrne, N. M., & Hills, A. P. (2017). Determination of new anthropometric cut-off values for obesity screening in Indonesian adults. *Asia Pacific Journal of Clinical Nutrition*, 26(4), 650–656.

- He, S., & Chen, X. (2013). Could the new body shape index predict the new onset of diabetes mellitus in the Chinese population? *PLoS ONE*, 8(1), 8–11.
- Jackson, C., & Wenger, N. (2011). Cardiovascular disease in the elderly. *Revista Española de Cardiología*, 64(8), 697–712.
- Ji, M., Zhang, S., & An, R. (2018). Effectiveness of A Body Shape Index (ABSI) in predicting chronic diseases and mortality: a systematic review and meta-analysis. *Obesity Reviews*, 19:737–59
- Jiang, S., Lu, W., Zong, X., Ruan, H., & Liu, Y. (2016). Obesity and hypertension (review). *Experimental and Therapeutic Medicine*, 12, 2395–2399.
- Kalani, Z., Salimi, T., & Rafiei, M. (2015). Comparison of obesity indexes BMI, WHR and WC in association with hypertension: results from a blood pressure status survey in Iran. *Journal of Cardiovascular Disease Research*, 6(2), 72–77.
- Kaplan, N. (2006). Primary hypertension: pathogenesis. In: Kaplan, N., editor. *Kaplan's Clinical Hypertension*. 9. Philadelphia: Lippincott Williams & Wilkins.
- Kementrian Kesehatan RI. (2018). *Riset kesehatan dasar (RISKESDAS) 2018*. Jakarta: Badan Penelitian dan Pengembangan Kesehatan.
- Kim, D., Kim, C., Ding, E., Townsend, M., & Lipsitz, L. (2013). Adiponectin levels and the risk of hypertension: a systematic review and meta-analysis. *Hypertension*, 62(1), 27–32.
- Kokkinos, P., Giannelou, A., Athanasiosmanolis, & Pittaras, A. (2009). Physical activity in the prevention and management of high blood pressure. *Hellenic Journal of Cardiology*, 50, 52–59.
- Kotsis, V., Stabouli, S., Papakatsika, S., Rizos, Z., & Parati, G. (2010). Mechanisms of obesity-induced hypertension. *Hypertension Research*, 33(5), 386–393.
- Krakauer, N. Y., & Krakauer, J. C. (2012). A new body shape index predicts mortality hazard independently of body mass index. *PLoS ONE*, 7(7), 1–10.
- Leon, B. G. C., Jensen, M. D., Hartman, J. J., & Jensen, T. B. (2016). Self-measured vs professionally measured waist circumference. *Annals of Family Medicine*, 14(3), 262–266.
- Li, W., Chen, I., Chang, Y., Loke, S., Wang, S., & Hsiao, K., (2013). Waist-to-height ratio, waist circumference, and body mass index as indices of cardiometabolic risk among 36,642 Taiwanese adults. *European Journal Nutrition*, 52, 57–65

- Luz, R.H., Barbosa, A.R., & d'Orsi, E. (2016). Waist circumference, body mass index and waist-height ratio: are two indices better than one for identifying hypertension risk in older adults? *Preventive Medicine*, 93, 76-81.
- Ma, W. Y., Yang, C. Y., Shih, S. R., Hsieh, H. J., Hung, C. S., Chiu, F. C., Lin, M. S., Liu, P. H., Hua, C. H., Hsein, Y. C., Chuang, L. M., Lin, J. W., Wei, J. N., & Li, H. Y. (2013). Measurement of waist circumference: midabdominal or iliac crest? *Diabetes Care*, 36(6), 1660–1666.
- Mauvais-Jarvis, F. (2015). Sex differences in metabolic homeostasis, diabetes, and obesity. *Biology of Sex Differences*, 6(14): 1-9.
- Menekşe, D., & Balci, S. (2017). Effect of body mass index and waist-hip ratio on blood pressure in adolescents: a descriptive and correlational study. *Turkiye Klinikleri Journal of Nursing*, 9(2), 99–106.
- Muñoz-Durango, N., Fuentes, C. A., Castillo, A. E., González-Gómez, L. M., Vecchiola, A., Fardella, C. E., & Kalergis, A. M. (2016). Role of the renin-angiotensin-aldosterone system beyond blood pressure regulation: molecular and cellular mechanisms involved in end-organ damage during arterial hypertension. *International Journal of Molecular Sciences*, 17(7), 1–17.
- NHANES. (2007). *Anthropometry Procedure Manual*. Washington: US Department of Health and Human Services.
- Ononamadu, C. J., Ezekwesili, C. N., Onyeukwu, O. F., Umeuguaju, U. F., Ezeigwe, O. C., & Ihegboro, G. O. (2017). Comparative analysis of anthropometric indices of obesity as correlates and potential predictors of risk for hypertension and prehypertension in a population in Nigeria. *Cardiovascular Journal of Africa*, 28(2): 92-99.
- Oparil, S., Zaman, M., & Calhoun, D. (2003). Pathogenesis of hypertension. *Annals of Internal Medicine*, 139, 761–776.
- Oparil, S., & Miller, A. (2005). Gender and blood pressure. *The Journal of Clinical Hypertension*, 7(5), 300–309.
- Pertiwi, R. (2016). *Nilai sensitivitas dan spesifisitas titik potong rasio lingkar pinggang tinggi badan sebagai prediktor kejadian hipertensi pada orang dewasa di Indonesia*. UIN Syarif Hidayatullah Jakarta.
- Power, M. L., & Schulkin, J. (2008). Sex differences in fat storage, fat metabolism, and the health risks from obesity: possible evolutionary origins. *British Journal of Nutrition*, 99, 931–940.
- Rahajeng, E., & Tuminah, S. (2009). Prevalensi hipertensi dan determinannya di Indonesia. *Majalah Kedokteran Indonesia*, 580–587.

- Rezende, A. C., Souza, L. G., Jardim, T. V., Perillo, N. B., Araújo, Y. C. L., Garcia de Souza, S., Sousa, A. L. L., Moreira, H. G., Barroso de Souza, W. K. S., Peixoto, M. G., & Jardim, P. C. B. V. (2018). Is waist-to-height ratio the best predictive indicator of hypertension incidence? A cohort study. *Biomedical Central Public Health, 18*(1), 281–291.
- Sastroasmoro, S. (2014). *Dasar-dasar metodologi penelitian klinis*. Jakarta: Sagung Seto.
- Strauss, J., Witoelar, F., & Sikoki, B. (2016). *The fifth wave of the indonesia family life survey: overview and field report: volume 1*. California: RAND Labor and Population Working Paper Series
- Talukder, M. A. H., Johnson, W. M., Varadharaj, S., Lian, J., Kearns, P. N., El-Mahdy, M. A., Liu, X., & Zweier, J. L. (2011). Chronic cigarette smoking-causes hypertension, increased oxidative stress, impaired no bioavailability, endothelial dysfunction, and cardiac remodeling in mice. *American Journal of Physiology-Heart and Circulatory Physiology, 300*(1), H388–H396.
- Tawfik, H. M. (2018). Waist height ratio and waist circumference in relation to hypertension, framingham risk score in hospitalized elderly Egyptians. *Egyptian Heart Journal, 70*(3), 213–216.
- Thuy, A., Blizzard, L., Schmidt, M., Luc, P., Granger, R., & Dwyer, T. (2010). The association between smoking and hypertension in a population-based sample of Vietnamese men. *Journal of Hypertension, 28*(2), 245–250.
- Tomicki, C., Gerage, A. M., Ritti-Dias, R. M., Silva, D. A. S., & Benedetti, T. R. B. (2017). Diagnostic property of anthropometric indicators in the prediction of high body fat estimated by DXA in hypertensive women. *Revista Brasileira de Cineantropometria e Desempenho Humano, 17*(3), 299–308.
- Tuan, N. T., Adair, L. S., Suchindran, C. M., He, K., & Popkin, B. M. (2009). The association between body mass index and hypertension is different between East and Southeast Asians. *American Journal of Clinical Nutrition, 89*, 1905 - 12.
- Tuan, N. T., Adair, L. S., Stevens, J., & Popkin, B. M. (2010). Prediction of hypertension by different anthropometric indices in adults: the change in estimate approach. *Public Health Nutrition, 13*(5), 639 - 646.
- Wenger, N. K., Arnold, A., Merz, N. B., Cooper-DeHoff, C., Ferdinand, K. C., Fleg, J. L., Gulati, M., Isadinso, I., Itchhaporia, D., Light-McGroary, K., Lindley, K. J., Mieres, J. H., Rosser, M. L., Saade, G. R., Walsh, M. N., Pepine, C. J. Hypertention accros a woman's cycle. *Journal of the American College of Cardiology, 71*(16), 1797-1813.

WHO. (2018). Obesity and overweight. Retrieved December 20, 2018, from <https://www.who.int/news-room/fact-sheets/detail/obesity-and-overweight>

Wirawan, N. N. (2016). Sensitifitas dan spesifisitas IMT dan lingkar pinggang-panggul dalam mengklasifikasikan kegemukan pada wanita. *Indonesian Journal of Human Nutrition*, 3(1), 49–59.

WHO. (2004). World health organization (WHO) expert consultation: appropriate body-mass index for asian populations and its implications for policy and intervention strategies. *Lancet*, 363(9403), 157–163.

WHO. (2011). *Waist circumference and waist- hip ratio WHO expert consultation*. Swiss: WHO Press.

Xu, R. Y., Zhou, Y. Q., Zhang, X. M., Wan, Y. P., & Gao, X. (2018). Body mass index, waist circumference, body fat mass, and risk of developing hypertension in normal-weight children and adolescents. *Nutrition, Metabolism, and Cardiovascular Diseases*, 28(10), 1061–1066.

Yoo, E. (2016). Waist-to-height ratio as a screening tool for obesity and cardiometabolic risk. *Korean Journal Pediatric*, 59(11), 425–431.