



## DAFTAR PUSTAKA

- Al-Kawaz, S. A. ,2014, 'Factors that predict mortality rate in biliary atresia', *J Fac Med Baghdad*. Baghdad: J Fac Med Baghdad, 56, pp. 136–140.
- Aterials, M. and Ethods, M. ,2000, 'Initial Response to Portoenterostomy Determines Long-Term Outcome in Patients With Biliary Atresia', *J Pediatr Surg*, 35, pp. 593–597.
- Bittmann, S. ,2005, 'Surgical experience in children with biliary atresia treated with portoenterostomy', *Curr Surg.*, 62(4), pp. 439–443.
- Bjørnland, K., Hinna, M., Aksnes, G., Stensrud, K. J., Ertresvåg, K., Bjørnstad-Østensen, A., *et al.* ,2018, 'Outcome for biliary atresia patients treated at a low-volume centre', *Scand J Gastroenterol*. Informa UK Limited, trading as Taylor & Francis Group, 53(4), pp. 471–474.
- Broue, P., Chardot, C., Gauthier, F., Lachaux, A., Sarles, J., Serinet, M.-O., *et al.* ,2009, 'Impact of Age at Kasai Operation on Its Results in Late Childhood and Adolescence: A Rational Basis for Biliary Atresia Screening', *Pediatrics*, 123(5), pp. 1280–1286.
- Carvalho, E. de, Ivantes, C. A. P. and Bezerra, J. A. ,2007, 'Extrahepatic biliary atresia: current concepts and future directions', *J. Pediatr. (Rio. J.)*, 83(2), pp. 105–120.
- Chardot, C. and Serinet, M. O. ,2006, 'Prognosis of biliary atresia: What can be further improved?', *J Pediatr*, 148(4), pp. 432–435.
- D'Agata, I. D. and Balistreri, W. F. ,2018, 'Evaluation of Liver Disease in the Pediatric Patient', *Pediatr Rev*, 20(11), pp. 376–390.
- Davenport, M. ,2006, 'Biliary atresia: Outcome and management', *Indian J. Pediatr.*, 73(9), pp. 825–828.
- Davenport, M., Ong, E., Sharif, K., Alizai, N., McClean, P., Hadzic, N., *et al.* ,2011, 'Biliary atresia in England and Wales: Results of centralization and new benchmark', *J. Pediatr. Surg.*, 46(9), pp. 1689–1694.
- Desmet, V. J. and Roskams, T. A. D. ,2007, 'The Cholangiopathies', in Suchy, F. J., Sokol, R. J., and Balisteri, W. F. (eds) *Liver Dis. Child*. Third. New York: Cambridge University Press, pp. 35–59.
- Ferreira, A. R., Queiroz, T. C. N., Vidigal, P. V. T., Ferreira, R. di P., Wanderley, D. C. and Fagundes, E. D. T. ,2019, 'Multivariate Analysis of Biliary Flow-Related Factors and Post-Kasai Survival in Biliary Atresia Patients', *Arg Gastroenterol*, 56(1), pp. 71–78.
- Goda, T., Kawahara, H., Kubota, A., Hirano, K., Umeda, S., Tani, G., *et al.* ,2013, 'The most reliable early predictors of outcome in patients with biliary atresia after Kasai's operation', *J. Pediatr. Surg*. Elsevier Inc., 48(12), pp. 2373–2377.



- Goedeke, J., Scholz, S., The, P. and Jagged, T. ,2014, 'Biliary Atresia', *SAGE*, (10), pp. 1–15.
- Gunadi, Gunawan, T. A., Widiyanto, G., Yuanita, A., Mulyani, N. S. and Makhmudi, A. ,2018, 'Liver transplant score for prediction of biliary atresia patients' survival following Kasai procedure', *BMC Res Notes*. BioMed Central, 11(1), pp. 1–5.
- Hinds, R., Davenport, M., Mieli-Vergani, G. and Hadžić, N. ,2004, 'Antenatal presentation of biliary atresia', *J. Pediatr.*, 144(1), pp. 43–46.
- Hundt, M., Basit, H. and John, S. ,2019, *Physiology, Bile Secretion, StatPearls Publ. LLC*. (Accessed: 24 July 2019).
- Karangwa, O. ,2016, *A Retrospective Review of the Outcome of Children Presenting to Tygerberg Children 's Hospital with Biliary Atresia* By. the University of Stellenbosch.
- Khanna, K., Bhatnagar, V., Agarwala, S., Srinivas, M. and Gupta, S. D. ,2018, 'Ratio of Preoperative and Postoperative Serum Bilirubin Levels Predicts Early Outcome Following Biliary Atresia Surgery', *J Indian Assoc Pediatr Surg*, 23, pp. 10–15.
- Lai, H., Chen, W., Chen, C., Hung, W. and Chang, M. ,2006, 'Long-Term Prognosis and Factors Affecting Biliary Atresia from Experience over A 25 Year Period', *Chang Gung Med J*, 29(7), pp. 234–239.
- Lampela, H. ,2013, *BILIARY ATRESIA Treatment Results and Native Liver Function*. Children's Hospital Institute of Clinical Medicine, University of Helsinki and National Graduate School of Clinical Investigation Helsinki.
- Laohapensang, M. and Srikuanchaen, P. ,2017, 'The Relationship between Clinical Outcomes After Kasai Operation and Related Factors in Infants with Biliary Atresia', *J Med Assoc Thai*, 100(4), pp. 99–104.
- Levin, L. N., Goddard, E., De Lacy, R. and Pillay, K. ,2012, *Biliary Atresia at Red Cross War Memorial Children's Hospital: A retrospective descriptive study reviewing the age of presentation, clinical course and outcome of infants presenting to RCWMCH with biliary atresia*, Uct. University of Cape Town.
- Mack, C. L. and Sokol, R. J. ,2005, 'Unraveling the Pathogenesis and Etiology of Biliary Atresia', *Pediatr Res*, 57(5), pp. 87–94.
- Maheranny, M. ,2016, 'Factors Affecting the Success Rate of Kasai Portoenterostomy', *New Ropanasuri J Surg*, 1(1), pp. 1–7.
- Namasemayam, D. and Nallusamy, M. ,2017, 'Factors influencing outcome after hepatic portoenterostomy among extrahepatic bile duct atresia patients in hospital Sultanah Bahiyah, Alor Setar', *Med J Malaysia*, 72(6), pp. 329–332.
- Ohhama, Y., Shinkai, M., Fujita, S., Nishi, T. and Yamamoto, H. ,2000, 'Early



- prediction of long-term survival and the timing of liver transplantation after the Kasai operation', *J Pediatr Surg*, 35(7), pp. 1031–1034.
- Qisthi, S. A., Gunadi, Makhmudi, A. and Dwihantoro, A. ,2019, *Faktor prognosis bilirubin total dan albumin 3 bulan pasca operasi Kasai terhadap kesintasan pasien atresia bilier di RSUP DR. Sardjito*. Universitas Gadjah Mada.
- Sanghai, S. R., Shah, I., Bhatnagar, S. and Murthy, A. ,2009, 'Incidence and prognostic factors associated with biliary atresia in western India.', *Ann hepatol*, 8(2), pp. 120–2.
- Saragih, D. S. P., Gunadi and Makhmudi, A. ,2019, *Faktor prognosis bilirubin total dan albumin 1 bulan pasca operasi Kasai terhadap kesintasan pasien atresia bilier di RSUP Dr. Sardjito*. Universitas Gadjah Mada.
- Schoen, B. T., Lee, H., Sullivan, K. and Ricketts, R. R. ,2001, 'The Kasai portoenterostomy: When is it too late?', *J Pediatr Surg*, 36(1), pp. 97–99.
- Shanmugam, N. P., Narasimhan, G. and Rajindrajith, S. ,2016, 'Living with biliary atresia : From Kasai portoenterostomy to liver transplantation', *Sri Lanka J. Child Heal.*, 45(2), pp. 116–122.
- Shneider, B. L., Brown, M. B., Haber, B., Whittington, P. F., Schwarz, K., Squires, R., *et al.* ,2006, 'A multicenter study of the outcome of biliary atresia in the United States, 1997 to 2000', *J Pediatr*, 148(4).
- Sookpotarom, P., Vejchapipat, P., Chittmitrapap, S., Chongsrisawat, V., Chandrakamol, B. and Poovorawan, Y. ,2006, 'Short-term results of Kasai operation for biliary atresia: Experience from one institution', *Asian J Surg*, 29(3), pp. 188–192.
- Superina, R., Magee, J. C., Brandt, M. L., Healey, P. J., Tiao, G., Ryckman, F., *et al.* ,2012, 'The anatomic pattern of biliary atresia identified at time of Kasai hepatportoenterostomy and early postoperative clearance of jaundice are significant predictors of transplant-free survival', *Ann Surg*, 254(4), pp. 577–585.
- Wang, Z., Chen, Y., Peng, C., Pang, W., Zhang, T., Wu, D., *et al.* ,2018, 'Five-Year Native Liver Survival Analysis in Biliary Atresia from a Single Large Chinese Center: the Death/Liver Transplantation Hazard Change and the Importance of Rapid Early Clearance of Jaundice', *J Pediatr Surg*. Elsevier Inc, 3468(18)30, pp. 1–5.