

## DAFTAR PUSTAKA

- Baig, M., Klein, J. and Mechtler, L. (2016). Imaging of brain tumors. *Continuum: Lifelong Learning in Neurology*, 22(5), pp.1529-1552.
- Bulakbasi, N., Guvenc, I., Onguru, O., Erdogan, E., Tayfun, C. and Ucoz, T. (2004). The added value of the apparent diffusion coefficient calculation to magnetic resonance imaging in the differentiation and grading of malignant brain tumors. *Journal of Computer Assisted Tomography*, 28(6), pp.735-746.
- Cha, S. (2006) 'Update on brain tumor imaging: From anatomy to physiology', *American Journal of Neuroradiology*, 27(3), pp. 475–487.
- Engelhorn, T., Savaskan, N.E., Schwarz, M.A., Kreutzer, J., Meyer, E.P., Hahnen, E., Ganslandt, O., Dörfler, A., Nimsky, C., Buchfelder, M. and Eyüpoglu, I.Y., 2009. Cellular characterization of the peritumoral edema zone in malignant brain tumors. *Cancer Science*, 100(10), pp.1856-1862.
- Ferlay, J., Colombet, M., Soerjomataram, I., Mathers, C., Parkin, D.M., Piñeros, M., Znaor, A. and Bray, F., 2019. Estimating the global cancer incidence and mortality in 2018: GLOBOCAN sources and methods. *International Journal of Cancer*, 144(8), pp.1941-1953.
- Fan, G.G., Deng, Q.L., Wu, Z.H. and Guo, Q.Y., 2006. Usefulness of diffusion/perfusion-weighted MRI in patients with non-enhancing supratentorial brain gliomas: a valuable tool to predict tumour grading?. *The British journal of radiology*, 79(944), pp.652-658.
- Guzman, R., Altrichter, S., El-Koussy, M., Gralla, J., Weis, J., Barth, A., Seiler, R.W., Schroth, G. and Lövblad, K.O., 2008. Contribution of the apparent diffusion coefficient in perilesional edema for the assessment of brain tumors. *Journal of neuroradiology*, 35(4), pp.224-229.
- Hartmann, M., Jansen, O., Heiland, S., Sommer, C., Münkler, K. and Sartor, K., 2001. Restricted diffusion within ring enhancement is not pathognomonic for brain abscess. *American Journal of Neuroradiology*, 22(9), pp.1738-1742.
- Kang, Y., Choi, S.H., Kim, Y.J., Kim, K.G., Sohn, C.H., Kim, J.H., Yun, T.J. and Chang, K.H., 2011. Gliomas: histogram analysis of apparent diffusion coefficient maps with standard-or high-b-value diffusion-weighted MR imaging—correlation with tumor grade. *Radiology*, 261(3), pp.882-890.
- Kesari, S. (2011) 'Tumor Biology: The Potential to Improve Current Diagnosis and Treatments', *YSONC*. Elsevier Inc., 38(6), pp. S2–S10. doi: 10.1053/j.seminoncol.2011.09.005.

Kim, B. and Gutierrez, J. (2012). Contrast-enhanced MR imaging in neuroimaging. *Magnetic Resonance Imaging Clinics of North America*, 20(4), pp.649-685

- Kimura, M. and da Cruz, L.C.H., 2016. Multiparametric MR imaging in the assessment of brain tumors. *Magnetic Resonance Imaging Clinics*, 24(1), pp.87-122.
- Kono, K., Inoue, Y., Nakayama, K., Shakudo, M., Morino, M., Ohata, K., *et al.*, (2001). The role of diffusion-weighted imaging in patients with brain tumors. *American Journal of Neuroradiology*, 22(6): 1081–1088.
- Lequin, M. and Hendrikse, J. (2017). Advanced MR imaging in pediatric brain tumors, Clinical Applications. *Neuroimaging Clinics of North America*, 27(1), pp.167-190.
- Louis, D.N., Perry, A., Reifenberger, G., Von Deimling, A., Figarella-Branger, D., Cavenee, W.K., Ohgaki, H., Wiestler, O.D., Kleihues, P. and Ellison, D.W., 2016. The 2016 World Health Organization classification of tumors of the central nervous system: a summary. *Acta neuropathologica*, 131(6), pp.803-820
- McKinney, P.A., 2004. Brain tumours: incidence, survival, and aetiology. *Journal of Neurology, Neurosurgery & Psychiatry*, 75(suppl 2), pp.ii12-ii17.
- Nabors, L.B., 2016. Central nervous system: notable developments in the management of primary and recurrent gliomas. *Journal of the National Comprehensive Cancer Network*, 14(5S), pp.681-684.
- Ono, Y., Chernov, M.F., Muragaki, Y., Maruyama, T., Abe, K. and Iseki, H., 2018. Imaging of intracranial gliomas. In *Intracranial Gliomas Part I-Surgery* (Vol. 30, pp. 12-62). Karger Publishers.
- Ostrom, Q.T., Gittleman, H., Truitt, G., Boscia, A., Kruchko, C. and Barnholtz-Sloan, J.S., 2018. CBTRUS statistical report: primary brain and other central nervous system tumors diagnosed in the United States in 2011–2015. *Neuro-oncology*, 20(suppl\_4), pp.iv1-iv86.
- Wen, P. and Kesari, S. (2008). Malignant Gliomas in Adults. *New England Journal of Medicine*, 359(5), pp.492-507.
- Raisi-nafchi, M., Faeghi, F., Zali, A., Haghghatkah, H. and Jalal-Shokouhi, J. (2016). Preoperative grading of astrocytic supratentorial brain tumors with diffusion-weighted magnetic resonance imaging and apparent diffusion coefficient. *Iranian Journal of Radiology*, 13(3):e30426.
- Rees, J.H., Smirniotopoulos, J.G., Jones, R.V. and Wong, K., 1996. Glioblastoma multiforme: radiologic-pathologic correlation. *Radiographics*, 16(6), pp.1413-1438.
- Schaefer, P. W., Grant, P. E. and Gonzalez, R. G. (2000). Diffusion-weighted MR Imaging of the Brain. *Radiology*, 217(2):331–345.
- Schoenegger, K., Oberndorfer, S., Wuschitz, B., Struhal, W., Hainfellner, J., Prayer, D., Heinzl, H., Lahrmann, H., Marosi, C. and Grisold, W., 2009.

- Peritumoral edema on MRI at initial diagnosis: an independent prognostic factor for glioblastoma?. *European journal of neurology*, 16(7), pp.874-878.
- Shakal, A.A.S. and Mokbel, E.A.H., 2014. Hemorrhage after stereotactic biopsy from intra-axial brain lesions: incidence and avoidance. *Journal of Neurological Surgery Part A: Central European Neurosurgery*, 75(03), pp.177-182.
- Stoyanov, G.S. and Dzhenkov, D.L., 2018. On the concepts and history of glioblastoma multiforme-morphology, genetics and epigenetics. *Folia medica*, 60(1), pp.48-66.
- Svolos, P., Tsolaki, E., Kapsalaki, E., Theodorou, K., Fountas, K., Fezoulidis, I. and Tsougos, I., 2013. Investigating brain tumor differentiation with diffusion and perfusion metrics at 3T MRI using pattern recognition techniques. *Magnetic Resonance Imaging*, 31(9), pp.1567-1577.
- Wang, S., Meng, M., Zhang, X., Wu, C., Wang, R., Wu, J., Sami, M.U. and Xu, K., 2018. Texture analysis of diffusion weighted imaging for the evaluation of glioma heterogeneity based on different regions of interest. *Oncology letters*, 15(5), pp.7297-7304.
- Wesseling, P., Kros, J.M. and Jeuken, J.W., 2011. The pathological diagnosis of diffuse gliomas: towards a smart synthesis of microscopic and molecular information in a multidisciplinary context. *Diagnostic Histopathology*, 17(11), pp.486-494.
- Wu, C.X., Lin, G.S., Lin, Z.X., Zhang, J.D., Liu, S.Y. and Zhou, C.F., 2015. Peritumoral edema shown by MRI predicts poor clinical outcome in glioblastoma. *World Journal of Surgical Oncology*, 13(1), p.97.
- Zhang, L., Min, Z., Tang, M., Chen, S., Lei, X. and Zhang, X., 2017. The utility of diffusion MRI with quantitative ADC measurements for differentiating high-grade from low-grade cerebral gliomas: evidence from a meta-analysis. *Journal of the Neurological Sciences*, 373, pp.9-15.