

REFERENCES

- Abd Ghani, A. (2017). *Encapsulation of Functional Oil by Spray Drying and their Stability in Spray-Dried Powder*. EHIME UNIVERSITY.
- Acosta-Esquivarosa, J., Jáuregui-Haza, U., Amaro-González, D., & Sordo-Martínez, L. (2009). Spray drying of aqueous extract of *Mangifera indica* L (Vimang): scale up for the process. *World Applied Sciences Journal*, 6(3), 408–412.
- Aghbashlo, M., Mobli, H., Madadlou, A., & Rafiee, S. (2013). Fish oil microencapsulation as influenced by spray dryer operational variables. *International Journal of Food Science and Technology*, 48(8), 1707–1713. <https://doi.org/10.1111/ijfs.12141>
- Ahn, J. H., Kim, Y. P., Lee, Y. M., Seo, E. M., Lee, K. W., & Kim, H. S. (2008). Optimization of microencapsulation of seed oil by response surface methodology. *Food Chemistry*, 107(1), 98–105. <https://doi.org/10.1016/j.foodchem.2007.07.067>
- Amiri Samani, S., & Najji, M. H. (2019). Effect of homogenizer pressure and temperature on physicochemical, oxidative stability, viscosity, droplet size, and sensory properties of Sesame vegetable cream. *Food Science and Nutrition*, 7(3), 899–906. <https://doi.org/10.1002/fsn3.680>
- Azuma, G., Kimura, N., Hosokawa, M., & Miyashita, K. (2009). Effect of droplet size on the oxidative stability of soybean oil tag and fish oil tag in oil-in-water emulsion. *Journal of Oleo Science*, 58(6), 329–338. <https://doi.org/10.5650/jos.58.329>
- Beltrán, G., Bucheli, M. E., Aguilera, M. P., Belaj, A., & Jimenez, A. (2016). Squalene in virgin olive oil: Screening of variability in olive cultivars. *European Journal of Lipid Science and Technology*, 118(8), 1250–1253. <https://doi.org/10.1002/ejlt.201500295>
- Binsi, P. K., Nayak, N., Sarkar, P. C., Jeyakumari, A., Muhamed Ashraf, P., Ninan, G., & Ravishankar, C. N. (2017). Structural and oxidative stabilization of spray dried fish oil microencapsulates with gum arabic and sage polyphenols: Characterization and release kinetics. *Food Chemistry*, 219, 158–168. <https://doi.org/10.1016/j.foodchem.2016.09.126>
- Botrel, D. A., Borges, S. V., Fernandes, R. V. D. B., & Lourenço Do Carmo, E. (2014). Optimization of Fish Oil Spray Drying Using a Protein:Inulin System. *Drying Technology*, 32(3), 279–290. <https://doi.org/10.1080/07373937.2013.823621>
- Carmona, P. A. O., Tonon, R. V., da Cunha, R. L., & Hubinger, M. D. (2013). Influence of Emulsion Properties on the Microencapsulation of Orange Essential Oil by Spray Drying. *Journal of Colloid Science and Biotechnology*, 2(2), 130–139. <https://doi.org/10.1166/jcsb.2013.1042>
- Carneiro, H. C. F., Tonon, R. V., Grosso, C. R. F., & Hubinger, M. D. (2013). Encapsulation efficiency and oxidative stability of flaxseed oil microencapsulated by spray drying using different combinations of wall materials. *Journal of Food Engineering*, 115(4), 443–451. <https://doi.org/10.1016/j.jfoodeng.2012.03.033>

- Chegini, G., & Taheri, M. (2013). Whey powder: Process technology and physical properties: A review. *Middle East Journal of Scientific Research*, 13(10), 1377–1387. <https://doi.org/10.5829/idosi.mejsr.2013.13.10.1239>
- Corzo, O., Bracho, N., & Alvarez, C. (2010). Weibull model for thin-layer drying of mango slices at different maturity stages. *Journal of Food Processing and Preservation*, 34(6), 993–1008. <https://doi.org/10.1111/j.1745-4549.2009.00433.x>
- Da Silva, A. C., & Jorge, N. (2014). Influence of *Lentinus edodes* and *Agaricus blazei* extracts on the prevention of oxidation and retention of tocopherols in soybean oil in an accelerated storage test. *Journal of Food Science and Technology*, 51(6), 1208–1212. <https://doi.org/10.1007/s13197-012-0623-1>
- Dahlquist-Willard, R. M., Marshall, M. N., Betts, S. L., Tuell-Todd, C. C., VanderGheynst, J. S., & Stapleton, J. J. (2016). Development and validation of a Weibull–Arrhenius model to predict thermal inactivation of black mustard (*Brassica nigra*) seeds under fluctuating temperature regimens. *Biosystems Engineering*, 151, 350–360. <https://doi.org/10.1016/j.biosystemseng.2016.09.015>
- Di Giorgio, L., Salgado, P. R., & Mauri, A. N. (2019). Encapsulation of fish oil in soybean protein particles by emulsification and spray drying. *Food Hydrocolloids*, 87(September 2018), 891–901. <https://doi.org/10.1016/j.foodhyd.2018.09.024>
- Edris, A. E., Kalemba, D., Adamiec, J., & Piaotkowski, M. (2016). Microencapsulation of *Nigella sativa* oleoresin by spray drying for food and nutraceutical applications. *Food Chemistry*, 204, 326–333. <https://doi.org/10.1016/j.foodchem.2016.02.143>
- Fang, X., Shima, M., & Adachi, S. (2005). Effects of drying conditions on the oxidation of linoleic acid encapsulated with gum arabic by spray-drying. *Food Science and Technology Research*, 11(4), 380–384. <https://doi.org/10.3136/fstr.11.380>
- Fermin Jimenez, J. A. (2019). イサダオイルの乳化噴霧乾燥粉末化と作成粉末の特質. Kagawa University.
- Fernandes, L. P., Turatti, I. C. C., Lopes, N. P., Ferreira, J. C., Candido, R. C., & Oliveira, W. P. (2008). Volatile retention and antifungal properties of spray-dried microparticles of *Lippia sidoides* essential oil. *Drying Technology*, 26(12), 1534–1542. <https://doi.org/10.1080/07373930802464034>
- Fernandes, R. V. D. B., Borges, S. V., & Botrel, D. A. (2014). Gum arabic/starch/maltodextrin/inulin as wall materials on the microencapsulation of rosemary essential oil. *Carbohydrate Polymers*, 101(1), 524–532. <https://doi.org/10.1016/j.carbpol.2013.09.083>
- Frascareli, E. C., Silva, V. M., Tonon, R. V., & Hubinger, M. D. (2012). Effect of process conditions on the microencapsulation of coffee oil by spray drying. *Food and Bioprocess Processing*, 90(3), 413–424. <https://doi.org/10.1016/j.fbp.2011.12.002>
- Gallardo, G., Guida, L., Martinez, V., López, M. C., Bernhardt, D., Blasco, R., ... Hermida, L. G. (2013). Microencapsulation of linseed oil by spray drying for functional food application. *Food Research International*, 52(2), 473–482.

- <https://doi.org/10.1016/j.foodres.2013.01.020>
- Gallego, M. G., Gordon, M. H., Segovia, F. J., Skowrya, M., & Almajano, M. P. (2013). Antioxidant properties of three aromatic herbs (rosemary, thyme and lavender) in oil-in-water emulsions. *JAACS, Journal of the American Oil Chemists' Society*, *90*(10), 1559–1568. <https://doi.org/10.1007/s11746-013-2303-3>
- Garcia, L. C., Tonon, R. V., & Hubinger, M. D. (2012). Effect of Homogenization Pressure and Oil Load on the Emulsion Properties and the Oil Retention of Microencapsulated Basil Essential Oil (*Ocimum basilicum* L.). *Drying Technology*, *30*(13), 1413–1421. <https://doi.org/10.1080/07373937.2012.685998>
- Gharsallaoui, A., Roudaut, G., Beney, L., Chambin, O., Voilley, A., & Saurel, R. (2012). Properties of spray-dried food flavours microencapsulated with two-layered membranes: Roles of interfacial interactions and water. *Food Chemistry*, *132*(4), 1713–1720. <https://doi.org/10.1016/j.foodchem.2011.03.028>
- Ghosh, M., Upadhyay, R., Mahato, D. K., & Mishra, H. N. (2019). Kinetics of lipid oxidation in omega fatty acids rich blends of sunflower and sesame oils using Rancimat. *Food Chemistry*, *272*(August 2018), 471–477. <https://doi.org/10.1016/j.foodchem.2018.08.072>
- Grigorakis, K., Giogios, I., Vasilaki, A., & Nengas, I. (2010). Effect of the fish oil, oxidation status and of heat treatment temperature on the volatile compounds of the produced fish feeds. *Animal Feed Science and Technology*, *158*(1–2), 73–84. <https://doi.org/10.1016/j.anifeedsci.2010.03.012>
- Heidarpour, M., & Farhoosh, R. (2018). A preliminary Rancimat-based kinetic approach of detecting olive oil adulteration. *LWT - Food Science and Technology*, *90*(July 2017), 77–82. <https://doi.org/10.1016/j.lwt.2017.12.015>
- Hoang Thi, T. H., Lemdani, M., & Flament, M. P. (2013). Use of calcium caseinate in association with lecithin for masking the bitterness of acetaminophen - Comparative study with sodium caseinate. *International Journal of Pharmaceutics*, *456*(2), 382–389. <https://doi.org/10.1016/j.ijpharm.2013.08.075>
- Hogan, S. A., McNamee, B. F., O’Riordan, E. D., & O’Sullivan, M. (2001). Emulsification and microencapsulation properties of sodium caseinate/carbohydrate blends. *International Dairy Journal*, *11*(3), 137–144. [https://doi.org/10.1016/S0958-6946\(01\)00091-7](https://doi.org/10.1016/S0958-6946(01)00091-7)
- Huang, H., Hao, S., Li, L., Yang, X., Cen, J., Lin, W., & Wei, Y. (2014). Influence of emulsion composition and spray-drying conditions on microencapsulation of tilapia oil. *Journal of Food Science and Technology*, *51*(9), 2148–2154. <https://doi.org/10.1007/s13197-012-0711-2>
- Huynh, T. V., Caffin, N., Dykes, G., & Bhandari, B. (2008). Optimization of the microencapsulation of lemon myrtle oil using response surface methodology. *Drying Technology*, *26*(3), 357–368. <https://doi.org/10.1080/07373930801898182>
- Ixtaina, V. Y., Julio, L. M., Wagner, J. R., Nolasco, S. M., & Tomás, M. C.

- (2015). Physicochemical characterization and stability of chia oil microencapsulated with sodium caseinate and lactose by spray-drying. *Powder Technology*, 271, 26–34.
<https://doi.org/10.1016/j.powtec.2014.11.006>
- Jafari, S. M., Assadpoor, E., He, Y., & Bhandari, B. (2008). Encapsulation efficiency of food flavours and oils during spray drying. *Drying Technology*, 26(7), 816–835. <https://doi.org/10.1080/07373930802135972>
- Jayasundera, M., Adhikari, B., Aldred, P., & Ghandi, A. (2009). Surface modification of spray dried food and emulsion powders with surface-active proteins: A review. *Journal of Food Engineering*, 93(3), 266–277.
<https://doi.org/10.1016/j.jfoodeng.2009.01.036>
- Jin, R., Yin, H., Wang, H., Zhang, D., Cao, K., & Yuan, C. (2018). Preparation and characterization of squalene microcapsules by complex coacervation. *Journal of Food Process Engineering*, 41(6), 1–6.
<https://doi.org/10.1111/jfpe.12847>
- Jin, Y., Liu, Z., Liu, D., Shi, G., Liu, D., Yang, Y., ... Zhou, Z. (2019). Natural antioxidant of rosemary extract used as an additive in the ultrasound-assisted extraction of anthocyanins from lingonberry (*Vaccinium vitis-idaea* L.) pomace. *Industrial Crops and Products*, 138(June), 111425.
<https://doi.org/10.1016/j.indcrop.2019.05.074>
- Jinapong, N., Suphantharika, M., & Jamnong, P. (2008). Production of instant soymilk powders by ultrafiltration, spray drying and fluidized bed agglomeration. *Journal of Food Engineering*, 84(2), 194–205.
<https://doi.org/10.1016/j.jfoodeng.2007.04.032>
- Kim, J. Y., Kim, M. J., Yi, B., Oh, S., & Lee, J. (2015). Antioxidant properties of ascorbic acid in bulk oils at different relative humidity. *Food Chemistry*, 176, 302–307. <https://doi.org/10.1016/j.foodchem.2014.12.079>
- Klinkesorn, U., Sophanodora, P., Chinachoti, P., Decker, E. A., & McClements, D. J. (2006). Characterization of spray-dried tuna oil emulsified in two-layered interfacial membranes prepared using electrostatic layer-by-layer deposition. *Food Research International*, 39(4), 449–457.
<https://doi.org/10.1016/j.foodres.2005.09.008>
- Kuriakose, R., & Anandharamakrishnan, C. (2010). Computational fluid dynamics (CFD) applications in spray drying of food products. *Trends in Food Science and Technology*, 21(8), 383–398.
<https://doi.org/10.1016/j.tifs.2010.04.009>
- Let, M. B., Jacobsen, C., Sørensen, A. D. M., & Meyer, A. S. (2007). Homogenization conditions affect the oxidative stability of fish oil enriched milk emulsions: Lipid oxidation. *Journal of Agricultural and Food Chemistry*, 55(5), 1773–1780. <https://doi.org/10.1021/jf062391s>
- Lethuaut, L., Métro, F., & Genot, C. (2002). *Effect of Droplet Size on Lipid Oxidation Rates of Oil Emulsion Stabilized By Protein*. 79(5), 425–430.
- Li, X., Li, Y., Yang, F., Liu, R., Zhao, C., Jin, Q., & Wang, X. (2019). Oxidation degree of soybean oil at induction time point under Rancimat test condition: Theoretical derivation and experimental observation. *Food Research International*, 120(March), 756–762.

- <https://doi.org/10.1016/j.foodres.2018.11.036>
- Liu, Y. H., Li, X. F., Zhu, W. X., Luo, L., Duan, X., & Yin, Y. (2016). Drying characteristics, kinetics model and effective moisture diffusivity of vacuum far-infrared dried *Rehmanniae*. *International Journal of Agricultural and Biological Engineering*, 9(5), 208–217. <https://doi.org/10.3965/j.ijabe.20160905.2082>
- Liu, Y., Hou, Z., Yang, J., & Gao, Y. (2015). Effects of antioxidants on the stability of β -Carotene in O/W emulsions stabilized by Gum Arabic. *Journal of Food Science and Technology*, 52(6), 3300–3311. <https://doi.org/10.1007/s13197-014-1380-0>
- Luo, Y., Zhang, B., Whent, M., Yu, L. L., & Wang, Q. (2011). Preparation and characterization of zein/chitosan complex for encapsulation of α -tocopherol, and its in vitro controlled release study. *Colloids and Surfaces B: Biointerfaces*, 85(2), 145–152. <https://doi.org/10.1016/j.colsurfb.2011.02.020>
- Martínez, M. L., Curti, M. I., Rocchia, P., Llabot, J. M., Penci, M. C., Bodoira, R. M., & Ribotta, P. D. (2015). Oxidative stability of walnut (*Juglans regia* L.) and chia (*Salvia hispanica* L.) oils microencapsulated by spray drying. *Powder Technology*, 270(Part A), 271–277. <https://doi.org/10.1016/j.powtec.2014.10.031>
- Miranda, M., Vega-Gálvez, A., García, P., Di Scala, K., Shi, J., Xue, S., & Uribe, E. (2010). Effect of temperature on structural properties of Aloe vera (*Aloe barbadensis* Miller) gel and Weibull distribution for modelling drying process. *Food and Bioprocess Processing*, 88(2–3), 138–144. <https://doi.org/10.1016/j.fbp.2009.06.001>
- Munin, A., & Edwards-Lévy, F. (2011). Encapsulation of natural polyphenolic compounds; a review. In *Pharmaceutics* (Vol. 3). <https://doi.org/10.3390/pharmaceutics3040793>
- Na, H. S., Kim, J. N., Kim, J. M., & Lee, K. Y. (2011). Encapsulation of fish oil using Cyclodextrin and whey protein concentrate. *Biotechnology and Bioprocess Engineering*, 16(6), 1077–1082. <https://doi.org/10.1007/s12257-011-0099-2>
- Nakazawa, R., Shima, M., & Adachi, S. (2008). Effect of oil-droplet size on the oxidation of microencapsulated methyl linoleate. *Journal of Oleo Science*, 57(4), 225–232. <https://doi.org/10.5650/jos.57.225>
- Naziri, E., Consonni, R., & Tsimidou, M. Z. (2014). Squalene oxidation products: Monitoring the formation, characterisation and pro-oxidant activity. *European Journal of Lipid Science and Technology*, 116(10), 1400–1411. <https://doi.org/10.1002/ejlt.201300506>
- Ng, S. K., Choong, Y. H., Tan, C. P., Long, K., & Nyam, K. L. (2014). Effect of total solids content in feed emulsion on the physical properties and oxidative stability of microencapsulated kenaf seed oil. *LWT - Food Science and Technology*, 58(2), 627–632. <https://doi.org/10.1016/j.lwt.2014.03.010>
- Peanparkdee, M., Iwamoto, S., & Yamauchi, R. (2016). Microencapsulation: a Review of Applications in the Food and Pharmaceutical Industries. *Reviews in Agricultural Science*, 4(0), 56–65. <https://doi.org/10.7831/ras.4.56>
- Perrier-Cornet, J. M., Marie, P., & Gervais, P. (2005). Comparison of

- emulsification efficiency of protein-stabilized oil-in-water emulsions using jet, high pressure and colloid mill homogenization. *Journal of Food Engineering*, 66(2), 211–217. <https://doi.org/10.1016/j.jfoodeng.2004.03.008>
- Phunpee, S., Ruktanonchai, U. R., Yoshii, H., Assabumrungrat, S., & Soottitantawat, A. (2017). Encapsulation of lemongrass oil with cyclodextrins by spray drying and its controlled release characteristics. *Bioscience, Biotechnology and Biochemistry*, 81(4), 718–723. <https://doi.org/10.1080/09168451.2016.1277942>
- Pourashouri, P., Shabanpour, B., Razavi, S. H., Jafari, S. M., Shabani, A., & Aubourg, S. P. (2014). Impact of Wall Materials on Physicochemical Properties of Microencapsulated Fish Oil by Spray Drying. *Food and Bioprocess Technology*, 7(8), 2354–2365. <https://doi.org/10.1007/s11947-013-1241-2>
- Pratap Singh, A., Siddiqui, J., & Diosady, L. L. (2018). Characterizing the pH-Dependent Release Kinetics of Food-Grade Spray Drying Encapsulated Iron Microcapsules for Food Fortification. *Food and Bioprocess Technology*, 11(2), 435–446. <https://doi.org/10.1007/s11947-017-2022-0>
- Qian, C., & McClements, D. J. (2011). Formation of nanoemulsions stabilized by model food-grade emulsifiers using high-pressure homogenization: Factors affecting particle size. *Food Hydrocolloids*, 25(5), 1000–1008. <https://doi.org/10.1016/j.foodhyd.2010.09.017>
- R.G. Kumar, L., Chatterjee, N. S., Tejpal, C. S., Vishnu, K. V., Anas, K. K., Asha, K. K., ... Mathew, S. (2017). Evaluation of chitosan as a wall material for microencapsulation of squalene by spray drying: Characterization and oxidative stability studies. *International Journal of Biological Macromolecules*, 104, 1986–1995. <https://doi.org/10.1016/j.ijbiomac.2017.03.114>
- Ramakrishnan, S., Ferrando, M., Aceña-Muñoz, L., Mestres, M., De Lamo-Castellví, S., & Güell, C. (2014). Influence of Emulsification Technique and Wall Composition on Physicochemical Properties and Oxidative Stability of Fish Oil Microcapsules Produced by Spray Drying. *Food and Bioprocess Technology*, 7(7), 1959–1972. <https://doi.org/10.1007/s11947-013-1187-4>
- Reddy, L. H., & Couvreur, P. (2009). Squalene: A natural triterpene for use in disease management and therapy. *Advanced Drug Delivery Reviews*, 61(15), 1412–1426. <https://doi.org/10.1016/j.addr.2009.09.005>
- Shamaei, S., Seiiedlou, S. S., Aghbashlo, M., Tsotsas, E., & Kharaghani, A. (2017). Microencapsulation of walnut oil by spray drying: Effects of wall material and drying conditions on physicochemical properties of microcapsules. *Innovative Food Science and Emerging Technologies*, 39, 101–112. <https://doi.org/10.1016/j.ifset.2016.11.011>
- Shen, Z., Augustin, M. A., Sanguansri, L., & Cheng, L. J. (2010). Oxidative stability of microencapsulated fish oil powders stabilized by blends of chitosan, modified starch, and glucose. *Journal of Agricultural and Food Chemistry*, 58(7), 4487–4493. <https://doi.org/10.1021/jf904102k>
- Shiga, H., Adachi, S., Adachi, S., & Yoshii, H. (2014). A simple method for determining the flaxseed or fish oil content with N, N-dimethylformamide in

- microcapsules prepared by spray drying. *Japan Journal of Food Engineering*, 15(3), 131–139. <https://doi.org/10.11301/jsfe.15.131>
- Shiga, H., Neoh, T. L., Ninomiya, A., Adachi, S., Pasten, I. L., Adachi, S., & Yoshii, H. (2017). Effect of oil droplet size on the oxidative stability of spray-dried flaxseed oil powders. *Bioscience, Biotechnology and Biochemistry*, 81(4), 698–704. <https://doi.org/10.1080/09168451.2017.1281720>
- Shishir, M. R. I., & Chen, W. (2017). Trends of spray drying: A critical review on drying of fruit and vegetable juices. *Trends in Food Science and Technology*, 65, 49–67. <https://doi.org/10.1016/j.tifs.2017.05.006>
- Soottitantawat, A., Yoshii, H., Furuta, T., Ohkawara, M., & Linko, P. (2003). Microencapsulation by spray drying: Influence of emulsion size on the retention of volatile compounds. *Journal of Food Science*, 68(7), 2256–2262. <https://doi.org/10.1111/j.1365-2621.2003.tb05756.x>
- Spanova, M., & Daum, G. (2011). Squalene - biochemistry, molecular biology, process biotechnology, and applications. *European Journal of Lipid Science and Technology*, 113(11), 1299–1320. <https://doi.org/10.1002/ejlt.201100203>
- Sultana, A., Tanaka, Y., Fushimi, Y., & Yoshii, H. (2018). Stability and release behavior of encapsulated flavor from spray-dried *Saccharomyces cerevisiae* and maltodextrin powder. *Food Research International*, 106(January), 809–816. <https://doi.org/10.1016/j.foodres.2018.01.059>
- Sultana, A., & Yoshii, H. (2019). Kinetic study of controlled release of flavor compounds from spray-dried encapsulated yeast powder using dynamic vapor sorption–gas chromatography. *Bioscience, Biotechnology and Biochemistry*, 83(4), 738–746. <https://doi.org/10.1080/09168451.2018.1564618>
- Tadros, T., Izquierdo, P., Esquena, J., & Solans, C. (2004). Formation and stability of nano-emulsions. *Advances in Colloid and Interface Science*, 108–109, 303–318. <https://doi.org/10.1016/j.cis.2003.10.023>
- Teixeira, M. I., Andrade, L. R., Farina, M., & Rocha-Leao, M. H. M. (2004). Characterization of short chain fatty acid microcapsules produced by spray drying. *Materials Science and Engineering C*, 24(5), 653–658. <https://doi.org/10.1016/j.msec.2004.08.008>
- Terpinc, P., Bezjak, M., & Abramovič, H. (2009). A kinetic model for evaluation of the antioxidant activity of several rosemary extracts. *Food Chemistry*, 115(2), 740–744. <https://doi.org/10.1016/j.foodchem.2008.12.033>
- Vehring, R. (2008). Pharmaceutical particle engineering via spray drying. *Pharmaceutical Research*, 25(5), 999–1022. <https://doi.org/10.1007/s11095-007-9475-1>
- Walzel, P. (2011). Influence of the spray method on product quality and morphology in spray drying. *Chemical Engineering and Technology*, 34(7), 1039–1048. <https://doi.org/10.1002/ceat.201100051>
- Wang, S., Shi, Y., Tu, Z., Zhang, L., Wang, H., Tian, M., & Zhang, N. (2017). Influence of soy lecithin concentration on the physical properties of whey protein isolate-stabilized emulsion and microcapsule formation. *Journal of Food Engineering*, 207, 73–80.

<https://doi.org/10.1016/j.jfoodeng.2017.03.020>

- Watanabe, Y., Fang, X., Adachi, S., Fukami, H., & Matsuno, R. (2004). Oxidation of 6-O-arachidonoyl L-ascorbate microencapsulated with a polysaccharide by spray-drying. *LWT - Food Science and Technology*, 37(4), 395–400. <https://doi.org/10.1016/j.lwt.2003.10.003>
- Wu, F., Shi, X., Zou, H., Zhang, T., Dong, X., Zhu, R., & Yu, C. (2019). Effects of high-pressure homogenization on physicochemical, rheological and emulsifying properties of myofibrillar protein. *Journal of Food Engineering*, 263(May), 272–279. <https://doi.org/10.1016/j.jfoodeng.2019.07.009>
- Wu, Q., Uluata, S., Cui, L., Wang, C., Li, D., McClements, J., & Decker, E. A. (2016). Physical and oxidation stability of self-emulsifying krill oil-in-water emulsions. *Food and Function*, 7(8), 3590–3598. <https://doi.org/10.1039/c6fo00045b>
- Xiao, H., Yao, Z., Peng, Q., Ni, F., Sun, Y., Zhang, C. X., & Zhong, Z. X. (2016). Extraction of squalene from camellia oil by silver ion complexation. *Separation and Purification Technology*, 169, 196–201. <https://doi.org/10.1016/j.seppur.2016.05.041>
- Yang, X., Tian, H., Ho, C. T., & Huang, Q. (2011). Inhibition of citral degradation by oil-in-water nanoemulsions combined with antioxidants. *Journal of Agricultural and Food Chemistry*, 59(11), 6113–6119. <https://doi.org/10.1021/jf2012375>
- Yang, Ying, Marshall-Breton, C., Leser, M. E., Sher, A. A., & McClements, D. J. (2012). Fabrication of ultrafine edible emulsions: Comparison of high-energy and low-energy homogenization methods. *Food Hydrocolloids*, 29(2), 398–406. <https://doi.org/10.1016/j.foodhyd.2012.04.009>
- Yang, Yong, Song, X., Sui, X., Qi, B., Wang, Z., Li, Y., & Jiang, L. (2016). Rosemary extract can be used as a synthetic antioxidant to improve vegetable oil oxidative stability. *Industrial Crops and Products*, 80, 141–147. <https://doi.org/10.1016/j.indcrop.2015.11.044>
- Yeşilsu, A. F., & Özyurt, G. (2019). Oxidative stability of microencapsulated fish oil with rosemary, thyme and laurel extracts: A kinetic assessment. *Journal of Food Engineering*, 240(14), 171–182. <https://doi.org/10.1016/j.jfoodeng.2018.07.021>
- Yoshii, H., Soottitawat, A., Liu, X. D., Atarashi, T., Furuta, T., Aishima, S., ... Linko, P. (2001). Flavor release from spray-dried maltodextrin/gum arabic or soy matrices as a function of storage relative humidity. *Innovative Food Science and Emerging Technologies*, 2(1), 55–61. [https://doi.org/10.1016/S1466-8564\(01\)00019-4](https://doi.org/10.1016/S1466-8564(01)00019-4)