



DAFTAR PUSTAKA

- Alken, S., Carceller, F., Mandeville, H. and Saran, F., 2018. High-grade glioma. In *Radiation Oncology for Pediatric CNS Tumors* (pp. 229-248). Springer, Cham.
- Baig, M., Klein, J. and Mechtler, L. (2016). Imaging of brain tumors. *Continuum: Lifelong Learning in Neurology*, 22(5), pp.1529-1552.
- Bosma, I., Vos, M.J., Heimans, J.J., Taphoorn, M.J., Aaronson, N.K., Postma, T.J., van der Ploeg, H.M., Muller, M., Vandertop, W.P., Slotman, B.J. and Klein, M., 2007. The course of neurocognitive functioning in high-grade glioma patients. *Neuro-oncology*, 9(1), pp.53-62.
- Bulakbasi, N., Guvenc, I., Onguru, O., Erdogan, E., Tayfun, C. and Ucoz, T. (2004). The added value of the apparent diffusion coefficient calculation to magnetic resonance imaging in the differentiation and grading of malignant brain tumors. *Journal of Computer Assisted Tomography*, 28(6), pp.735-746.
- Engelhorn, T., Savaskan, N.E., Schwarz, M.A., Kreutzer, J., Meyer, E.P., Hahnen, E., Ganslandt, O., Dörfler, A., Nimsky, C., Buchfelder, M. and Eyüpoglu, I.Y., 2009. Cellular characterization of the peritumoral edema zone in malignant brain tumors. *Cancer Science*, 100(10), pp.1856-1862.
- Ferlay, J., Colombet, M., Soerjomataram, I., Mathers, C., Parkin, D.M., Piñeros, M., Znaor, A. and Bray, F., 2019. Estimating the global cancer incidence and mortality in 2018: GLOBOCAN sources and methods. *International Journal of Cancer*, 144(8), pp.1941-1953.
- Flint, J., Hansen, B., Vestergaard-Poulsen, P. and Blackband, S. (2009). Diffusion weighted magnetic resonance imaging of neuronal activity in the hippocampal slice model. *NeuroImage*, 46(2), pp.411-418.
- Geneidi, E., Habib, L., Chalabi, N. and Haschim, M. (2016). Potential role of quantitative MRI assessment in differentiating high from low-grade gliomas. *The Egyptian Journal of Radiology and Nuclear Medicine*, 47(1), pp.243-253.
- Johansen-Berg, H. and Behrens, T. (2014). *Diffusion MRI*. Amsterdam: Elsevier Science.
- Johnson, D.R., Guerin, J.B., Giannini, C., Morris, J.M., Eckel, L.J. and Kaufmann, T.J., 2017. 2016 updates to the WHO brain tumor classification system: what the radiologist needs to know. *Radiographics*, 37(7), pp.2164-2180



- Kesari, S., 2011, December. Understanding glioblastoma tumor biology: the potential to improve current diagnosis and treatments. In *Seminars in oncology* (Vol. 38, pp. S2-S10). WB Saunders.
- Kim, B. and Gutierrez, J. (2012). Contrast-enhanced MR imaging in neuroimaging. *Magnetic Resonance Imaging Clinics of North America*, 20(4), pp.649-685.
- Kimura, M. and da Cruz, L.C.H., 2016. Multiparametric MR imaging in the assessment of brain tumors. *Magnetic Resonance Imaging Clinics*, 24(1), pp.87-122.
- Kono, K., Inoue, Y., Nakayama, K., Shakudo, M., Morino, M., Ohata, K., et al., (2001). The role of diffusion-weighted imaging in patients with brain tumors. *American Journal of Neuroradiology*, 22(6): 1081–1088.
- Larjavaara, S., Mäntylä, R., Salminen, T., Haapasalo, H., Raitanen, J., Jääskeläinen, J. and Auvinen, A., 2007. Incidence of gliomas by anatomic location. *Neuro-oncology*, 9(3), pp.319-325.
- Lemercier, P., Maya, S., Patrie, J., Flors, L. and Leiva-Salinas, C. (2014). Gradient of apparent diffusion coefficient values in peritumoral edema helps in differentiation of glioblastoma from solitary metastatic lesions. *American Journal of Roentgenology*, 203(1), pp.163-169.
- Lequin, M. and Hendrikse, J. (2017). Advanced MR imaging in pediatric brain tumors, Clinical Applications. *Neuroimaging Clinics of North America*, 27(1), pp.167-190.
- Louis, D.N., Perry, A., Reifenberger, G., Von Deimling, A., Figarella-Branger, D., Cavenee, W.K., Ohgaki, H., Wiestler, O.D., Kleihues, P. and Ellison, D.W., 2016. The 2016 World Health Organization classification of tumors of the central nervous system: a summary. *Acta neuropathologica*, 131(6), pp.803-820.
- Nabors, L.B., 2016. Central nervous system: notable developments in the management of primary and recurrent gliomas. *Journal of the National Comprehensive Cancer Network*, 14(5S), pp.681-684.
- Ono, Y., Chernov, M.F., Muragaki, Y., Maruyama, T., Abe, K. and Iseki, H., 2018. Imaging of intracranial gliomas. In *Intracranial Gliomas Part I-Surgery* (Vol. 30, pp. 12-62). Karger Publishers.
- Ostrom, Q.T., Gittleman, H., Truitt, G., Boscia, A., Kruchko, C. and Barnholtz-Sloan, J.S., 2018. CBTRUS statistical report: primary brain and other central nervous system tumors diagnosed in the United States in 2011–2015. *Neuro-oncology*, 20(suppl_4), pp.iv1-iv86.
- Raisi-nafchi, M., Faeghi, F., Zali, A., Haghishatkhah, H. and Jalal-Shokouhi, J. (2016). Preoperative grading of astrocytic supratentorial brain tumors with



diffusion-weighted magnetic resonance imaging and apparent diffusion coefficient. *Iranian Journal of Radiology*, 13(3):e30426.

- Rees, J.H., Smirniotopoulos, J.G., Jones, R.V. and Wong, K., 1996. Glioblastoma multiforme: radiologic-pathologic correlation. *Radiographics*, 16(6), pp.1413-1438.
- Sastroasmoro, S. and Ismael, S. (2011). Usulan Penelitian. in *Dasar-Dasar Metodologi Penelitian Klinis*. 4th edn. Jakarta: Sagung Seto, 31–63.
- Schaefer, P. W., Grant, P. E. and Gonzalez, R. G. (2000). Diffusion-weighted MR Imaging of the Brain. *Radiology*, 217(2):331–345.
- Schiffer, D., 2006. *Brain tumor pathology: current diagnostic hotspots and pitfalls*. Springer Science & Business Media.
- Schoenegger, K., Oberndorfer, S., Wuschitz, B., Struhal, W., Hainfellner, J., Prayer, D., Heinzl, H., Lahrmann, H., Marosi, C. and Grisold, W., 2009. Peritumoral edema on MRI at initial diagnosis: an independent prognostic factor for glioblastoma?. *European Journal of Neurology*, 16(7), pp.874-878.
- Shakal, A.A.S. and Mokbel, E.A.H., 2014. Hemorrhage after stereotactic biopsy from intra-axial brain lesions: incidence and avoidance. *Journal of Neurological Surgery Part A: Central European Neurosurgery*, 75(03), pp.177-182.
- Smirniotopoulos, J., Murphy, F., Rushing, E., Rees, J. and Schroeder, J. (2007). Patterns of Contrast Enhancement in the Brain and Meninges. *RadioGraphics*, 27(2), pp.525-551.
- Svolos, P., Tsolaki, E., Kapsalaki, E., Theodorou, K., Fountas, K., Fezoulidis, I. and Tsougos, I., 2013. Investigating brain tumor differentiation with diffusion and perfusion metrics at 3T MRI using pattern recognition techniques. *Magnetic Resonance Imaging*, 31(9), pp.1567-1577.
- Wang, S., Meng, M., Zhang, X., Wu, C., Wang, R., Wu, J., Sami, M.U. and Xu, K., 2018. Texture analysis of diffusion weighted imaging for the evaluation of glioma heterogeneity based on different regions of interest. *Oncology letters*, 15(5), pp.7297-7304.
- Wen, P.Y. and Kesari, S., 2008. Malignant gliomas in adults. *New England Journal of Medicine*, 359(5), pp.492-507.
- Withiwej, T., Thongnual, K., Wongprapairoj, S., Chawalparit, O., Piyapittayanan, S. and Sangruji, T., 2017. Correlation between magnetic resonance spectroscopy and perfusion at non-enhancing edematous area and tumor recurrence in high-grade gliomas after maximal safe resection. *Journal of The Medical Association of Thailand*, 100(4), p.79.



UNIVERSITAS
GADJAH MADA

**AKURASI NILAI DIFFUSION WEIGHTED IMAGING DAN APPARENT DIFFUSION COEFFICIENT PADA
EDEMA PERITUMORAL
DALAM MENENTUKAN DERAJAT KEGANASAN ASTROSITOMA**
MUHAMMAD ICHSAN N, dr. Yana Supriatna, Ph.D, SpRad (K); dr. Sri Retna Dwidianarti, SpRad (K)

Universitas Gadjah Mada, 2019 | Diunduh dari <http://etd.repository.ugm.ac.id/>

- Wu, C.X., Lin, G.S., Lin, Z.X., Zhang, J.D., Liu, S.Y. and Zhou, C.F., 2015. Peritumoral edema shown by MRI predicts poor clinical outcome in glioblastoma. *World Journal of Surgical Oncology*, 13(1), p.97.
- Xiao, H.F., Chen, Z.Y., Lou, X., Wang, Y.L., Gui, Q.P., Wang, Y., Shi, K.N., Zhou, Z.Y., Zheng, D.D., Wang, D.J. and Ma, L., 2015. Astrocytic tumour grading: a comparative study of three-dimensional pseudocontinuous arterial spin labelling, dynamic susceptibility contrast-enhanced perfusion-weighted imaging, and diffusion-weighted imaging. *European Radiology*, 25(12), pp.3423-3430.
- Zhang, L., Min, Z., Tang, M., Chen, S., Lei, X. and Zhang, X., 2017. The utility of diffusion MRI with quantitative ADC measurements for differentiating high-grade from low-grade cerebral gliomas: evidence from a meta-analysis. *Journal of the Neurological Sciences*, 373, pp.9-15.