

## DAFTAR PUSTAKA

1. Jiang H, Qiu X, Dai J, et al. Association of rs11190870 near LBX1 with adolescent idiopathic scoliosis susceptibility in a Han Chinese population. *Eur Spine J.* 2013;22(2):282-286. doi:10.1007/s00586-012-2532-4
2. Choudhry MN, Ahmad Z, Verma R. The Open Orthopaedics Journal Adolescent Idiopathic Scoliosis. *Open Orthop J.* 2016;10(16):143-154. doi:10.2174/1874325001610010143
3. Miller MD, Brinker MR. *Review of Orthopaedics.* 3rd ed. Saunders Elseviers; 2000.
4. Konieczny MR, Senyurt H, Krauspe R. Epidemiology of adolescent idiopathic scoliosis. *J Child Orthop.* 2013;7(1):3-9. doi:10.1007/s11832-012-0457-4
5. Wong C, Gosvig K, Sonne-Holm S. The role of the paravertebral muscles in adolescent idiopathic scoliosis evaluated by temporary paralysis. *Scoliosis Spinal Disord.* 2017;12(1):1-8. doi:10.1186/s13013-017-0138-7
6. L.S. N, R. H, P.C. N, K.P. N, A.T. S. Prevalence of scoliosis among school students in a town in southern Brazil. *Sao Paulo Med J.* 2010;128(2):69-73. <http://www.embase.com/search/results?subaction=viewrecord&from=export&id=L359380812%5Cnhttp://www.scielo.br/pdf/spmj/v128n2/a05v1282.pdf%5Cnhttp://elvis.ubvu.vu.nl:9003/vulink?sid=EMBASE&issn=15163180&id=doi:&atitle=Prevalence+of+scoliosis+among+school+st.>
7. Takahashi Y, Kou I, Takahashi A, et al. A genome-wide association study

- identifies common variants near LBX1 associated with adolescent idiopathic scoliosis. *Nat Genet.* 2011;43(12):1237-1240. doi:10.1038/ng.974
8. Guo L, Yamashita H, Kou I, et al. Functional Investigation of a Non-coding Variant Associated with Adolescent Idiopathic Scoliosis in Zebrafish: Elevated Expression of the Ladybird Homeobox Gene Causes Body Axis Deformation. *PLoS Genet.* 2016;12(1):1-28. doi:10.1371/journal.pgen.1005802
  9. Jagla K, Dollé P, Mattei MG, et al. Mouse Lbx1 and human LBX1 define a novel mammalian homeobox gene family related to the Drosophila lady bird genes. *Mech Dev.* 1995;53(3):345-356. <http://www.ncbi.nlm.nih.gov/pubmed/8645601>. Accessed June 12, 2019.
  10. Cao Y, Min J, Zhang Q, Li H, Li H. Associations of LBX1 gene and adolescent idiopathic scoliosis susceptibility: A meta-analysis based on 34,626 subjects. *BMC Musculoskelet Disord.* 2016;17(1):1-10. doi:10.1186/s12891-016-1139-z
  11. Gross MK, Dottori M, Goulding M. Lbx1 specifies somatosensory association interneurons in the dorsal spinal cord. *Neuron.* 2002;34(4):535-549. <http://www.ncbi.nlm.nih.gov/pubmed/12062038>. Accessed June 12, 2019.
  12. Leboeuf D, Letellier K, Alos N, Edery P, Moldovan F. Do estrogens impact adolescent idiopathic scoliosis? *Trends Endocrinol Metab.* 2009;20(4):147-152. doi:10.1016/j.tem.2008.12.004
  13. Skibinska I, Tomaszewski M, Andrusiewicz M, et al. Expression of estrogen

- receptor coactivator proline-, glutamic acid- and leucine-rich protein 1 within paraspinal muscles in adolescents with idiopathic scoliosis. *PLoS One*. 2016;11(4):1-13. doi:10.1371/journal.pone.0152286
14. Grauers A, Einarsdottir E, Gerdhem P. Genetics and pathogenesis of idiopathic scoliosis. *Scoliosis Spinal Disord*. 2016;11(1):1-7. doi:10.1186/s13013-016-0105-8
  15. Apley AG, Solomon L. *Apley and Solomon's System of Orthopaedics and Trauma*. 10th ed. (Blom AW, Warwick D, Whitehouse MR, eds.). CRC Press; 2018.
  16. Herkowitz HN, Garfin SR, Eismont FJ, Bell GR. *Rothman - Simeone : The Spine 6th Edition*. 6th ed. Philadelphia: Elseviers; 2011.
  17. Garland HG. HEREDITARY SCOLIOSIS. *BMJ*. 1934;1(3816):328-328. doi:10.1136/bmj.1.3816.328
  18. Wynne-Davies R. Familial (idiopathic) scoliosis. A family survey. *J Bone Joint Surg Br*. 1968;50(1):24-30. <http://www.ncbi.nlm.nih.gov/pubmed/5641594>. Accessed October 14, 2019.
  19. Riseborough EJ, Wynne-Davies R. A genetic survey of idiopathic scoliosis in Boston, Massachusetts. *J Bone Joint Surg Am*. 1973;55(5):974-982. <http://www.ncbi.nlm.nih.gov/pubmed/4760104>. Accessed October 14, 2019.
  20. Knott P, Pappo E, Cameron M, et al. SOSORT 2012 consensus paper: reducing x-ray exposure in pediatric patients with scoliosis The epidemiology of x-ray exposure in pediatric patients with spinal deformity. 2014:1-9.

21. Vaccarro AR. *Spine Core Knowledge in Orthopaedics*. Philadelphia: Mosby; 2005.
22. Lenke LG, Betz RR, Harms J, et al. Adolescent Idiopathic Scoliosis. *J Bone Jt Surgery-American Vol.* 2001;83(8):1169-1181. doi:10.2106/00004623-200108000-00006
23. Carlson BB, Burton DC, Asher MA. Comparison of trunk and spine deformity in adolescent idiopathic scoliosis. *Scoliosis.* 2013;8(1):2-5. doi:10.1186/1748-7161-8-2
24. Marengo L, Cravino M, Canavese F, Andreacchio A. Classification of adolescent idiopathic scoliosis. *Scoliosis Diagnosis, Classif Manag Options.* 2018:35-44. doi:10.1007/s11832-012-0459-2
25. Jiang J, Qiu Y, Mao S, Zhao Q, Qian B, Zhu F. The influence of elastic orthotic belt on sagittal profile in adolescent idiopathic thoracic scoliosis: A comparative radiographic study with Milwaukee brace. *BMC Musculoskelet Disord.* 2010;11(Figure 1):1-5. doi:10.1186/1471-2474-11-219
26. Weinstein SL, Zavala DC, Ponseti I V. Idiopathic scoliosis. Long-term follow-up and prognosis in untreated patients. *J Bone Jt Surg - Ser A.* 1981;63(5):702-712. doi:10.2106/00004623-198163050-00003
27. Moreau A, Wang DS, Forget S, et al. Melatonin signaling dysfunction in adolescent idiopathic scoliosis. *Spine (Phila Pa 1976).* 2004;29(16):1772-1781. <http://www.ncbi.nlm.nih.gov/pubmed/15303021>. Accessed June 12, 2019.
28. Burwell RG, Dangerfield PH, Moulton A, Grivas TB. Adolescent idiopathic

- scoliosis (AIS), environment, exposome and epigenetics: A molecular perspective of postnatal normal spinal growth and the etiopathogenesis of AIS with consideration of a network approach and possible implications for medical therapy. *Scoliosis*. 2011;6(1):1-20. doi:10.1186/1748-7161-6-26
29. Sharma S, Londono D, Eckalbar WL, et al. A PAX1 enhancer locus is associated with susceptibility to idiopathic scoliosis in females. *Nat Commun*. 2015;6:1-10. doi:10.1038/ncomms7452
  30. LBX1. <https://www.genecards.org/cgi-bin/carddisp.pl?gene=LBX1>.
  31. Kudo D, Miyakoshi N, Hongo M, et al. LBX1 mRNA Expression in Paravertebral Muscles of Patients with Adolescent Idiopathic Scoliosis : A Preliminary Study. *Akita Univ*. 2013;(40):151-156.
  32. Livak KJ, Schmittgen TD. Analysis of Relative Gene Expression Data Using Real- Time Quantitative PCR and the 2<sup>-ΔΔC<sub>T</sub></sup> Method. 2001;408:402-408. doi:10.1006/meth.2001.1262
  33. Hitesh SS, Yang NMJ. Idiopathic scoliosis in Korean schoolchildren : a prospective screening study of over 1 million children. 2011:1087-1094. doi:10.1007/s00586-011-1695-8
  34. Schlösser TPC, Van Der Heijden GJMG, Versteeg AL, Castelein RM. How “idiopathic” is adolescent idiopathic scoliosis? A systematic review on associated abnormalities. *PLoS One*. 2014;9(5):1-10. doi:10.1371/journal.pone.0097461
  35. Kudo D, Miyakoshi N, Hongo M, et al. Nerve growth factor and estrogen receptor mRNA expression in paravertebral muscles of patients with

- adolescent idiopathic scoliosis: A preliminary study. *Spine Deform.* 2015;3(2):122-127. doi:10.1016/j.jspd.2014.07.006
36. Saka K. [Biomechanical analysis of scoliosis and back muscles using CT evaluation and the finite element method]. *Nihon Seikeigeka Gakkai Zasshi.* 1987;61(4):299-310. <http://www.ncbi.nlm.nih.gov/pubmed/3624957>. Accessed October 14, 2019.
37. Modi HN, Suh SW, Yang JH, Hong JY, Venkatesh KP, Muzaffar N. Spontaneous regression of curve in immature idiopathic scoliosis - does spinal column play a role to balance? An observation with literature review. *J Orthop Surg Res.* 2010;5(1):1-8. doi:10.1186/1749-799X-5-80