

DAFTAR PUSTAKA

- Access, O., Nardone, R., Brigo, F. & Trinka, E. 2016. Acute symptomatic seizures caused by electrolyte disturbance, 12(1), pp. 21–33. doi: 10.3988/jcn.2016.12.1.21.
- Agius Anastasi, A., Falzon, O., Camilleri, K., Vella, M. & Muscat, R. 2017. Brain symmetry index in healthy and stroke patients for assessment and prognosis', *Stroke Research and Treatment*, 2017. doi: 10.1155/2017/8276136.
- Alia, C., Spalletti, C., Lai, S., Panarese, A., Lamola, G., Bertolucci, F., Vallone, F., Di Garbo, A., Chisari, C., Micera, S. & Caleo, M. 2017. Neuroplastic Changes Following Brain Ischemia and their Contribution to Stroke Recovery: Novel Approaches in Neurorehabilitation, *Frontiers in Cellular Neuroscience*, 11(March), pp. 1–22. doi: 10.3389/fncel.2017.00076.
- Alvarez-Sabín, J., Montaner, J., Padró, L., Molina, C. A., Rovira, R., Codina, A. & Quintana, M. 2002. Gabapentin in late-onset poststroke seizures', *Neurology*, 59(12), pp. 1991–1993. doi: 10.1212/01.WNL.0000038388.57824.B6.
- Appelros, P., Nydevik, I. & Viitanen, M. 2003. Poor outcome after first-ever stroke: Predictors for death, dependency, and recurrent stroke within the first year, *Stroke*, 34(1), pp. 122–126. doi: 10.1161/01.STR.0000047852.05842.3C.
- Assenza, G., Zappasodi, F., Pasqualetti, P., Vernieri, F. & Tecchio, F. 2013. A contralesional EEG power increase mediated by interhemispheric disconnection provides negative prognosis in acute stroke, *Restorative Neurology and Neuroscience*, 31(2), pp. 177–188. doi: 10.3233/RNN-120244.
- Ayerbe, L., Ayis, S., Crichton, S., Wolfe, C. D. A. & Rudd, A. G. 2014. The long-term outcomes of depression up to 10 years after stroke; The south london stroke register, *Journal of Neurology, Neurosurgery and Psychiatry*, 85(5), pp. 514–521. doi: 10.1136/jnnp-2013-306448.
- Bauer, G., Bauer, R. 2016. EEG, Drug Effects, and Central Nervous System Poisoning. Available at www.neuropsychology.com
- Beckham, J. D. & Tyler, K. L. 2012. Neuro-Intensive Care of Patients with Acute CNS Infections, *Neurotherapeutics*, 9, pp. 124–138. doi: 10.1007/s13311-011-0086-5.
- Beghi, E. Alessandro, R. & Beretta, S. 2011. Incidence and predictors of acute symptomatic seizures after stroke, *Neurology*, 77(20), pp. 1785–1793. doi: 10.1212/WNL.0b013e3182364878.
- Bentes, C., Peralta, A. R., Viana, P., Martins, H., Morgado, C., Casimiro, C., Franco, A. C., Fonseca, A. C., Geraldes, R., Canhão, P., Pinho e Melo, T., Paiva, T. & Ferro, J. M. 2018. Quantitative EEG and functional outcome following acute ischemic stroke, *Clinical Neurophysiology*. International Federation of Clinical Neurophysiology, 129(8), pp. 1680–1687. doi: 10.1016/j.clinph.2018.05.021.
- Between, R., Deficit, N., Functional, F., Shifts, O., First, D. & After, H. 2013. Relation Between Neurologic Deficit Severity, 43(6), pp. 1537–1541. doi:

10.1161/STROKEAHA.111.636928.RELATION.

- Bhardwaj, A. 2006. Neurological impact of vasopressin dysregulation and hyponatremia, *Annals of Neurology*, 59(2), pp. 229–236. doi: 10.1002/ana.20788.
- Bhattarai, S., Ning, Z.X. 2014. EEG and SPECT Change in Acute Ischemic Stroke. *J Neurol Neurophysiol* 5: 190
- Buckingham, S. C., Campbell, S. L., Haas, B. R., Montana, V., Robel, S., Ogunrinu, T. & Sontheimer, H. 2011. Glutamate release by primary brain tumors induces epileptic activity, *Nature Medicine*. Nature Publishing Group, 17(10), pp. 1269–1274. doi: 10.1038/nm.2453.
- Bureau, M., Azkoitia, J. M., Ezmendi, G., Manterola, I., Zabaleta, H., Perez, M. & Medina, J. 2007. Non-invasive, wireless and universal interface for the control of peripheral devices by means of head movements, *2007 IEEE 10th International Conference on Rehabilitation Robotics, ICORR'07*, 4(10), pp. 124–131. doi: 10.1109/ICORR.2007.4428417.
- Burneo, J. G., Fang, J. and Saposnik, G. 2010. Impact of seizures on morbidity and mortality after stroke: A Canadian multi-centre cohort study, *European Journal of Neurology*, 17(1), pp. 52–58. doi: 10.1111/j.1468-1331.2009.02739.x.
- Capes, S. E., Hunt, D., Malmberg, K., Pathak, P. & Gerstein, H. C. 2001. Stress hyperglycemia and prognosis of stroke in nondiabetic and diabetic patients: A systematic overview, *Stroke*, 32(10), pp. 2426–2432. doi: 10.1161/hs1001.096194.
- Castillo, J., Leira, R., García, M. M., Serena, J., Blanco, M. & Dávalos, A. 2004. Blood Pressure Decrease during the Acute Phase of Ischemic Stroke Is Associated with Brain Injury and Poor Stroke Outcome, *Stroke*, 35(2), pp. 520–527. doi: 10.1161/01.STR.0000109769.22917.B0.
- Cernik, D. Sanak, D. & Divisova, P. 2019. Impact of blood pressure levels within first 24 hours after mechanical thrombectomy on clinical outcome in acute ischemic stroke patients, *Journal of NeuroInterventional Surgery*, 11(8), pp. 735–739. doi: 10.1136/neurintsurg-2018-014548.
- Chen, W., Liu, G., Fang, J., Wang, Y., Song, Y., Pan, Y., Li, H., & Liu, L. 2016. External Validation of the Total Health Risks in Vascular Events Score to Predict Functional Outcome and Mortality in Patients Entered into the China National Stroke Registry. *J. Stroke Cerebrovasc. Dis.* 25.2331-2337.
- Cheng, B. Forkert, N.D, & Zavaglia, M. 2014. Influence of stroke infarct location on functional outcome measured by the modified rankin scale, *Stroke*, 45(6), pp. 1695–1702. doi: 10.1161/STROKEAHA.114.005152.
- Choi-Kwon, S., Han, S. W., Kwon, S. U. & Kim, J. S. 2005. Poststroke fatigue: Characteristics and related factors, *Cerebrovascular Diseases*, 19(2), pp. 84–90. doi: 10.1159/000082784.
- Cioncoloni, D., Piu, P., Tassi, R., Acampa, M., Guideri, F., Taddei, S., Bielli, S., Martini, G. & Mazzocchio, R. 2012. Relationship between the modified Rankin Scale and the Barthel Index in the process of functional recovery after stroke, *NeuroRehabilitation*, 30(4), pp. 315–322. doi: 10.3233/NRE-2012-0761.

- Colhoun, H. M., Betteridge, D. J., Durrington, P. N., Hitman, G. A., Neil, H. A. W., Livingstone, S. J., Thomason, M. J., Mackness, M. I., Charlton-Menys, V. & Fuller, J. H. 2004. Primary prevention of cardiovascular disease with atorvastatin in type 2 diabetes in the Collaborative Atorvastatin Diabetes Study (CARDS): Multicentre randomised placebo-controlled trial, *Lancet*, 364(9435), pp. 685–696. doi: 10.1016/S0140-6736(04)16895-5.
- Cramer, S. C. 2004. Functional imaging in stroke recovery, *Stroke*, 35(11 SUPPL. 1), pp. 2695–2698. doi: 10.1161/01.STR.0000143326.36847.b0.
- Cuspinada E, C. Machado, L. Galan, E. Aubert. 2007. QEEG Prognostic Value, 38(3).
- Cuspinada, E., Machado, C., Aubert, E., Galán, L., Liopis, F. & Avila, Y. 2003. Predicting Outcome in Acute Stroke: A Comparison between QEEG and the Canadian Neurological Scale, *Clinical EEG and Neuroscience*, 34(1), pp. 1–4. doi: 10.1177/155005940303400104.
- Diedler, J., Sykora, M., Jüttler, E., Veltkamp, R., Steiner, T. & Rupp, A. 2010 ‘EEG power spectrum to predict prognosis after hemicraniectomy for space-occupying middle cerebral artery infarction, *Cerebrovascular Diseases*, 29(2), pp. 162–169. doi: 10.1159/000262313.
- Dien & Doehner. 2008. Public Access’, *Bone*, 23(1), pp. 1–7. doi: 10.1038/jid.2014.371.
- Dimyan, M. A. & Cohen, L. G. 2011. Neuroplasticity in the context of motor rehabilitation after stroke’, *Nature Reviews Neurology*. Nature Publishing Group, 7(2), pp. 76–85. doi: 10.1038/nrneuro.2010.200.
- Doehner, W., Schenkel, J., Anker, S. D., Springer, J. & Audebert, H. 2013. Overweight and obesity are associated with improved survival, functional outcome, and stroke recurrence after acute stroke or transient ischaemic attack: Observations from the tempis trial. *European Heart Journal*, 34(4), pp. 268–277. doi: 10.1093/eurheartj/ehs340.
- Donnelly, E. M. & Blum, A. S. 2007. Focal and Generalized Slowing, Coma, and Brain Death. in Blum, A. S. and Rutkove, S. B. (eds) *The Clinical Neurophysiology Primer*. Totowa, NJ: Humana Press Inc, pp. 127–140.
- Dromerick, A. W., Edwards, D. F. & Dinger, M. N. 2003. Sensitivity to changes in disability after stroke: A comparison of four scales useful in clinical trials. *The Journal of Rehabilitation Research and Development*, 40(1), p. 1. doi: 10.1682/JRRD.2003.01.0001.
- Duffy, L., Gajree, S., Langhorne, P., Stott, D. J. & Quinn, T. J. 2013. Reliability (Inter-rater Agreement) of the barthel index for assessment of stroke survivors: Systematic review and meta-analysis, *Stroke*, 44(2), pp. 462–468. doi: 10.1161/STROKEAHA.112.678615.
- Dzialowski, I., Hill, M. D., Coutts, S. B., Demchuk, A. M., Kent, D. M., Wunderlich, O. & Von Kummer, R. 2006. Extent of early ischemic changes on computed tomography (CT) before thrombolysis: Prognostic value of the Alberta Stroke Program early CT score in ECASS II, *Stroke*, 37(4), pp. 973–978. doi: 10.1161/01.STR.0000206215.62441.56.
- Ernst, M., Boers, A. M. M., Aigner, A., Berkhemer, O. A., Yoo, A. J., Roos, Y. B., Dippel, D. W. J., Van Der Lugt, A., Van Oostenbrugge, R. J., Van Zwam,

- W. H., Fiehler, J., Marquering, H. A. & Majoie, C. B. L. M. 2017 . Association of Computed Tomography Ischemic Lesion Location with Functional Outcome in Acute Large Vessel Occlusion Ischemic Stroke. *Stroke*, 48(9), pp. 2426–2433. doi: 10.1161/STROKEAHA.117.017513.
- Ervasti, J., Virtanen, M., Lallukka, T., Friberg, E., Mittendorfer-Rutz, E., Lundström, E. & Alexanderson, K. 2017. Permanent work disability before and after ischaemic heart disease or stroke event: A nationwide population-based cohort study in Sweden, *BMJ Open*, 7(9). doi: 10.1136/bmjopen-2017-017910.
- Ezzat Nazzal, M., Ahmed Saadah, M., Mohammed Saadah, L. & Mustafa Trebinjac, S. 2009. Acute ischemic stroke: Relationship of brain lesion location & functional outcome, *Disability and Rehabilitation*, 31(18), pp. 1501–1506. doi: 10.1080/09638280802627702.
- Finnigan, S. P., Rose, S. E., Walsh, M., Griffin, M., Janke, A. L., McMahon, K. L., Gillies, R., Strudwick, M. W., Pettigrew, C. M., Semple, J., Brown, J., Brown, P. & Chalk, J. B. 2004. Correlation of Quantitative EEG in Acute Ischemic Stroke with 30-Day NIHSS Score: Comparison with Diffusion and Perfusion MRI, *Stroke*, 35(4), pp. 899–903. doi: 10.1161/01.STR.0000122622.73916.d2.
- Finnigan, S. & van Putten, M. J. A. M. 2013. EEG in ischaemic stroke: Quantitative EEG can uniquely inform (sub-)acute prognoses and clinical management. *Clinical Neurophysiology*. International Federation of Clinical Neurophysiology, 124(1), pp. 10–19. doi: 10.1016/j.clinph.2012.07.003.
- Finnigan, S., Wong, A. & Read, S. 2016. Defining abnormal slow EEG activity in acute ischaemic stroke: Delta/alpha ratio as an optimal QEEG index, *Clinical Neurophysiology*. International Federation of Clinical Neurophysiology, 127(2), pp. 1452–1459. doi: 10.1016/j.clinph.2015.07.014.
- Frost, S. B. 2003. Reorganization of Remote Cortical Regions After Ischemic Brain Injury: A Potential Substrate for Stroke Recovery, *Journal of Neurophysiology*, 89(6), pp. 3205–3214. doi: 10.1152/jn.01143.2002.
- Frykholm, P., 2002. Cerebral ischemia studied with PET and microdialysis. *Comprehensive Summaries Uppsala Dissertations* 91 : 5319 - 5328
- Goyal, N., Tsivgoulis, G., Iftikhar, S., Khorchid, Y., Fawad Ishfaq, M., Doss, V. T., Zand, R., Chang, J., Alsherbini, K., Choudhri, A., Hoit, D., Alexandrov, A. V., Arthur, A. S. & Eljovich, L. 2017. Admission systolic blood pressure and outcomes in large vessel occlusion strokes treated with endovascular treatment, *Journal of NeuroInterventional Surgery*, 9(5), pp. 451–454. doi: 10.1136/neurintsurg-2016-012386.
- Gray, V., Rice, C. L. & Garland, S. J. 2012. Factors that influence muscle weakness following stroke and their clinical implications: A critical review, *Physiotherapy Canada*, 64(4), pp. 415–426. doi: 10.3138/ptc.2011-03.
- Guo, X., Wu, W. & Tong, S. 2017. Asymmetry of hemispheric interdependences in the early hours following unilateral stroke: An electrophysiological study in rats, *Proceedings of the Annual International Conference of the IEEE Engineering in Medicine and Biology Society, EMBS*, pp. 4363–

4366. doi: 10.1109/EMBC.2017.8037822.

- Gupta, S., Yadav, R. & Malhotra, A. 2016. Assessment of physical disability using Barthel index among elderly of rural areas of district Jhansi (U.P), India, *Journal of Family Medicine and Primary Care*, 5(4), p. 853. doi: 10.4103/2249-4863.201178.
- Hara, Y. 2015. Brain Plasticity and Rehabilitation in Stroke Patients. *Journal of Nippon Medical School*, 82(1), pp. 4–13. doi: 10.1272/jnms.82.4.
- Harvey, R. L. 2015. Predictors of Functional Outcome Following Stroke. *Physical Medicine and Rehabilitation Clinics of North America*. Elsevier Inc, 26(4), pp. 583–598. doi: 10.1016/j.pmr.2015.07.002.
- Hatfield, L. A., Crone, N. E., Kossoff, E. H., Ewen, J. B., Pyzik, P. L., Lin, D. D. M., Kelley, T. M. & Comi, A. M. 2007. Quantitative EEG asymmetry correlates with clinical severity in unilateral Sturge-Weber syndrome, *Epilepsia*, 48(1), pp. 191–195. doi: 10.1111/j.1528-1167.2006.00630.x.
- Hauser, W. A. 1992. Seizure Disorders: The Changes With Age, *Epilepsia*, 33, pp. 6–14. doi: 10.1111/j.1528-1157.1992.tb06222.x.
- Heiss, W. D. & Kidwell, C. S. 2014. Imaging for prediction of functional outcome and assessment of recovery in ischemic stroke, *Stroke*, 45(4), pp. 1195–1201. doi: 10.1161/STROKEAHA.113.003611.
- Higashida, R. T., Furlan, A. J., Roberts, H., Tomsick, T., Connors, B., Barr, J., Dillon, W., Warach, S., Broderick, J., Tilley, B. & Sacks, D. 2003. Erratum: Trial Design and Reporting Standards for Intra-Arterial Cerebral Thrombolysis for Acute Ischemic Stroke (Stroke (August 2003) 34:8 (1156-1162))', *Stroke*, 34(11), p. 2774. doi: 10.1161/01.STR.0000101763.87465.A4.
- Huybrechts, K. F. & Caro, J. J. 2007. The Barthel Index and modified Rankin Scale as prognostic tools for long-term outcomes after stroke: A qualitative review of the literature, *Current Medical Research and Opinion*, 23(7), pp. 1627–1636. doi: 10.1185/030079907X210444.
- Jia, Q., Zhao, X., Wang, C., Wang, Yilong, Yan, Y., Li, H., Zhong, L., Liu, L., Zheng, H., Zhou, Y. & Wang, Yongjun. 2011. Diabetes and poor outcomes within 6 months after acute ischemic stroke: The China national stroke registry, *Stroke*, 42(10), pp. 2758–2762. doi: 10.1161/STROKEAHA.111.621649.
- Johnston, K. C., Bruno, A., Pauls, Q., Hall, C. E., Barrett, K. M., Barsan, W., Fansler, A., Van De Bruinhorst, K., Janis, S. & Durkalski-Mauldin, V. L. 2019. Intensive vs Standard Treatment of Hyperglycemia and Functional Outcome in Patients with Acute Ischemic Stroke: The SHINE Randomized Clinical Trial, *JAMA - Journal of the American Medical Association*, 322(4), pp. 326–335. doi: 10.1001/jama.2019.9346.
- Jönsson, A. C., Lindgren, I., Norrving, B. & Lindgren, A. 2008. Weight loss after stroke: A population-based study from the Lund Stroke Register. *Stroke*, 39(3), pp. 918–923. doi: 10.1161/STROKEAHA.107.497602.
- Katzan, I. L., Dawson, N. V., Thomas, C. L., Votruba, M. E. & Cebul, R. D. 2007. The cost of pneumonia after acute stroke, *Neurology*, 68(22), pp. 1938–1943. doi: 10.1212/01.wnl.0000263187.08969.45.

- Kelly, J., Hunt, B. J., Lewis, R. R., Swaminathan, R., Moody, A., Seed, P. T. & Rudd, A. 2004. Dehydration and venous thromboembolism after acute stroke. *QJM - Monthly Journal of the Association of Physicians*, 97(5), pp. 293–296. doi: 10.1093/qjmed/hch050.
- Khakshour, M. 2015. all things about QEEG (Rochester Institute of Technology). (August), pp. 0–37. doi: 10.13140/RG.2.1.2174.6403.
- Khalaf, G. M., Abdkadhim, M. & Suhail, I. M. 2018. Value of Alberta Stroke Program early Computed Tomography Score (ASPECTS) in the Evaluation of Middle Cerebral Artery Acute Ischemic Stroke using CT and DW-MRI, *Journal of Medical Diagnostic Methods*, 07(04), pp. 1–6. doi: 10.4172/2168-9784.1000280.
- Khan, M., Baird, G. L., Goddeau, R. P., Silver, B. & Henninger, N. 2017. Alberta stroke program early CT score infarct location predicts outcome following M2 occlusion, *Frontiers in Neurology*, 8(MAR), pp. 1–6. doi: 10.3389/fneur.2017.00098.
- Kirchhof, P. & Lambert, E. 2016. 2016 ESC Guidelines for the management of atrial fibrillation developed in collaboration with EACTS, *European Heart Journal*, 37(38). doi: 10.1093/eurheartj/ehw210.
- van der Knaap, L. J. & van der Ham, I. J. M. 2011. How does the corpus callosum mediate interhemispheric transfer? A review. *Behavioural Brain Research*. Elsevier B.V., 223(1), pp. 211–221. doi: 10.1016/j.bbr.2011.04.018.
- Kongsawasdi, S., Klaphajone, J., Wivatvongvana, P. & Watcharasaksilp, K. 2019. Prognostic Factors of Functional Outcome Assessed by Using the Modified Rankin Scale in Subacute Ischemic Stroke, *Journal of Clinical Medicine Research*, 11(5), pp. 375–382. doi: 10.14740/jocmr3799.
- Koton, S., Molshatzki, N., Bornstein, N. M. & Tanne, D. 2012. Low cholesterol, statins and outcomes in patients with first-ever acute ischemic stroke, *Cerebrovascular Diseases*, 34(3), pp. 213–220. doi: 10.1159/000342302.
- Leon-Carrion, J., Martin-Rodriguez, J. F., Damas-Lopez, J., Barroso y Martin, J. M. & Dominguez-Morales, M. R. 2009. Delta-alpha ratio correlates with level of recovery after neurorehabilitation in patients with acquired brain injury, *Clinical Neurophysiology*. International Federation of Clinical Neurophysiology, 120(6), pp. 1039–1045. doi: 10.1016/j.clinph.2009.01.021.
- Li, J., Zhang, P., Wu, S., Yi, X., Wang, C. & Liu, M. 2018. Factors associated with favourable outcome in large hemispheric infarctions, *BMC Neurology*. BMC Neurology, 18(1), pp. 1–8. doi: 10.1186/s12883-018-1148-7.
- Li, S. 2017. Spasticity, motor recovery, & neural plasticity after stroke. *Frontiers in Neurology*, 8(APR), pp. 1–8. doi: 10.3389/fneur.2017.00120.
- Liu-ambrose, T., Eng, J. J. & Ot, P. T. 2015. Exercise Training and Recreational Activities to Promote Executive Functions in Chronic Stroke : A Proof-of-concept Study, *Journal of Stroke and Cerebrovascular Diseases*. Elsevier Ltd, 24(1), pp. 130–137. doi: 10.1016/j.jstrokecerebrovasdis.2014.08.012.
- Logar, C. & Boswell, M. 1991. The value of EEG-Mapping in focal cerebral lesions, *Brain Topography*, 3(4), pp. 441–446. doi: 10.1007/BF01129003.
- Lu, Q., Huang, G., Chen, L., Li, W. & Liang, Z. 2019. Structural and functional

- reorganization following unilateral internal capsule infarction contribute to neurological function recovery, *Neuroradiology*. *Neuroradiology*, 61(10), pp. 1181–1190. doi: 10.1007/s00234-019-02278-x.
- Luo, H., Wang, X., Chen, C., Wang, J., Li, C., Xu, Z., Yang, X., Shi, W. & Zeng, C. 2015. Oxidative stress causes imbalance of renal renin angiotensin system (RAS) components and hypertension in obese Zucker rats, *Journal of the American Heart Association*, 4(2), pp. 1–14. doi: 10.1161/JAHA.114.001559.
- Mahmoud Fouad, M., Mohamed Farag, S., Hegazy, M. I. & Abd Elalem Aziz, M. 2017. Prediction of Functional Outcome in Ischemic Stroke Patients: An Observational Study on Egyptian Population, *Cureus*, 9(6). doi: 10.7759/cureus.1392.
- Mang, C. S., Campbell, K. L., Ross, C. J. D. & Boyd, L. A. 2013. Promoting Neuroplasticity for Motor Rehabilitation After Stroke: Considering the Effects of Aerobic Exercise & Genetic Variation on Brain-Derived Neurotrophic Factor, *Physical Therapy*, 93(12), pp. 1707–1716. doi: 10.2522/ptj.20130053.
- Martin-schild, S., Albright, K. C., Mph, D. O., Tanksley, J., Jones, E. B., Grotta, J. C. & Savitz, S. I. 2012. NIH Public Access, 57(February 2010), pp. 42–45. doi: 10.1016/j.annemergmed.2010.06.564.Zero.
- van Meer, M. P. A., Otte, W. M., van der Marel, K., Nijboer, C. H., Kavelaars, A., van der Sprenkel, J. W. B., Viergever, M. A. & Dijkhuizen, R. M. 2012. Extent of bilateral neuronal network reorganization and functional recovery in relation to stroke severity', *Journal of Neuroscience*, 32(13), pp. 4495–4507. doi: 10.1523/JNEUROSCI.3662-11.2012.
- Mehta, B., Leslie-Mazwi, T. M., Chandra, R. V., Chaudhry, Z. A., Rabinov, J. D., Hirsch, J. A., Schwamm, L. H., Rost, N. S. & Yoo, A. J. 2013. Assessing variability in neurointerventional practice patterns for acute ischemic stroke, *Journal of NeuroInterventional Surgery*, 5(SUPPL.1), pp. 4–6. doi: 10.1136/neurintsurg-2012-010565.
- Melikian, N., Seddon, M. D., Casadei, B., Chowienzyk, P. J. & Shah, A. M. 2009. Neuronal nitric oxide synthase and human vascular regulation, *Trends in Cardiovascular Medicine*. Elsevier Inc., 19(8), pp. 256–262. doi: 10.1016/j.tcm.2010.02.007.
- Miller, C. M. & Behrouz, R. 2016. Impact of Infection on Stroke Morbidity and Outcomes, *Current Neurology and Neuroscience Reports*. *Current Neurology and Neuroscience Reports*, 16(9), pp. 1–5. doi: 10.1007/s11910-016-0679-9.
- Mori, M., Naganuma, M., Okada, Y., Hasegawa, Y., Shiokawa, Y., Nakagawara, J., Furui, E., Kimura, K., Yamagami, H., Kario, K., Okuda, S., Koga, M., Minematsu, K. & Toyoda, K. 2012. Early neurological deterioration within 24 hours after intravenous rt-PA therapy for stroke patients: The stroke acute management with urgent risk factor assessment and improvement rt-PA registry. *Cerebrovascular Diseases*, 34(2), pp. 140–146. doi: 10.1159/000339759.
- Murphy, T. H. & Corbett, D. 2009. Plasticity during stroke recovery: From synapse

- to behaviour. *Nature Reviews Neuroscience*. Nature Publishing Group, 10(12), pp. 861–872. doi: 10.1038/nrn2735.
- Murri, L., Gori, S., Massetani, R., Bonanni, E., Marcella, F. & Milani, S. 1998. Evaluation of acute ischemic stroke using quantitative EEG: A comparison with conventional EEG and CT scan. *Neurophysiologie Clinique*, 28(3), pp. 249–257. doi: 10.1016/S0987-7053(98)80115-9.
- Musa, K. I. & Keegan, T. J. 2018. The change of Barthel Index scores from the time of discharge until 3-month post-discharge among acute stroke patients in Malaysia: A random intercept model. *PLoS ONE*, 13(12), pp. 1–13. doi: 10.1371/journal.pone.0208594.
- Muscari, A., Puddu, G. M., Santoro, N. & Zoli, M. 2011. A simple scoring system for outcome prediction of ischemic stroke. *Acta Neurologica Scandinavica*, 124(5), pp. 334–342. doi: 10.1111/j.1600-0404.2010.01479.x.
- Naito, H., Nezu, T., Hosomi, N., Aoki, S., Kinoshita, N., Kuga, J., Shimomura, R., Araki, M., Ueno, H., Ochi, K. & Maruyama, H. 2018. Controlling Nutritional Status score for predicting 3-mo functional outcome in acute ischemic stroke. *Nutrition*. Elsevier Inc., 55–56, pp. 1–6. doi: 10.1016/j.nut.2018.02.025.
- Nakayama, E., Tohara, H., Hino, T., Sato, M., Hiraba, H., Abe, K. & Ueda, K. . 2014. The effects of ADL on recovery of swallowing function in stroke patients after acute phase. *Journal of Oral Rehabilitation*, 41(12), pp. 904–911. doi: 10.1111/joor.12212.
- Ntaios, G., Gioulekas, F., Papavasileiou, V., Strbian, D. & Michel, P. 2016. ASTRAL, DRAGON and SEDAN scores predict stroke outcome more accurately than physicians. *European Journal of Neurology*, 23(11), pp. 1651–1657. doi: 10.1111/ene.13100.
- Nudo, R. J. 2013. Recovery after brain injury: mechanisms and principles. *Frontiers in Human Neuroscience*, 7(December), pp. 1–14. doi: 10.3389/fnhum.2013.00887.
- Pan, S. L., Lien, I. N. & Chen, T. H. H. 2010. Is Higher Serum Total Cholesterol Level Associated With Better Long-Term Functional Outcomes After Noncardioembolic Ischemic Stroke?. *Archives of Physical Medicine and Rehabilitation*. Elsevier Inc., 91(6), pp. 913–918. doi: 10.1016/j.apmr.2010.02.002.
- Paolucci, S., Antonucci, G., Grasso, M. G. & Pizzamiglio, L. 2001. The role of unilateral spatial neglect in rehabilitation of right brain-damaged ischemic stroke patients: A matched comparison. *Archives of Physical Medicine and Rehabilitation*, 82(6), pp. 743–749. doi: 10.1053/apmr.2001.23191.
- Park, J. S., Lee, J. M., Kwak, H. S. & Chung, G. H. 2019. Predictive value of CT angiography source image ASPECTS in patients with anterior circulation acute ischemic stroke after endovascular treatment: Ultimate infarct size and clinical outcome. *Journal of NeuroInterventional Surgery*, 11(4), pp. 342–346. doi: 10.1136/neurintsurg-2018-014359.
- Parsons, M. W., Barber, P. A., Desmond, P. M., Baird, T. A., Darby, D. G., Byrnes, G., Tress, B. M. & Davis, S. M. 2002. Acute hyperglycemia adversely

- affects stroke outcome: A magnetic resonance imaging and spectroscopy study. *Annals of Neurology*, 52(1), pp. 20–28. doi: 10.1002/ana.10241.
- Penry, K. 1981. Proposal for Revised Clinical and Electroencephalographic Classification of Epileptic Seizures.. *Epilepsia*, 22(4), pp. 489–501. doi: 10.1111/j.1528-1157.1981.tb06159.x.
- van Putten, M. J. A. M. 2007. The revised brain symmetry index. *Clinical Neurophysiology*, 118(11), pp. 2362–2367. doi: 10.1016/j.clinph.2007.07.019.
- Van Putten, M. J. A. M. & Tavy, D. L. J. 2004. Continuous quantitative EEG monitoring in hemispheric stroke patients using the brain symmetry index. *Stroke*, 35(11), pp. 2489–2492. doi: 10.1161/01.STR.0000144649.49861.1d.
- Quinn, T. J., Langhorne, P. & Stott, D. J. 2011. Barthel index for stroke trials: Development, properties, and application. *Stroke*, 42(4), pp. 1146–1151. doi: 10.1161/STROKEAHA.110.598540.
- Ribo, M., Molina, C. A., Delgado, P., Rubiera, M., Delgado-Mederos, R., Rovira, A., Munuera, J. & Alvarez-Sabin, J. 2007. Hyperglycemia during ischemia rapidly accelerates brain damage in stroke patients treated with tPA, *Journal of Cerebral Blood Flow and Metabolism*, 27(9), pp. 1616–1622. doi: 10.1038/sj.jcbfm.9600460.
- Rosso, C., Colliot, O., Pires, C., Delmaire, C., Valabrègue, R., Crozier, S., Dormont, D., Baillet, S., Samson, Y. & Lehericy, S. 2011. Prédiction du handicap par les valeurs d'ADC dans les structures motrices à la phase hyper-aiguë de l'AVC ischémique étude comparative avec le volume en diffusion. *Journal of Neuroradiology*. Elsevier Masson SAS, 38(2), pp. 105–112. doi: 10.1016/j.neurad.2010.05.001.
- Ryu, W. S., Schellingerhout, D., Jeong, S. W., Nahrendorf, M. & Kim, D. E. 2016. Association between Serum Lipid Profiles and Early Neurological Deterioration in Acute Ischemic Stroke. *Journal of Stroke and Cerebrovascular Diseases*. Elsevier Inc., 25(8), pp. 2024–2030. doi: 10.1016/j.jstrokecerebrovasdis.2016.05.009.
- Saif, S. & Fazal, N. 2014. Association between NHI stroke scale score and functional outcome in acute ischemic stroke. *NIH Stroke Scale Score*, 64(4), pp. 585–590.
- Sajobi, T. T. & Timothy, I.P. 2017. Early trajectory of stroke severity predicts long-term functional outcomes in ischemic stroke subjects. *Stroke*, 48(1), pp. 105–110. doi: 10.1161/STROKEAHA.116.014456.
- Salinsky, M. C., Oken, B. S., Kramer, R. E. & Morehead, L. 1992. A comparison of quantitative EEG frequency analysis and conventional EEG in patients with focal brain lesions. *Electroencephalography and Clinical Neurophysiology*, 83(6), pp. 358–366. doi: 10.1016/0013-4694(92)90071-O.
- Salter, K., Campbell, N., Richardson, M., Mehta, S., Jutai, J., Zettler, L., Ba, M. M., McClure, A., Mays, R., Foley, N. & Teasell, R. 2013. EBRSR [Evidence-Based Review of Stroke Rehabilitation] 20 Outcome Measures in Stroke Rehabilitation. *Evidence-Based Review of Stroke Rehabilitation*,

- p. 141. Available at: <http://www.ebrsr.com/evidence-review>.
- Santalucia, P., Pezzella, F. R., Sessa, M., Monaco, S., Torgano, G., Anticoli, S., Zanolli, E., Maimone Baronello, M., Paciaroni, M. & Caso, V. 2013. Sex differences in clinical presentation, severity and outcome of stroke: Results from a hospital-based registry. *European Journal of Internal Medicine*. European Federation of Internal Medicine., 24(2), pp. 167–171. doi: 10.1016/j.ejim.2012.10.004.
- Sato, M., Ido, Y., Yoshimura, Y. & Mutai, H. 2019. Relationship of Malnutrition During Hospitalization With Functional Recovery and Postdischarge Destination in Elderly Stroke Patients. *Journal of Stroke and Cerebrovascular Diseases*. Elsevier Inc., 28(7), pp. 1866–1872. doi: 10.1016/j.jstrokecerebrovasdis.2019.04.012.
- Scherbakov, N., Von Haehling, S., Anker, S. D., Dirnagl, U. & Doehner, W. 2013. Stroke induced Sarcopenia: Muscle wasting and disability after stroke. *International Journal of Cardiology*. Elsevier Ireland Ltd, 170(2), pp. 89–94. doi: 10.1016/j.ijcard.2013.10.031.
- Scherbakov, N., Dirnagl, U. & Doehner, W. 2011. Body weight after stroke lessons from the obesity paradox. *Stroke*, 42(12), pp. 3646–3650. doi: 10.1161/STROKEAHA.111.619163.
- Scherbakov, N. & Doehner, W. 2011. Sarcopenia in stroke-facts and numbers on muscle loss accounting for disability after stroke. *Journal of Cachexia, Sarcopenia and Muscle*, 2(1), pp. 5–8. doi: 10.1007/s13539-011-0024-8.
- Setyopranoto, I., Wibowo, S., Sadewa, A. & Lamsudin, R. 2016. Reduced Vascular Endothelial Growth Factor A Levels are Correlated with the Incidence of Brain Edema in Acute Ischemic Stroke Patients. *British Journal of Medicine and Medical Research*, 13(2), pp. 1–7. doi: 10.9734/BJMMR/2016/23156.
- Sharanreddy, M. & Kulkarni, P. K. 2013. Automated EEG signal analysis for identification of epilepsy seizures and brain tumour. *Journal of Medical Engineering and Technology*, 37(8), pp. 511–519. doi: 10.3109/03091902.2013.837530.
- Sheorajpanday, R. V. A., Nagels, G., Weeren, A. J. T. M., Surgeloose, D. De & Deyn, P. P. De. 2010. Clinical Neurophysiology Additional value of quantitative EEG in acute anterior circulation syndrome of presumed ischemic origin. *Clinical Neurophysiology*. International Federation of Clinical Neurophysiology, 121(10), pp. 1719–1725. doi: 10.1016/j.clinph.2009.10.037.
- Sheorajpanday, R. V. A., Nagels, G., Weeren, A. J. T. M. & Deyn, P. P. De. 2011. Clinical Neurophysiology Quantitative EEG in ischemic stroke: Correlation with infarct volume and functional status in posterior circulation and lacunar syndromes. *Clinical Neurophysiology*. International Federation of Clinical Neurophysiology, 122(5), pp. 884–890. doi: 10.1016/j.clinph.2010.08.020.
- Sheorajpanday, R. V. A., Nagels, G., Weeren, A. J. T. M., van Putten, M. J. A. M. & De Deyn, P. P. 2011. Quantitative EEG in ischemic stroke: Correlation with functional status after 6 months. *Clinical Neurophysiology*.

- International Federation of Clinical Neurophysiology, 122(5), pp. 874–883. doi: 10.1016/j.clinph.2010.07.028.
- Shimoyama, T., Kimura, K., Uemura, J., Saji, N. & Shibazaki, K. 2014. Elevated glucose level adversely affects infarct volume growth and neurological deterioration in non-diabetic stroke patients, but not diabetic stroke patients. *European Journal of Neurology*, 21(3), pp. 402–410. doi: 10.1111/ene.12280.
- Slot, K. B., Berge, E., Sandercock, P., Lewis, S. C., Dorman, P. & Dennis, M. 2009. Causes of death by level of dependency at 6 months after ischemic stroke in 3 large cohorts. *Stroke*, 40(5), pp. 1585–1589. doi: 10.1161/STROKEAHA.108.531533.
- Song, J., Davey, C., Poulsen, C., Luu, P., Turovets, S., Anderson, E., Li, K. & Tucker, D. 2015. EEG source localization: Sensor density and head surface coverage. *Journal of Neuroscience Methods*. Elsevier B.V., 256, pp. 9–21. doi: 10.1016/j.jneumeth.2015.08.015.
- Strong, K., Mathers, C. & Bonita, R. 2007. Preventing stroke : saving lives around the world’, 6(February), pp. 182–187.
- Syeban, Z., Octaviana, F. 2014. EEG Praktis. *Badan Penerbit Fakultas Kedokteran Universitas Indonesia*. hal 1-17
- Takeuchi, N., Oouchida, Y. & Izumi, S. I. 2012. Motor control and neural plasticity through interhemispheric interactions. *Neural Plasticity*, 2012. doi: 10.1155/2012/823285.
- Thorpe, K. E. (2013) ‘outcomes’.
- Toschke, A. M., Tilling, K., Cox, A. M., Rudd, A. G., Heuschmann, P. U. & Wolfe, C. D. A. 2010. Patient-specific recovery patterns over time measured by dependence in activities of daily living after stroke and post-stroke care : The South London Stroke Register (SLSR)’, pp. 219–225. doi: 10.1111/j.1468-1331.2009.02774.x.
- Tunjungsari, A.P., Asmedi, A. Paryono. 2018. Abnormalitas Gelombang Elektroensefalografi sebagai Prediktor Perburukan Defisit Neurologis Pada Stroke Iskemik Akut.
- Tsai, Y. H., Yuan, R., Huang, Y. C., Yeh, M. Y., Lin, C. P. & Biswal, B. B. 2014. Disruption of brain connectivity in acute stroke patients with early impairment in consciousness. *Frontiers in Psychology*, 4(JAN), pp. 1–10. doi: 10.3389/fpsyg.2013.00956.
- Tsao, J. W., Hemphill, J. C., Johnston, S. C., Smith, W. S. & Bonovich, D. C. 2005. Initial Glasgow Coma Scale score predicts outcome following thrombolysis for posterior circulation stroke. *Archives of Neurology*, 62(7), pp. 1126–1129. doi: 10.1001/archneur.62.7.1126.
- Tsivgoulis, G., Kotsis, V. & Giannopoulos, S. 2011. Intravenous thrombolysis for acute ischaemic stroke: Effective blood pressure control matters’, *International Journal of Stroke*, 6(2), pp. 125–127. doi: 10.1111/j.1747-4949.2010.00570.x.
- Tsolaki, A., Kazis, D., Kompatsiaris, I., Kosmidou, V. & Tsolaki, M. 2014. Electroencephalogram and alzheimer’s disease: Clinical and research approaches. *International Journal of Alzheimer’s Disease*. Hindawi

- Publishing Corporation, 2014. doi: 10.1155/2014/349249.
- Ultra1, P. 2001. qEEG mechanism of , 117(September), pp. 1–9. doi: 10.1111/j.1747-4949.2009.00337.x.
- Vemmos, K. N., Tsivgoulis, G., Spengos, K., Zakopoulos, N., Synetos, A., Kotsis, V., Vassilopoulos, D. & Mavrikakis, M. 2003. Association between 24-h blood pressure monitoring variables and brain oedema in patients with hyperacute stroke. *Journal of Hypertension*, 21(11), pp. 2167–2173. doi: 10.1097/00004872-200311000-00027.
- Wang, H., Camicia, M., Terdiman, J., Hung, Y.Y., Sandel, M.E., 2011. Time to inpatient rehabilitation hospital admission and functional outcomes of stroke patients. *PMR* 3, 296-304.
- Ward, N. 2017. 20 Neural plasticity for functional rehabilitation after stroke. *Journal of Neurology, Neurosurgery & Psychiatry*, 88(8), p. A10.2-A10. doi: 10.1136/jnnp-2017-BNPA.20.
- Ward, N. S. & Cohen, L. G. 2004. Europe PMC Funders Group Mechanisms Underlying Recovery of Motor Function After Stroke. *Arch. Neurol.*, 61(12), pp. 1844–1848. doi: 10.1001/archneur.61.12.1844.Mechanisms.
- Winstein, C. J. & Wu. L.M. 2016. *AHA / ASA Guideline Guidelines for Adult Stroke Rehabilitation and Recovery*. doi: 10.1161/STR.0000000000000098.
- De Wit, L., Putman, K., Devos, H., Brinkmann, N., Dejaeger, E., De Weerd, W., Jenni, W., Lincoln, N., Schuback, B. & Schupp, W. 2014. Long-term prediction of functional outcome after stroke using single items of the Barthel Index at discharge from rehabilitation centre. *Disability and Rehabilitation*, 36(5), pp. 353–358. doi: 10.3109/09638288.2013.793411.
- Wu, J., Srinivasan, R., Burke Quinlan, E., Solodkin, A., Small, S. L. & Cramer, S. C. 2016. Utility of EEG measures of brain function in patients with acute stroke. *Journal of Neurophysiology*, 115(5), pp. 2399–2405. doi: 10.1152/jn.00978.2015.
- Yagi, M., Yasunaga, H., Matsui, H., Morita, K., Fushimi, K., Fujimoto, M., Koyama, T. & Fujitani, J. 2017. Impact of Rehabilitation on Outcomes in Patients with Ischemic Stroke: A Nationwide Retrospective Cohort Study in Japan', *Stroke*, 48(3), pp. 740–746. doi: 10.1161/STROKEAHA.116.015147.
- Yoo, A. J., Zaidat, O. O., Chaudhry, Z. A., Berkhemer, O. A., Gilberto Gonzalez, R., Goyal, M., Demchuk, A. M., Menon, B. K., Mualem, E., Ueda, D., Buell, H., Sit, S. P. & Bose, A. 2014. Impact of pretreatment noncontrast CT Alberta stroke program early ct score on clinical outcome after intra arterial stroke therapy', *Stroke*, 45(3), pp. 746–751. doi: 10.1161/STROKEAHA.113.004260.
- Yu, A. Y. X., Holodinsky, J. K., Zerna, C., Svenson, L. W., Jetté, N., Quan, H. & Hill, M. D. 2016. Use and utility of administrative health data for stroke research and surveillance', *Stroke*, 47(7), pp. 1946–1952. doi: 10.1161/STROKEAHA.116.012390.
- Zhao, Y., Zou, C., Wang, C., Zhang, Y. & Wang, S. 2016. Long-term outcomes after stroke in elderly patients with atrial fibrillation: A hospital-based

follow-up study in China', *Frontiers in Aging Neuroscience*, 8(MAR), pp.
1–6. doi: 10.3389/fnagi.2016.00056.