UNIVERSITAS
GADJAH MADA
PARAMETER KUANTITATIF EEG (QEEG) SEBAGAI PREDIKTOR LUARAN FUNGSIONAL JANGKA PANJANG PASIEN STROKE ISKEMIK
DYANNE PARAMITA A P, dr. Astuti, Sp.S (K).; Dr. dr. Ahmad Asmedi, M.Kes, Sp.S (K)
Universitas Gadjah Mada, 2019 | Diunduh dari http://etd.repository.ugm.ac.id/

DAFTAR PUSTAKA

Access, O., Nardone, R., Brigo, F. & Trinka, E. 2016. Acute symptomatics seizures caused by electrolyte disturbance, 12(1), pp. 21–33. doi: 10.3988/jcn.2016.12.1.21.

Agius Anastasi, A., Falzon, O., Camilleri, K., Vella, M. & Muscat, R. 2017. Brain symmetry index in healthy and stroke patients for assessment and prognosis', *Stroke Research and Treatment*, 2017. doi: 10.1155/2017/8276136.

Alia, C., Spalletti, C., Lai, S., Panarese, A., Lamola, G., Bertolucci, F., Vallone, F., Di Garbo, A., Chisari, C., Micera, S. & Caleo, M. 2017. Neuroplastic Changes Following Brain Ischemia and their Contribution to Stroke Recovery: Novel Approaches in Neurorehabilitation, *Frontiers in Cellular Neuroscience*, 11(March), pp. 1–22. doi: 10.3389/fncel.2017.00076.

Alvarez-Sabín, J., Montaner, J., Padró, L., Molina, C. A., Rovira, R., Codina, A. & Quintana, M. 2002. Gabapentin in late-onset poststroke seizures', *Neurology*, 59(12), pp. 1991–1993. doi: 10.1212/01.WNL.0000038388.57824.B6.

Appelros, P., Nydevik, I. & Viitanen, M. 2003. Poor outcome after first-ever stroke: Predictors for death, dependency, and recurrent stroke within the first year, *Stroke*, 34(1), pp. 122–126. doi: 10.1161/01.STR.0000047852.05842.3C.

Assenza, G., Zappasodi, F., Pasqualetti, P., Vernieri, F. & Tecchio, F. 2013. A contralesional EEG power increase mediated by interhemispheric disconnection provides negative prognosis in acute stroke, *Restorative Neurology and Neuroscience*, 31(2), pp. 177–188. doi: 10.3233/RNN-120244.

Ayerbe, L., Ayis, S., Crichton, S., Wolfe, C. D. A. & Rudd, A. G. 2014. The long-term outcomes of depression up to 10 years after stroke; The south london stroke rregister, *Journal of Neurology, Neurosurgery and Psychiatry*, 85(5), pp. 514–521. doi: 10.1136/jnnp-2013-306448.

Bauer, G., Bauer, R. 2016. EEG, Drug Effects, and Central Nervous System Poisoning. Available at www.neuropsykey.com

Beckham, J. D. & Tyler, K. L. 2012. Neuro-Intensive Care of Patients with Acute CNS Infections, *Neurotherapeutics*, 9, pp. 124–138. doi: 10.1007/s13311-011-0086-5.

Beghi, E. Alessandro, R. & Beretta, S. 2011. Incidence and predictors of acute symptomatic seizures after stroke, *Neurology*, 77(20), pp. 1785–1793. doi: 10.1212/WNL.0b013e3182364878.

Bentes, C., Peralta, A. R., Viana, P., Martins, H., Morgado, C., Casimiro, C., Franco, A. C., Fonseca, A. C., Geraldes, R., Canhão, P., Pinho e Melo, T., Paiva, T. & Ferro, J. M. 2018. Quantitative EEG and functional outcome following acute ischemic stroke, *Clinical Neurophysiology*. International Federation of Clinical Neurophysiology, 129(8), pp. 1680–1687. doi: 10.1016/j.clinph.2018.05.021.

Between, R., Deficit, N., Functional, F., Shifts, O., First, D. & After, H. 2013. Relation Between Neurologic Deficit Severity, 43(6), pp. 1537–1541. doi:

UNIVERSITAS GADJAH MADA
PARAMETER KUANTITATIF EEG (QEEG) SEBAGAI PREDIKTOR LUARAN FUNGSIONAL JANGKA PANJANG PASIEN STROKE ISKEMIK
DYANNE PARAMITA A P, dr. Astuti, Sp.S (K).; Dr. dr. Ahmad Asmedi, M.Kes, Sp.S (K)
Universitas Gadjah Mada, 2019 | Diunduh dari http://etd.repository.ugm.ac.id/

10.1161/STROKEAHA.111.636928.RELATION.

Bhardwaj, A. 2006. Neurological impact of vasopressin dysregulation and hyponatremia, *Annals of Neurology*, 59(2), pp. 229–236. doi: 10.1002/ana.20788.

Bhattarai, S., Ning, Z.X. 2014. EEG and SPECT Change in Acute Ischemic Stroke. *J Neurol Neurophysiol* 5: 190

Buckingham, S. C., Campbell, S. L., Haas, B. R., Montana, V., Robel, S., Ogunrinu, T. & Sontheimer, H. 2011. Glutamate release by primary brain tumors induces epileptic activity, *Nature Medicine*. Nature Publishing Group, 17(10), pp. 1269–1274. doi: 10.1038/nm.2453.

Bureau, M., Azkoitia, J. M., Ezmendi, G., Manterola, I., Zabaleta, H., Perez, M. & Medina, J. 2007. Non-invasive, wireless and universal interface for the control of peripheral devices by means of head movements, *2007 IEEE 10th International Conference on Rehabilitation Robotics, ICORR'07*, 4(10), pp. 124–131. doi: 10.1109/ICORR.2007.4428417.

Burneo, J. G., Fang, J. and Saposnik, G. 2010. Impact of seizures on morbidity and mortality after stroke: A Canadian multi-centre cohort study, *European Journal of Neurology*, 17(1), pp. 52–58. doi: 10.1111/j.1468-1331.2009.02739.x.

Capes, S. E., Hunt, D., Malmberg, K., Pathak, P. & Gerstein, H. C. 2001. Stress hyperglycemia and prognosis of stroke in nondiabetic and diabetic patients: A systematic overview, *Stroke*, 32(10), pp. 2426–2432. doi: 10.1161/hs1001.096194.

Castillo, J., Leira, R., García, M. M., Serena, J., Blanco, M. & Dávalos, A. 2004. Blood Pressure Decrease during the Acute Phase of Ischemic Stroke Is Associated with Brain Injury and Poor Stroke Outcome, *Stroke*, 35(2), pp. 520–527. doi: 10.1161/01.STR.0000109769.22917.B0.

Cernik, D. Sanak, D. & Divisova, P. 2019. Impact of blood pressure levels within first 24 hours after mechanical thrombectomy on clinical outcome in acute ischemic stroke patients, *Journal of NeuroInterventional Surgery*, 11(8), pp. 735–739. doi: 10.1136/neurintsurg-2018-014548.

Chen, W., Liu, G., Fang, J., Wang, Y., Song, Y., Pan, Y., Li, H.,& Liu, L. 2016. External Validation of the Totaled Health Risks in Vascular Events Score to Predict Functional Outcome and Mortality in Patients Entered into the China National Stroke Registry. *J. StrokeCerebrovasc. Dis.* 25.2331-2337.

Cheng, B. Forkert, N.D, & Zavaglia, M. 2014. Influence of stroke infarct location on functional outcome measured by the modified rankin scale, *Stroke*, 45(6), pp. 1695–1702. doi: 10.1161/STROKEAHA.114.005152.

Choi-Kwon, S., Han, S. W., Kwon, S. U. & Kim, J. S. 2005. Poststroke fatigue: Characteristics and related factors, *Cerebrovascular Diseases*, 19(2), pp. 84–90. doi: 10.1159/000082784.

Cioncoloni, D., Piu, P., Tassi, R., Acampa, M., Guideri, F., Taddei, S., Bielli, S., Martini, G. & Mazzocchio, R. 2012. Relationship between the modified Rankin Scale and the Barthel Index in the process of functional recovery after stroke, *NeuroRehabilitation*, 30(4), pp. 315–322. doi: 10.3233/NRE-2012-0761.

UNIVERSITAS
GADJAH MADA
PARAMETER KUANTITATIF EEG (QEEG) SEBAGAI PREDIKTOR LUARAN FUNGSIONAL JANGKA PANJANG PASIEN STROKE ISKEMIK
DYANNE PARAMITA A P, dr. Astuti, Sp.S (K).; Dr. dr. Ahmad Asmedi, M.Kes, Sp.S (K)
Universitas Gadjah Mada, 2019 | Diunduh dari http://etd.repository.ugm.ac.id/

Colhoun, H. M., Betteridge, D. J., Durrington, P. N., Hitman, G. A., Neil, H. A. W., Livingstone, S. J., Thomason, M. J., Mackness, M. I., Charlton-Menys, V. & Fuller, J. H. 2004. Primary prevention of cardiovascular disease with atorvastatin in type 2 diabetes in the Collaborative Atorvastatin Diabetes Study (CARDS): Multicentre randomised placebo-controlled trial, *Lancet*, 364(9435), pp. 685–696. doi: 10.1016/S0140-6736(04)16895-5.

Cramer, S. C. 2004. Functional imaging in stroke recovery, *Stroke*, 35(11 SUPPL. 1), pp. 2695–2698. doi: 10.1161/01.STR.0000143326.36847.b0.

Cuspineda E, C. Machado, L. Galan, E. Aubert. 2007. QEEG Prognostic Value, 38(3).

Cuspineda, E., Machado, C., Aubert, E., Galán, L., Liopis, F. & Avila, Y. 2003. Predicting Outcome in Acute Stroke: A Comparison between QEEG and the Canadian Neurological Scale, *Clinical EEG and Neuroscience*, 34(1), pp. 1–4. doi: 10.1177/155005940303400104.

Diedler, J., Sykora, M., Jüttler, E., Veltkamp, R., Steiner, T. & Rupp, A. 2010 'EEG power spectrum to predict prognosis after hemicraniectomy for space-occupying middle cerebral artery infarction, *Cerebrovascular Diseases*, 29(2), pp. 162–169. doi: 10.1159/000262313.

Dien & Doehner. 2008. Public Access', *Bone*, 23(1), pp. 1–7. doi: 10.1038/jid.2014.371.

Dimyan, M. A. & Cohen, L. G. 2011. Neuroplasticity in the context of motor rehabilitation after stroke', *Nature Reviews Neurology*. Nature Publishing Group, 7(2), pp. 76–85. doi: 10.1038/nrneurol.2010.200.

Doehner, W., Schenkel, J., Anker, S. D., Springer, J. & Audebert, H. 2013. Overweight and obesity are associated with improved survival, functional outcome, and stroke recurrence after acute stroke or transient ischaemic attack: Observations from the tempis trial. *European Heart Journal*, 34(4), pp. 268–277. doi: 10.1093/eurheartj/ehs340.

Donnelly, E. M. & Blum, A. S. 2007. Focal and Generalized Slowing, Coma, and Brain Death. in Blum, A. S. and Rutkove, S. B. (eds) *The Clinical Neurophysiology Primer*. Totowa, NJ: Humana Press Inc, pp. 127–140.

Dromerick, A. W., Edwards, D. F.& Diringer, M. N. 2003. Sensitivity to changes in disability after stroke: A comparison of four scales useful in clinical trials. *The Journal of Rehabilitation Research and Development*, 40(1), p. 1. doi: 10.1682/JRRD.2003.01.0001.

Duffy, L., Gajree, S., Langhorne, P., Stott, D. J.& Quinn, T. J. 2013. Reliability (Inter-rater Agreement) of the barthel index for assessment of stroke survivors: Systematic review and meta-analysis, *Stroke*, 44(2), pp. 462–468. doi: 10.1161/STROKEAHA.112.678615.

Dzialowski, I., Hill, M. D., Coutts, S. B., Demchuk, A. M., Kent, D. M., Wunderlich, O. & Von Kummer, R. 2006. Extent of early ischemic changes on computed tomography (CT) before thrombolysis: Prognostic value of the Alberta Stroke Program early CT score in ECASS II, *Stroke*, 37(4), pp. 973–978. doi: 10.1161/01.STR.0000206215.62441.56.

Ernst, M., Boers, A. M. M., Aigner, A., Berkhemer, O. A., Yoo, A. J., Roos, Y. B., Dippel, D. W. J., Van Der Lugt, A., Van Oostenbrugge, R. J., Van Zwam,

UNIVERSITAS GADJAH MADA
PARAMETER KUANTITATIF EEG (QEEG) SEBAGAI PREDIKTOR LUARAN FUNGSIONAL JANGKA PANJANG PASIEN STROKE ISKEMIK
DYANNE PARAMITA A P, dr. Astuti, Sp.S (K).; Dr. dr. Ahmad Asmedi, M.Kes, Sp.S (K)
Universitas Gadjah Mada, 2019 | Diunduh dari http://etd.repository.ugm.ac.id/

W. H., Fiehler, J., Marquering, H. A. & Majoie, C. B. L. M. 2017 . Association of Computed Tomography Ischemic Lesion Location with Functional Outcome in Acute Large Vessel Occlusion Ischemic Stroke. *Stroke*, 48(9), pp. 2426–2433. doi: 10.1161/STROKEAHA.117.017513.

Ervasti, J., Virtanen, M., Lallukka, T., Friberg, E., Mittendorfer-Rutz, E., Lundström, E. & Alexanderson, K. 2017. Permanent work disability before and after ischaemic heart disease or stroke event: A nationwide population-based cohort study in Sweden, *BMJ Open*, 7(9). doi: 10.1136/bmjopen-2017-017910.

Ezzat Nazzal, M., Ahmed Saadah, M., Mohammed Saadah, L. & Mustafa Trebinjac, S. 2009. Acute ischemic stroke: Relationship of brain lesion location & functional outcome, *Disability and Rehabilitation*, 31(18), pp. 1501–1506. doi: 10.1080/09638280802627702.

Finnigan, S. P., Rose, S. E., Walsh, M., Griffin, M., Janke, A. L., McMahon, K. L., Gillies, R., Strudwick, M. W., Pettigrew, C. M., Semple, J., Brown, J., Brown, P. & Chalk, J. B. 2004. Correlation of Quantitative EEG in Acute Ischemic Stroke with 30-Day NIHSS Score: Comparison with Diffusion and Perfusion MRI, *Stroke*, 35(4), pp. 899–903. doi: 10.1161/01.STR.0000122622.73916.d2.

Finnigan, S. & van Putten, M. J. A. M. 2013. EEG in ischaemic stroke: Quantitative EEG can uniquely inform (sub-)acute prognoses and clinical management. *Clinical Neurophysiology*. International Federation of Clinical Neurophysiology, 124(1), pp. 10–19. doi: 10.1016/j.clinph.2012.07.003.

Finnigan, S., Wong, A. & Read, S. 2016. Defining abnormal slow EEG activity in acute ischaemic stroke: Delta/alpha ratio as an optimal QEEG index, *Clinical Neurophysiology*. International Federation of Clinical Neurophysiology, 127(2), pp. 1452–1459. doi: 10.1016/j.clinph.2015.07.014.

Frost, S. B. 2003. Reorganization of Remote Cortical Regions After Ischemic Brain Injury: A Potential Substrate for Stroke Recovery, *Journal of Neurophysiology*, 89(6), pp. 3205–3214. doi: 10.1152/jn.01143.2002.

Frykholm, P., 2002. Cerebral ischemia studied with PET and microdialysis. *Comprehensive Summaries Uppsala Dissertations* 91 : 5319 - 5328

Goyal, N., Tsivgoulis, G., Iftikhar, S., Khorchid, Y., Fawad Ishfaq, M., Doss, V. T., Zand, R., Chang, J., Alsherbini, K., Choudhri, A., Hoit, D., Alexandrov, A. V., Arthur, A. S. & Elijovich, L. 2017. Admission systolic blood pressure and outcomes in large vessel occlusion strokes treated with endovascular treatment, *Journal of NeuroInterventional Surgery*, 9(5), pp. 451–454. doi: 10.1136/neurintsurg-2016-012386.

Gray, V., Rice, C. L. & Garland, S. J. 2012. Factors that influence muscle weakness following stroke and their clinical implications: A critical review, *Physiotherapy Canada*, 64(4), pp. 415–426. doi: 10.3138/ptc.2011-03.

Guo, X., Wu, W. & Tong, S. 2017. Asymmetry of hemispheric interdependences in the early hours following unilateral stroke: An electrophysiological study in rats, *Proceedings of the Annual International Conference of the IEEE Engineering in Medicine and Biology Society, EMBS*, pp. 4363–

UNIVERSITAS
GADJAH MADA
PARAMETER KUANTITATIF EEG (QEEG) SEBAGAI PREDIKTOR LUARAN FUNGSIONAL JANGKA PANJANG PASIEN STROKE ISKEMIK
DYANNE PARAMITA A P, dr. Astuti, Sp.S (K).; Dr. dr. Ahmad Asmedi, M.Kes, Sp.S (K)
Universitas Gadjah Mada, 2019 | Diunduh dari http://etd.repository.ugm.ac.id/

4366. doi: 10.1109/EMBC.2017.8037822.

Gupta, S., Yadav, R. & Malhotra, A. 2016. Assessment of physical disability using Barthel index among elderly of rural areas of district Jhansi (U.P), India, *Journal of Family Medicine and Primary Care*, 5(4), p. 853. doi: 10.4103/2249-4863.201178.

Hara, Y. 2015. Brain Plasticity and Rehabilitation in Stroke Patients. *Journal of Nippon Medical School*, 82(1), pp. 4–13. doi: 10.1272/jnms.82.4.

Harvey, R. L. 2015. Predictors of Functional Outcome Following Stroke. *Physical Medicine and Rehabilitation Clinics of North America*. Elsevier Inc, 26(4), pp. 583–598. doi: 10.1016/j.pmr.2015.07.002.

Hatfield, L. A., Crone, N. E., Kossoff, E. H., Ewen, J. B., Pyzik, P. L., Lin, D. D. M., Kelley, T. M. & Comi, A. M. 2007. Quantitative EEG asymmetry correlates with clinical severity in unilateral Sturge-Weber syndrome, *Epilepsia*, 48(1), pp. 191–195. doi: 10.1111/j.1528-1167.2006.00630.x.

Hauser, W. A. 1992. Seizure Disorders: The Changes With Age, *Epilepsia*, 33, pp. 6–14. doi: 10.1111/j.1528-1157.1992.tb06222.x.

Heiss, W. D. & Kidwell, C. S. 2014. Imaging for prediction of functional outcome and assessment of recovery in ischemic stroke, *Stroke*, 45(4), pp. 1195–1201. doi: 10.1161/STROKEAHA.113.003611.

Higashida, R. T., Furlan, A. J., Roberts, H., Tomsick, T., Connors, B., Barr, J., Dillon, W., Warach, S., Broderick, J., Tilley, B. & Sacks, D. 2003. Erratum: Trial Design and Reporting Standards for Intra-Arterial Cerebral Thrombolysis for Acute Ischemic Stroke (Stroke (August 2003) 34:8 (1156-1162))', *Stroke*, 34(11), p. 2774. doi: 10.1161/01.STR.0000101763.87465.A4.

Huybrechts, K. F. & Caro, J. J. 2007. The Barthel Index and modified Rankin Scale as prognostic tools for long-term outcomes after stroke: A qualitative review of the literature, *Current Medical Research and Opinion*, 23(7), pp. 1627–1636. doi: 10.1185/030079907X210444.

Jia, Q., Zhao, X., Wang, C., Wang, Yilong, Yan, Y., Li, H., Zhong, L., Liu, L., Zheng, H., Zhou, Y. & Wang, Yongjun. 2011. Diabetes and poor outcomes within 6 months after acute ischemic stroke: The China national stroke registry, *Stroke*, 42(10), pp. 2758–2762. doi: 10.1161/STROKEAHA.111.621649.

Johnston, K. C., Bruno, A., Pauls, Q., Hall, C. E., Barrett, K. M., Barsan, W., Fansler, A., Van De Bruinhorst, K., Janis, S. & Durkalski-Mauldin, V. L. 2019. Intensive vs Standard Treatment of Hyperglycemia and Functional Outcome in Patients with Acute Ischemic Stroke: The SHINE Randomized Clinical Trial, *JAMA - Journal of the American Medical Association*, 322(4), pp. 326–335. doi: 10.1001/jama.2019.9346.

Jöonsson, A. C., Lindgren, I., Norrving, B. & Lindgren, A. 2008. Weight loss after stroke: A population-based study from the Lund Stroke Register. *Stroke*, 39(3), pp. 918–923. doi: 10.1161/STROKEAHA.107.497602.

Katzan, I. L., Dawson, N. V., Thomas, C. L., Votruba, M. E. & Cebul, R. D. 2007. The cost of pneumonia after acute stroke, *Neurology*, 68(22), pp. 1938–1943. doi: 10.1212/01.wnl.0000263187.08969.45.

UNIVERSITAS GADJAH MADA
PARAMETER KUANTITATIF EEG (QEEG) SEBAGAI PREDIKTOR LUARAN FUNGSIONAL JANGKA PANJANG PASIEN STROKE ISKEMIK
DYANNE PARAMITA A P, dr. Astuti, Sp.S (K).; Dr. dr. Ahmad Asmedi, M.Kes, Sp.S (K)
Universitas Gadjah Mada, 2019 | Diunduh dari http://etd.repository.ugm.ac.id/

Kelly, J., Hunt, B. J., Lewis, R. R., Swaminathan, R., Moody, A., Seed, P. T. & Rudd, A. 2004. Dehydration and venous thromboembolism after acute stroke. *QJM - Monthly Journal of the Association of Physicians*, 97(5), pp. 293–296. doi: 10.1093/qjmed/hch050.

Khakshour, M. 2015. all things about QEEG (Rochester Institute of Technology). (August), pp. 0–37. doi: 10.13140/RG.2.1.2174.6403.

Khalaf, G. M., Abdkadhim, M. & Suhail, I. M. 2018. Value of Alberta Stroke Program early Computed Tomography Score (ASPECTS) in the Evaluation of Middle Cerebral Artery Acute Ischemic Stroke using CT and DW-MRI, *Journal of Medical Diagnostic Methods*, 07(04), pp. 1–6. doi: 10.4172/2168-9784.1000280.

Khan, M., Baird, G. L., Goddeau, R. P., Silver, B. & Henninger, N. 2017. Alberta stroke program early CT score infarct location predicts outcome following M2 occlusion, *Frontiers in Neurology*, 8(MAR), pp. 1–6. doi: 10.3389/fneur.2017.00098.

Kirchhof, P. &Lambert. E. 2016. 2016 ESC Guidelines for the management of atrial fibrillation developed in collaboration with EACTS, *European Heart Journal*, 37(38). doi: 10.1093/eurheartj/ehw210.

van der Knaap, L. J. & van der Ham, I. J. M. 2011. How does the corpus callosum mediate interhemispheric transfer? A review.*Behavioural Brain Research*. Elsevier B.V., 223(1), pp. 211–221. doi: 10.1016/j.bbr.2011.04.018.

Kongsawasdi, S., Klaphajone, J., Wivatvongvana, P. & Watcharasaksilp, K. 2019. Prognostic Factors of Functional Outcome Assessed by Using the Modified Rankin Scale in Subacute Ischemic Stroke, *Journal of Clinical Medicine Research*, 11(5), pp. 375–382. doi: 10.14740/jocmr3799.

Koton, S., Molshatzki, N., Bornstein, N. M. & Tanne, D.2012. Low cholesterol, statins and outcomes in patients with first-ever acute ischemic stroke, *Cerebrovascular Diseases*, 34(3), pp. 213–220. doi: 10.1159/000342302.

Leon-Carrion, J., Martin-Rodriguez, J. F., Damas-Lopez, J., Barroso y Martin, J. M. & Dominguez-Morales, M. R. 2009. Delta-alpha ratio correlates with level of recovery after neurorehabilitation in patients with acquired brain injury, *Clinical Neurophysiology*. International Federation of Clinical Neurophysiology, 120(6), pp. 1039–1045. doi: 10.1016/j.clinph.2009.01.021.

Li, J., Zhang, P., Wu, S., Yi, X., Wang, C. & Liu, M. 2018. Factors associated with favourable outcome in large hemispheric infarctions, *BMC Neurology*. BMC Neurology, 18(1), pp. 1–8. doi: 10.1186/s12883-018-1148-7.

Li, S. 2017. Spasticity, motor recovery, & neural plasticity after stroke. *Frontiers in Neurology*, 8(APR), pp. 1–8. doi: 10.3389/fneur.2017.00120.

Liu-ambrose, T., Eng, J. J. & Ot, P. T. 2015.Exercise Training and Recreational Activities to Promote Executive Functions in Chronic Stroke : A Proof-of-concept Study, *Journal of Stroke and Cerebrovascular Diseases*. Elsevier Ltd, 24(1), pp. 130–137. doi: 10.1016/j.jstrokecerebrovasdis.2014.08.012.

Logar, C. & Boswell, M. 1991. The value of EEG-Mapping in focal cerebral lesions, *Brain Topography*, 3(4), pp. 441–446. doi: 10.1007/BF01129003.

Lu, Q., Huang, G., Chen, L., Li, W. & Liang, Z. 2019. Structural and functional

UNIVERSITAS GADJAH MADA
PARAMETER KUANTITATIF EEG (QEEG) SEBAGAI PREDIKTOR LUARAN FUNGSIONAL JANGKA PANJANG PASIEN STROKE ISKEMIK
DYANNE PARAMITA A P, dr. Astuti, Sp.S (K).; Dr. dr. Ahmad Asmedi, M.Kes, Sp.S (K)
Universitas Gadjah Mada, 2019 | Diunduh dari http://etd.repository.ugm.ac.id/

reorganization following unilateral internal capsule infarction contribute to neurological function recovery, *Neuroradiology*. Neuroradiology, 61(10), pp. 1181–1190. doi: 10.1007/s00234-019-02278-x.

Luo, H., Wang, X., Chen, C., Wang, J., Li, C., Xu, Z., Yang, X., Shi, W. & Zeng, C. 2015. Oxidative stress causes imbalance of renal renin angiotensin system (RAS) components and hypertension in obese Zucker rats, *Journal of the American Heart Association*, 4(2), pp. 1–14. doi: 10.1161/JAHA.114.001559.

Mahmoud Fouad, M., Mohamed Farag, S., Hegazy, M. I. & Abd Elalem Aziz, M. 2017. Prediction of Functional Outcome in Ischemic Stroke Patients: An Observational Study on Egyptian Population, *Cureus*, 9(6). doi: 10.7759/cureus.1392.

Mang, C. S., Campbell, K. L., Ross, C. J. D. & Boyd, L. A. 2013. Promoting Neuroplasticity for Motor Rehabilitation After Stroke: Considering the Effects of Aerobic Exercise & Genetic Variation on Brain-Derived Neurotrophic Factor, *Physical Therapy*, 93(12), pp. 1707–1716. doi: 10.2522/ptj.20130053.

Martin-schild, S., Albright, K. C., Mph, D. O., Tanksley, J., Jones, E. B., Grotta, J. C. & Savitz, S. I. 2012. NIH Public Access, 57(February 2010), pp. 42–45. doi: 10.1016/j.annemergmed.2010.06.564.Zero.

van Meer, M. P. A., Otte, W. M., van der Marel, K., Nijboer, C. H., Kavelaars, A., van der Sprenkel, J. W. B., Viergever, M. A. & Dijkhuizen, R. M. 2012. Extent of bilateral neuronal network reorganization and functional recovery in relation to stroke severity', *Journal of Neuroscience*, 32(13), pp. 4495–4507. doi: 10.1523/JNEUROSCI.3662-11.2012.

Mehta, B., Leslie-Mazwi, T. M., Chandra, R. V., Chaudhry, Z. A., Rabinov, J. D., Hirsch, J. A., Schwamm, L. H., Rost, N. S. & Yoo, A. J. 2013. Assessing variability in neurointerventional practice patterns for acute ischemic stroke, *Journal of NeuroInterventional Surgery*, 5(SUPPL.1), pp. 4–6. doi: 10.1136/neurintsurg-2012-010565.

Melikian, N., Seddon, M. D., Casadei, B., Chowienczyk, P. J. & Shah, A. M. 2009. Neuronal nitric oxide synthase and human vascular regulation, *Trends in Cardiovascular Medicine*. Elsevier Inc., 19(8), pp. 256–262. doi: 10.1016/j.tcm.2010.02.007.

Miller, C. M. & Behrouz, R. 2016. Impact of Infection on Stroke Morbidity and Outcomes, *Current Neurology and Neuroscience Reports*. Current Neurology and Neuroscience Reports, 16(9), pp. 1–5. doi: 10.1007/s11910-016-0679-9.

Mori, M., Naganuma, M., Okada, Y., Hasegawa, Y., Shiokawa, Y., Nakagawara, J., Furui, E., Kimura, K., Yamagami, H., Kario, K., Okuda, S., Koga, M., Minematsu, K. & Toyoda, K. 2012. Early neurological deterioration within 24 hours after intravenous rt-PA therapy for stroke patients: The stroke acute management with urgent risk factor assessment and improvement rt-PA registry. *Cerebrovascular Diseases*, 34(2), pp. 140–146. doi: 10.1159/000339759.

Murphy, T. H. & Corbett, D. 2009. Plasticity during stroke recovery: From synapse

UNIVERSITAS GADJAH MADA
PARAMETER KUANTITATIF EEG (QEEG) SEBAGAI PREDIKTOR LUARAN FUNGSIONAL JANGKA PANJANG PASIEN STROKE ISKEMIK
DYANNE PARAMITA A P, dr. Astuti, Sp.S (K).; Dr. dr. Ahmad Asmedi, M.Kes, Sp.S (K)
Universitas Gadjah Mada, 2019 | Diunduh dari http://etd.repository.ugm.ac.id/

to behaviour. *Nature Reviews Neuroscience*. Nature Publishing Group, 10(12), pp. 861–872. doi: 10.1038/nrn2735.

Murri, L., Gori, S., Massetani, R., Bonanni, E., Marcella, F. & Milani, S.1998. Evaluation of acute ischemic stroke using quantitative EEG: A comparison with conventional EEG and CT scan. *Neurophysiologie Clinique*, 28(3), pp. 249–257. doi: 10.1016/S0987-7053(98)80115-9.

Musa, K. I. & Keegan, T. J. 2018. The change of Barthel Index scores from the time of discharge until 3-month post-discharge among acute stroke patients in Malaysia: A random intercept model. *PLoS ONE*, 13(12), pp. 1–13. doi: 10.1371/journal.pone.0208594.

Muscari, A., Puddu, G. M., Santoro, N. & Zoli, M. 2011. A simple scoring system for outcome prediction of ischemic stroke. *Acta Neurologica Scandinavica*, 124(5), pp. 334–342. doi: 10.1111/j.1600-0404.2010.01479.x.

Naito, H., Nezu, T., Hosomi, N., Aoki, S., Kinoshita, N., Kuga, J., Shimomura, R., Araki, M., Ueno, H., Ochi, K. & Maruyama, H. 2018. Controlling Nutritional Status score for predicting 3-mo functional outcome in acute ischemic stroke. *Nutrition*. Elsevier Inc., 55–56, pp. 1–6. doi: 10.1016/j.nut.2018.02.025.

Nakayama, E., Tohara, H., Hino, T., Sato, M., Hiraba, H., Abe, K. & Ueda, K. . 2014. The effects of ADL on recovery of swallowing function in stroke patients after acute phase. *Journal of Oral Rehabilitation*, 41(12), pp. 904–911. doi: 10.1111/joor.12212.

Ntaios, G., Gioulekas, F., Papavasileiou, V., Strbian, D. & Michel, P. 2016. ASTRAL, DRAGON and SEDAN scores predict stroke outcome more accurately than physicians. *European Journal of Neurology*, 23(11), pp. 1651–1657. doi: 10.1111/ene.13100.

Nudo, R. J. 2013. Recovery after brain injury: mechanisms and principles. *Frontiers in Human Neuroscience*, 7(December), pp. 1–14. doi: 10.3389/fnhum.2013.00887.

Pan, S. L., Lien, I. N. & Chen, T. H. H. 2010. Is Higher Serum Total Cholesterol Level Associated With Better Long-Term Functional Outcomes After Noncardioembolic Ischemic Stroke?. *Archives of Physical Medicine and Rehabilitation*. Elsevier Inc., 91(6), pp. 913–918. doi: 10.1016/j.apmr.2010.02.002.

Paolucci, S., Antonucci, G., Grasso, M. G. & Pizzamiglio, L. 2001. The role of unilateral spatial neglect in rehabilitation of right brain-damaged ischemic stroke patients: A matched comparison. *Archives of Physical Medicine and Rehabilitation*, 82(6), pp. 743–749. doi: 10.1053/apmr.2001.23191.

Park, J. S., Lee, J. M., Kwak, H. S. & Chung, G. H. 2019. Predictive value of CT angiography source image ASPECTS in patients with anterior circulation acute ischemic stroke after endovascular treatment: Ultimate infarct size and clinical outcome. *Journal of NeuroInterventional Surgery*, 11(4), pp. 342–346. doi: 10.1136/neurintsurg-2018-014359.

Parsons, M. W., Barber, P. A., Desmond, P. M., Baird, T. A., Darby, D. G., Byrnes, G., Tress, B. M. & Davis, S. M. 2002. Acute hyperglycemia adversely

UNIVERSITAS GADJAH MADA
PARAMETER KUANTITATIF EEG (QEEG) SEBAGAI PREDIKTOR LUARAN FUNGSIONAL JANGKA PANJANG PASIEN STROKE ISKEMIK
DYANNE PARAMITA A P, dr. Astuti, Sp.S (K).; Dr. dr. Ahmad Asmedi, M.Kes, Sp.S (K)
Universitas Gadjah Mada, 2019 | Diunduh dari http://etd.repository.ugm.ac.id/

affects stroke outcome: A magnetic resonance imaging and spectroscopy study. *Annals of Neurology*, 52(1), pp. 20–28. doi: 10.1002/ana.10241.

Penry, K. 1981. Proposal for Revised Clinical and Electroencephalographic Classification of Epileptic Seizures.. *Epilepsia*, 22(4), pp. 489–501. doi: 10.1111/j.1528-1157.1981.tb06159.x.

van Putten, M. J. A. M. 2007. The revised brain symmetry index. *Clinical Neurophysiology*, 118(11), pp. 2362–2367. doi: 10.1016/j.clinph.2007.07.019.

Van Putten, M. J. A. M. & Tavy, D. L. J. 2004. Continuous quantitative EEG monitoring in hemispheric stroke patients using the brain symmetry index. *Stroke*, 35(11), pp. 2489–2492. doi: 10.1161/01.STR.0000144649.49861.1d.

Quinn, T. J., Langhorne, P. & Stott, D. J. 2011. Barthel index for stroke trials: Development, properties, and application. *Stroke*, 42(4), pp. 1146–1151. doi: 10.1161/STROKEAHA.110.598540.

Ribo, M., Molina, C. A., Delgado, P., Rubiera, M., Delgado-Mederos, R., Rovira, A., Munuera, J. & Alvarez-Sabin, J. 2007. Hyperglycemia during ischemia rapidly accelerates brain damage in stroke patients treated with tPA, *Journal of Cerebral Blood Flow and Metabolism*, 27(9), pp. 1616–1622. doi: 10.1038/sj.jcbfm.9600460.

Rosso, C., Colliot, O., Pires, C., Delmaire, C., Valabrègue, R., Crozier, S., Dormont, D., Baillet, S., Samson, Y. & Lehéricy, S. 2011. Prédiction du handicap par les valeurs d'ADC dans les structures motrices à la phase hyper-aiguë de l'AVC ischémique étude comparative avec le volume en diffusion. *Journal of Neuroradiology*. Elsevier Masson SAS, 38(2), pp. 105–112. doi: 10.1016/j.neurad.2010.05.001.

Ryu, W. S., Schellingerhout, D., Jeong, S. W., Nahrendorf, M. & Kim, D. E. 2016. Association between Serum Lipid Profiles and Early Neurological Deterioration in Acute Ischemic Stroke. *Journal of Stroke and Cerebrovascular Diseases*. Elsevier Inc., 25(8), pp. 2024–2030. doi: 10.1016/j.jstrokecerebrovasdis.2016.05.009.

Saif, S. & Fazal, N. 2014. Association between NHI stroke scale score and functional outcome in acute ischemic stroke. *NIH Stroke Scale Score*, 64(4), pp. 585–590.

Sajobi, T. T. & Timothy, I.P. 2017. Early trajectory of stroke severity predicts long-term functional outcomes in ischemic stroke subjects. *Stroke*, 48(1), pp. 105–110. doi: 10.1161/STROKEAHA.116.014456.

Salinsky, M. C., Oken, B. S., Kramer, R. E. & Morehead, L. 1992. A comparison of quantitative EEG frequency analysis and conventional EEG in patients with focal brain lesions. *Electroencephalography and Clinical Neurophysiology*, 83(6), pp. 358–366. doi: 10.1016/0013-4694(92)90071-O.

Salter, K., Campbell, N., Richardson, M., Mehta, S., Jutai, J., Zettler, L., Ba, M. M., Mcclure, A., Mays, R., Foley, N. & Teasell, R. 2013. EBRSR [Evidence-Based Review of Stroke Rehabilitation] 20 Outcome Measures in Stroke Rehabilitation. *Evidence-Based Review of Stroke Rehabilitation*,

UNIVERSITAS GADJAH MADA
PARAMETER KUANTITATIF EEG (QEEG) SEBAGAI PREDIKTOR LUARAN FUNGSIONAL JANGKA PANJANG PASIEN STROKE ISKEMIK
DYANNE PARAMITA A P, dr. Astuti, Sp.S (K).; Dr. dr. Ahmad Asmedi, M.Kes, Sp.S (K)
Universitas Gadjah Mada, 2019 | Diunduh dari http://etd.repository.ugm.ac.id/

p. 141. Available at: http://www.ebrsr.com/evidence-review.

Santalucia, P., Pezzella, F. R., Sessa, M., Monaco, S., Torgano, G., Anticoli, S., Zanoli, E., Maimone Baronello, M., Paciaroni, M. & Caso, V. 2013.Sex differences in clinical presentation, severity and outcome of stroke: Results from a hospital-based registry. *European Journal of Internal Medicine*. European Federation of Internal Medicine., 24(2), pp. 167–171. doi: 10.1016/j.ejim.2012.10.004.

Sato, M., Ido, Y., Yoshimura, Y. & Mutai, H. 2019. Relationship of Malnutrition During Hospitalization With Functional Recovery and Postdischarge Destination in Elderly Stroke Patients. *Journal of Stroke and Cerebrovascular Diseases*. Elsevier Inc., 28(7), pp. 1866–1872. doi: 10.1016/j.jstrokecerebrovasdis.2019.04.012.

Scherbakov, N., Von Haehling, S., Anker, S. D., Dirnagl, U. & Doehner, W. 2013. Stroke induced Sarcopenia: Muscle wasting and disability after stroke. *International Journal of Cardiology*. Elsevier Ireland Ltd, 170(2), pp. 89–94. doi: 10.1016/j.ijcard.2013.10.031.

Scherbakov, N., Dirnagl, U. & Doehner, W. 2011. Body weight after stroke lessons from the obesity paradox. *Stroke*, 42(12), pp. 3646–3650. doi: 10.1161/STROKEAHA.111.619163.

Scherbakov, N. & Doehner, W. 2011. Sarcopenia in stroke-facts and numbers on muscle loss accounting for disability after stroke. *Journal of Cachexia, Sarcopenia and Muscle*, 2(1), pp. 5–8. doi: 10.1007/s13539-011-0024-8.

Setyopranoto, I., Wibowo, S., Sadewa, A. & Lamsudin, R. 2016. Reduced Vascular Endothelial Growth Factor A Levels are Correlated with the Incidence of Brain Edema in Acute Ischemic Stroke Patients. *British Journal of Medicine and Medical Research*, 13(2), pp. 1–7. doi: 10.9734/BJMMR/2016/23156.

Sharanreddy, M. & Kulkarni, P. K. 2013. Automated EEG signal analysis for identification of epilepsy seizures and brain tumour. *Journal of Medical Engineering and Technology*, 37(8), pp. 511–519. doi: 10.3109/03091902.2013.837530.

Sheorajpanday, R. V. A., Nagels, G., Weeren, A. J. T. M., Surgeloose, D. De & Deyn, P. P. De. 2010. Clinical Neurophysiology Additional value of quantitative EEG in acute anterior circulation syndrome of presumed ischemic origin. *Clinical Neurophysiology*. International Federation of Clinical Neurophysiology, 121(10), pp. 1719–1725. doi: 10.1016/j.clinph.2009.10.037.

Sheorajpanday, R. V. A., Nagels, G., Weeren, A. J. T. M. & Deyn, P. P. De. 2011. Clinical Neurophysiology Quantitative EEG in ischemic stroke : Correlation with infarct volume and functional status in posterior circulation and lacunar syndromes. *Clinical Neurophysiology*. International Federation of Clinical Neurophysiology, 122(5), pp. 884–890. doi: 10.1016/j.clinph.2010.08.020.

Sheorajpanday, R. V. A., Nagels, G., Weeren, A. J. T. M., van Putten, M. J. A. M. & De Deyn, P. P. 2011. Quantitative EEG in ischemic stroke: Correlation with functional status after 6months. *Clinical Neurophysiology*.

UNIVERSITAS GADJAH MADA
PARAMETER KUANTITATIF EEG (QEEG) SEBAGAI PREDIKTOR LUARAN FUNGSIONAL JANGKA PANJANG PASIEN STROKE ISKEMIK
DYANNE PARAMITA A P, dr. Astuti, Sp.S (K).; Dr. dr. Ahmad Asmedi, M.Kes, Sp.S (K)
Universitas Gadjah Mada, 2019 | Diunduh dari http://etd.repository.ugm.ac.id/

International Federation of Clinical Neurophysiology, 122(5), pp. 874–883. doi: 10.1016/j.clinph.2010.07.028.

Shimoyama, T., Kimura, K., Uemura, J., Saji, N. & Shibazaki, K. 2014. Elevated glucose level adversely affects infarct volume growth and neurological deterioration in non-diabetic stroke patients, but not diabetic stroke patients. *European Journal of Neurology*, 21(3), pp. 402–410. doi: 10.1111/ene.12280.

Slot, K. B., Berge, E., Sandercock, P., Lewis, S. C., Dorman, P. & Dennis, M. 2009. Causes of death by level of dependency at 6 months after ischemic stroke in 3 large cohorts. *Stroke*, 40(5), pp. 1585–1589. doi: 10.1161/STROKEAHA.108.531533.

Song, J., Davey, C., Poulsen, C., Luu, P., Turovets, S., Anderson, E., Li, K. & Tucker, D. 2015. EEG source localization: Sensor density and head surface coverage. *Journal of Neuroscience Methods*. Elsevier B.V., 256, pp. 9–21. doi: 10.1016/j.jneumeth.2015.08.015.

Strong, K., Mathers, C. & Bonita, R. 2007. Preventing stroke : saving lives around the world', 6(February), pp. 182–187.

Syeban, Z., Octaviana, F. 2014. EEG Praktis. *Badan Penerbit Fakultas Kedokteran Universitas Indonesia.* hal 1-17

Takeuchi, N., Oouchida, Y. & Izumi, S. I. 2012. Motor control and neural plasticity through interhemispheric interactions. *Neural Plasticity*, 2012. doi: 10.1155/2012/823285.

Thorpe, K. E. (2013) 'outcomes'.

Toschke, A. M., Tilling, K., Cox, A. M., Rudd, A. G., Heuschmann, P. U. & Wolfe, C. D. A. 2010. Patient-specific recovery patterns over time measured by dependence in activities of daily living after stroke and post-stroke care : The South London Stroke Register (SLSR)', pp. 219–225. doi: 10.1111/j.1468-1331.2009.02774.x.

Tunjungsari, A.P., Asmedi, A. Paryono. 2018. AbnormalitasGelombang Elektroensefalografi sebagai Prediktor Perburukan Defisit Neurologis Pada Stroke Iskemik Akut.

Tsai, Y. H., Yuan, R., Huang, Y. C., Yeh, M. Y., Lin, C. P. & Biswal, B. B. 2014. Disruption of brain connectivity in acute stroke patients with early impairment in consciousness. *Frontiers in Psychology*, 4(JAN), pp. 1–10. doi: 10.3389/fpsyg.2013.00956.

Tsao, J. W., Hemphill, J. C., Johnston, S. C., Smith, W. S. & Bonovich, D. C. 2005. Initial Glasgow Coma Scale score predicts outcome following thrombolysis for posterior circulation stroke. *Archives of Neurology*, 62(7), pp. 1126–1129. doi: 10.1001/archneur.62.7.1126.

Tsivgoulis, G., Kotsis, V. & Giannopoulos, S. 2011. Intravenous thrombolysis for acute ischaemic stroke: Effective blood pressure control matters', *International Journal of Stroke*, 6(2), pp. 125–127. doi: 10.1111/j.1747-4949.2010.00570.x.

Tsolaki, A., Kazis, D., Kompatsiaris, I., Kosmidou, V. & Tsolaki, M. 2014. Electroencephalogram and alzheimer's disease: Clinical and research approaches. *International Journal of Alzheimer's Disease*. Hindawi

UNIVERSITAS GADJAH MADA
PARAMETER KUANTITATIF EEG (QEEG) SEBAGAI PREDIKTOR LUARAN FUNGSIONAL JANGKA PANJANG PASIEN STROKE ISKEMIK
DYANNE PARAMITA A P, dr. Astuti, Sp.S (K).; Dr. dr. Ahmad Asmedi, M.Kes, Sp.S (K)
Universitas Gadjah Mada, 2019 | Diunduh dari http://etd.repository.ugm.ac.id/

Publishing Corporation, 2014. doi: 10.1155/2014/349249.

Ultra1, P. 2001. qEEG mechanism of , 117(September), pp. 1–9. doi: 10.1111/j.1747-4949.2009.00337.x.

Vemmos, K. N., Tsivgoulis, G., Spengos, K., Zakopoulos, N., Synetos, A., Kotsis, V., Vassilopoulos, D. & Mavrikakis, M. 2003. Association between 24-h blood pressure monitoring variables and brain oedema in patients with hyperacute stroke. *Journal of Hypertension*, 21(11), pp. 2167–2173. doi: 10.1097/00004872-200311000-00027.

Wang, H., Camicia, M., Terdiman, J., Hung, Y.Y., Sandel, M.E., 2011. Time to inoatient rehabilitation hospital admission and functional outcomes of stroke patients. *PMR* 3, 296-304.

Ward, N. 2017. 20 Neural plasticity for functional rehabilitation after stroke. *Journal of Neurology, Neurosurgery & Psychiatry*, 88(8), p. A10.2-A10. doi: 10.1136/jnnp-2017-BNPA.20.

Ward, N. S. & Cohen, L. G. 2004. Europe PMC Funders Group Mechanisms Underlying Recovery of Motor Function After Stroke. *Arch. Neurol.*, 61(12), pp. 1844–1848. doi: 10.1001/archneur.61.12.1844.Mechanisms.

Winstein, C. J. & Wuu. L.M. 2016. *AHA / ASA Guideline Guidelines for Adult Stroke Rehabilitation and Recovery*. doi: 10.1161/STR.0000000000000098.

De Wit, L., Putman, K., Devos, H., Brinkmann, N., Dejaeger, E., De Weerdt, W., Jenni, W., Lincoln, N., Schuback, B. & Schupp, W. 2014. Long-term prediction of functional outcome after stroke using single items of the Barthel Index at discharge from rehabilitation centre. *Disability and Rehabilitation*, 36(5), pp. 353–358. doi: 10.3109/09638288.2013.793411.

Wu, J., Srinivasan, R., Burke Quinlan, E., Solodkin, A., Small, S. L. & Cramer, S. C. 2016. Utility of EEG measures of brain function in patients with acute stroke. *Journal of Neurophysiology*, 115(5), pp. 2399–2405. doi: 10.1152/jn.00978.2015.

Yagi, M., Yasunaga, H., Matsui, H., Morita, K., Fushimi, K., Fujimoto, M., Koyama, T. & Fujitani, J. 2017.Impact of Rehabilitation on Outcomes in Patients with Ischemic Stroke: A Nationwide Retrospective Cohort Study in Japan', *Stroke*, 48(3), pp. 740–746. doi: 10.1161/STROKEAHA.116.015147.

Yoo, A. J., Zaidat, O. O., Chaudhry, Z. A., Berkhemer, O. A., Gilberto Gonzalez, R., Goyal, M., Demchuk, A. M., Menon, B. K., Mualem, E., Ueda, D., Buell, H., Sit, S. P. & Bose, A. 2014. Impact of pretreatment noncontrast CT Alberta stroke program early ct score on clinical outcome after intra arterial stroke therapy', *Stroke*, 45(3), pp. 746–751. doi: 10.1161/STROKEAHA.113.004260.

Yu, A. Y. X., Holodinsky, J. K., Zerna, C., Svenson, L. W., Jetté, N., Quan, H. & Hill, M. D. 2016. Use and utility of administrative health data for stroke research and surveillance', *Stroke*, 47(7), pp. 1946–1952. doi: 10.1161/STROKEAHA.116.012390.

Zhao, Y., Zou, C., Wang, C., Zhang, Y. & Wang, S. 2016. Long-term outcomes after stroke in elderly patients with atrial fibrillation: A hospital-based

UNIVERSITAS
GADJAH MADA
PARAMETER KUANTITATIF EEG (QEEG) SEBAGAI PREDIKTOR LUARAN FUNGSIONAL JANGKA PANJANG PASIEN STROKE ISKEMIK
DYANNE PARAMITA A P, dr. Astuti, Sp.S (K).; Dr. dr. Ahmad Asmedi, M.Kes, Sp.S (K)
Universitas Gadjah Mada, 2019 | Diunduh dari http://etd.repository.ugm.ac.id/

follow-up study in China', *Frontiers in Aging Neuroscience*, 8(MAR), pp. 1–6. doi: 10.3389/fnagi.2016.00056.