

DAFTAR PUSTAKA

- Abraham, S. & Weismann, C.G. 2016. Left ventricular end-systolic eccentricity index for assessment of pulmonary hypertension in infants. *Echocardiogr.* 33(6):910–915.
- Al-naamani, N., Trammell, A.W., Woods, D.R., Mellor, A., Begley, J., Stacey, et al. 2014. Recommendations for the use of natriuretic peptides in acute cardiac care: A position statement from the Study Group on Biomarkers in Cardiology of the ESC Working Group on Acute Cardiac Care. *Eur Heart J.* 33(4):1190–1194.
- Al-Naamani, N., Trammell, A.W., Safdar, Z. 2019. Circulating biomarkers in pulmonary arterial hypertension. *Advances PH.* 14(1):21–27.
- Amdani, S.M., Mian, M.U.M., Thomas, R.L., Ross, R.D. 2018. NT-pro BNP—A marker for worsening respiratory status and mortality in infants and young children with pulmonary hypertension. *Congenit Heart Dis.* 13(4):499–505.
- Baggen, V.J.M., Van Den Bosch, A.E., Eindhoven, J.A., Schut, A.R.W., Cuypers, J.A.A.E., Witsenburg, M., et al. 2017. Prognostic value of n-terminal pro-b-type natriuretic peptide, troponin-t, and growth-differentiation factor 15 in adult congenital heart disease. *Circ.* 135(3):264–279.
- Barst, R.J., Ertel, S.I., Beghetti, M., Ivy, D.D. 2011. Pulmonary arterial hypertension: a comparison between children and adults. *The European respiratory journal.* 37(3):665–77.
- Bossone, E., Andrea, A.D., Alto, M.D., Citro, R., Argiento, P., Ferrara, F., et al. 2013. Echocardiography in pulmonary arterial hypertension : from diagnosis to prognosis. *J Am Soc Echocardiogr.* 26(1):1–14.
- D’Alto, M. & Mahadevan, V.S. 2012. Pulmonary arterial hypertension associated with congenital heart disease. *Eur Respir Rev.* 21(126):328–337.
- Decroo, T., Henríquez-trujillo, A.R., Weggheleire, A. De., Lynen, L.

2017. Rational use of Xpert testing in patients with presumptive TB : clinicians should be encouraged to use the test-treat threshold. (December). *BMC Inf dis.* 17(6):674-80
- Dimopoulos, K., Wort, S.J., Gatzoulis, M. a. 2014. Pulmonary hypertension related to congenital heart disease: A call for action. *Eur Heart J.* 35(11):691–700.
- Ebell, M.H. 2009. Diagnosis: Making the best use of medical data. *American Family Physician.* 79(6):478–480.
- El-Korashy, R.I.M., Amin, Y.M., Eissa, A.I., Thabet, T.S. 2014. Echocardiography versus right heart catheterization in class I pulmonary hypertension. *Egypt J Chest Dis Tuberc.* 63(2):419–422.
- Elsharawy, S., Hassan, B., Morsy, S. & Khalifa, N. 2012. Diagnostic value of N-terminal pro-brain natriuretic peptide levels in pediatric patients with ventricular septal defect. *Egypt Heart J.* 64(4):241–246.
- Gan, C.T., McCann, G.P., Marcus, J.T., van Wolferen, S. a., Twisk, J.W., Boonstra, et al 2006. NT-proBNP reflects right ventricular structure and function in pulmonary hypertension. *Eur Respi J.* 28(6):1190–1194.
- Grimes, D.A. & Schulz, K.F. 2005. Refining clinical diagnosis with likelihood ratios. *Lancet.* 365(panel 1):1500–1505.
- Guazzi, M. & Naeije, R. 2017. Pulmonary Hypertension in Heart Failure: Pathophysiology, Pathobiology, and Emerging Clinical Perspectives. *Journal of the American College of Cardiology.* 69(13):1718–1734.
- Hajian-Tilaki, K. 2013. Receiver operating characteristic (ROC) curve analysis for medical diagnostic test evaluation. *Caspian Journal of Internal Medicine.* 4(2):627–635.
- Hall, C. 2004. Essential biochemistry and physiology of (NT-pro)BNP. *Eur Journal Heart Fail.* 6(3):257–260.
- Hammerstingl, C., Schueler, R., Bors, L., Momcilovic, D., Pabst, S., Nickenig, G. & Skowasch, D. 2012. Diagnostic value of echocardiography in the diagnosis of pulmonary hypertension. *PLoS*

ONE. 7(6):2–8.

- Harbaum, L., Hennigs, J.K., Baumann, H.J., Lüneburg, N., Griesch, E., Bokemeyer, et al. 2014. N-terminal pro-brain natriuretic peptide is a useful prognostic marker in patients with pre-capillary pulmonary hypertension and renal insufficiency. *PLoS ONE*. 9(4):2–9.
- Heart, I.J.C., Parasuraman, S., Walker, S., Loudon, B.L., Gollop, N.D., Wilson, A.M., et al. 2016. Assessment of pulmonary artery pressure by echocardiography — A comprehensive review. *IJCHA*. 12:45–51.
- Hu, D., Liu, Y., Tao, H., Gao, J. 2015. Clinical value of plasma B-type natriuretic peptide assay in pediatric pneumonia accompanied by heart failure. *Exp Ther Med*. 10(6):2175–2179.
- Janda, S., Shahidi, N., Gin, K. & Swiston, J. 2011. Diagnostic accuracy of echocardiography for pulmonary hypertension : a systematic review and. *Heart*. 97:612–622.
- Jone, P.-N. & Ivy, D.D. 2014. Echocardiography in pediatric pulmonary hypertension. *Front Pediatr*. 2(November):1–15.
- Kangin Murat, Pirinccioglu, A.G. & Alyan, Ö. 2012. Plasma natriuretic peptides as predictors of hemodynamically in children with congenital heart disease . *Int Arch Med Res*. 4(2):8–14.
- Koch, A., Zink, S. & Singer, H. 2006. B-type natriuretic peptide in paediatric patients with congenital heart disease. *Eur Heart J*. 27(7):861–866.
- Koestenberger, M., Apitz, C., Abdul-Khaliq, H. & Hansmann, G. (in press). Transthoracic echocardiography for the evaluation of children and adolescents with suspected or confirmed pulmonary hypertension. *The European Paediatric Pulmo. Heart*. 102:14-22.
- Koura, H.M., Abdalla, N.M., Ibrahim, M.H., Abo Hashish, M.M.A. & Zaki, S.M. 2016. NT-proBNP in children with left to right shunt and dilated cardiomyopathy. *Iran J Pediatr*. 26(3).
- Leeflang, M.M.G., Rutjes, A.W., Reitsma, J.B., Hooft, L. & Bossuyt, P.M.M. 2013. Variation of a test’s sensitivity and specificity with

- disease prevalence. *CMAJ*. 185(11):537–544.
- Mocerri, P., Baudouy, D., Chiche, O., Cerboni, P., Bouvier, P., Chaussade, C., et al. 2014. Imaging in pulmonary hypertension: Focus on the role of echocardiography. *Arch Cardiovasc Dis*. 107(4):261–271.
- Montani, D. & Simonneau, G. 2012. Updated clinical classification of pulmonary hypertension. In *Pulmo Vasc Dis*. V. 62. 1–13.
- Nir, A. & Nasser, N. 2005. Clinical value of NT-ProBNP and BNP in pediatric cardiology. *J Card Fail*. 11(5 Suppl):S76–S80.
- Nir, A., Lindinger, A., Rauh, M., Bar-Oz, B., Laer, S., Schwachtgen, L., et al. 2009. NT-pro-B-type natriuretic peptide in infants and children: Reference values based on combined data from four studies. *Pediatr Card*. 30(1):3–8.
- Pauker, S.G. & Kassirer, J.P. 1980. The threshold approach to clinical decision making. *NEJM*. 302(20):1110–1117.
- Rashid, U., Qureshi, A., Hyder, S. & Sadiq, M. 2016. Pattern of congenital heart disease in a developing country tertiary care center: Factors associated with delayed diagnosis. *Ann Pediatr Car*. 9(3):210.
- Safari, S., Baratloo, A., Elfil, M. & Negida, A. 2016. Evidence Based Emergency Medicine ; Part 5 Receiver Operating Curve and Area under the Curve. 4(2):111–113.
- Skinner, G.J. 2017. Echocardiographic Assessment of Pulmonary Arterial Hypertension for Pediatricians and Neonatologists. *Front Pediatr*. 5(September):1–8.
- Sugimoto, M., Kuwata, S., Kurishima, C., Kim, J.H., Iwamoto, Y. & Senzaki, H. 2015. Cardiac biomarkers in children with congenital heart disease. *World J Pediatr*. 11(4):309–315.
- Ten Kate, C.A., Tibboel, D. & Kraemer, U.S. 2015. B-type natriuretic peptide as a parameter for pulmonary hypertension in children. A systematic review. *Eur J Pediatr*. 174(10):1267–1275.
- Tenny, S. & Hoffman, M.R. 2019. Prevalence. In *NCBI book shelf. A service of the National Library of Medicine, National Institutes of*

Health. NCBI. 1–5.

Woods, D.R., Mellor, a., Begley, J., Stacey, M., O’Hara, J., Hawkins, a., Yarker, J., Foxen, S., et al. 2013. Brain natriuretic peptide and NT-proBNP levels reflect pulmonary artery systolic pressure in trekkers at high altitude. *Physiological Research*. 62(6):597–603.

Zhao, Q.M., Ma, X.J., Jia, B. & Huang, G.Y. 2013. Prevalence of congenital heart disease at live birth: An accurate assessment by echocardiographic screening. *Acta Paediatr*. 102(4):397–402.