



DAFTAR PUSTAKA

- Almeida, S.M., Furtado, J.M., Mascarenhas, P., Ferraz, M.E., Silva, L.R., Ferreira, J.C., Monteiro, M., Vilanova, M., Ferraz, F.P. 2016. Anthropometric predictors of body fat in a large population of 9-year-old school-aged children. *Obes Sci Pract*, 2(3): 272–281.
- Badan Penelitian dan Pengembangan Kesehatan Kementerian Kesehatan RI. 2018. *Laporan Nasional RISKESDAS 2018*. Badan Penelitian dan Pengembangan Kesehatan Kementerian Kesehatan RI, Jakarta.
- Badan Penelitian dan Pengembangan Kesehatan Kementerian Kesehatan RI. 2013. *Riset Kesehatan Dasar RISKESDAS 2013*. Badan Penelitian dan Pengembangan Kesehatan Kementerian Kesehatan RI, Jakarta.
- Batubara, J.R. 2016. Practices of growth assessment in children: Is anthropometric measurement important? *Paediatr Indones*, 45(4): 145.
- Butcher, A., Kabiri, L.S., Brewer, W., Ortiz, A. 2018. Criterion validity and sensitivity to change of a pediatric bioelectrical impedance analysis scale in adolescents. *Child Obes*, 15(2): 142–148.
- Chula de Castro, J.A., Lima, T.R. de, Silva, D.A.S. 2018. Body composition estimation in children and adolescents by bioelectrical impedance analysis: A systematic review. *J Bodyw Mov Ther*, 22(1): 134–146.
- Collins, S., Duffield, A., Myatt, M. 2000. Assessment of nutritional status in emergency-affected populations. Center for Disease Control and Prevention.
- Daniels, S.R. 2009. The use of BMI in the clinical setting. *Pediatrics*, 124 Suppl(Supplement 1): S35-41.
- De Ferranti, S., Mozaffarian, D. 2008. The perfect storm: Obesity, adipocyte dysfunction, and metabolic consequences. *Clin Chem*, 54(6): 945–955.
- Ejtahed, H.S., Asghari, G., Mirmiran, P., Hosseinpour-Niazi, S., Sherafat-KazemZadeh, R., Azizi, F. 2014. Body mass index as a measure of percentage body fat prediction and excess adiposity diagnosis among Iranian adolescents. *Arch Iran Med*, 17(6): 400–405.
- Ferretti, R.D.L., Cintra, I.D.P., Passos, M.A.Z., Ferrari, G.L.D.M., Fisberg, M. 2015. Elevated neck circumference and associated factors in adolescents. *BMC Public Health*, 15(1): 1–10.
- Fink, B. 2012. Neck circumference: its usage in medicine and biology. In *Handbook of Anthropometry*. New York, NY: Springer New York: 665–675.
- Gamliel, A., Ziv-Baran, T., Siegel, R.M., Fogelman, Y., Dubnov-Raz, G. 2015. Using weight-for-age percentiles to screen for overweight and obese children and adolescents. *Prev Med (Baltim)*, 81: 174–179.
- Goluch-Koniuszy, Z., Kuchlewska, M. 2017. Body composition in 13-year-old adolescents with abdominal obesity, depending on the BMI value. *Adv Clin Exp Med*, 26(6): 973–979.
- Himah, R., Prawirohartono, E.P., Julia, M. 2008. Association between obesity and lipid profile in children 10-12 years of age. *Paediatr Indones*, 48(5): 257–260.
- Himes, J.H. 2009. Challenges of accurately measuring and using BMI and other indicators of obesity in children. *Pediatrics*, 124(supplement 1): s3–s22.



- Hingorjo, M.R., Qureshi, M.A., Mehdi, A. 2012. Neck circumference as a useful marker of obesity: A comparison with body mass index and waist circumference. *J Pak Med Assoc*, 62(1): 36–40.
- Horan, M., Gibney, E., Molloy, E., McAuliffe, F. 2015. Methodologies to assess paediatric adiposity. *Irish J Med Sci*, 184(1): 53–68.
- Kelishadi, R., Djalalinia, S., Motlagh, M.E., Rahimi, A., Bahreynian, M., Arefirad, T., Ardalan, G., Safiri, S., Hasani, M., Asayesh, H., Mansourian, M., Qorbani, M. 2016. Association of neck circumference with general and abdominal obesity in children and adolescents: The weight disorders survey of the CASPIAN-IV study. *BMJ Open*, 6(9): e011794.
- Kiess, W., Blüher, S., Kapellen, T., Garten, A., Klammt, J., Kratzsch, J., Körner, A. 2006. Physiology of obesity in childhood and adolescence. *Curr Paediatr*, 16(2): 123–131.
- Kim, Y., Lee, J.-M., Laurson, K., Bai, Y., Gaesser, G.A., Welk, G.J. 2014. Accuracy of neck circumference in classifying overweight and obese US children. *ISRN Obes*, 2014: 1–6.
- Kyle, U.G., Bosaeus, I., De Lorenzo, A.D., Deurenberg, P., Elia, M., Gómez, J.M., Heitmann, B.L., Kent-Smith, L., Melchior, J.C., Pirllich, M., Scharfetter, H., Schols, A.M.W.J., Pichard, C. 2004. Bioelectrical impedance analysis - Part I: Review of principles and methods. *Clin Nutr*, 23(5): 1226–1243.
- Labor, C., Trevisol, D., Schuelter, F. 2018. Waist-to-height ratio as a screening tool for childhood obesity : A systematic literature review. *Ann Pediatr Child Heal*, 6: 1–6.
- Martin-Calvo, N., Moreno-Galarraga, L., Martinez-Gonzalez, M.A. 2016. Association between body mass index, waist-to-height ratio and adiposity in children: A systematic review and meta-analysis. *Nutrients*, 8(8): 512.
- Mazıcıoğlu, M.M., Hatipoğlu, N., Oztürk, A., Ciçek, B., Ustünbaş, H.B., Kurtoglu, S. 2010. Waist circumference and mid-upper arm circumference in evaluation of obesity in children aged between 6 and 17 years. *J Clin Res Pediatr Endocrinol*, 2(4): 144–150.
- McCarthy, H.D., Cole, T.J., Fry, T., Jebb, S.A., Prentice, A.M. 2006. Body fat reference curves for children. *Int J Obes*, 30(4): 598–602.
- Mulyasari, I., Pontang, G.S. 2018. Waist circumference and waist-to-height ratio as indicators for excess adiposity in adolescents. *J Gizi dan Pangan*, 13(3): 131–136.
- Pereira, P.F., Serrano, H.M.S., Carvalho, G.Q., Lamounier, J.A., Peluzio, M. do C.G., Franceschini, S. do C.C., Priore, S.E. 2010. Waist Circumference as an indicator of body fat and metabolic disorders in adolescents: A comparison of four criteria. *Rev Assoc Med Bras*, 56(6): 665–669.
- Sijtsma, A., Bocca, G., L'abée, C., Liem, E.T., Sauer, P.J.J., Corpeleijn, E. 2014. Waist-to-height ratio, waist circumference and BMI as indicators of percentage fat mass and cardiometabolic risk factors in children aged 3-7 years. *Clin Nutr*, 33(2): 311–315.
- Sjarif, D.R., Gultom, L.C., Hendarto, A., Lestari, E.D., Sidiartha, I.G.L., Mexitalia, M. 2014. *Diagnosis, tata laksana dan pencegahan obesitas*.



UNIVERSITAS
GADJAH MADA

INDEKS MASSA TUBUH, WAIST CIRCUMFERENCE, WAIST TO HEIGHT RATIO, DAN LINGKAR
LEHER SEBAGAI ALAT
SKRINING UNTUK MENILAI OBESITAS BERDASARKAN PERSENTASE LEMAK TUBUH PADA
REMAJA DI DAERAH ISTIMEWA

YOGYAKARTA

RAHMA ANINDITA, Prof. dr. Madarina Julia, Ph.D, Sp.AK.; dr. Neti Nurani, M.Sc, SP.AK

Universitas Gadjah Mada, 2019 | Diunduh dari <http://etd.repository.ugm.ac.id/>

Jakarta: IDAI.

- Talma, H., Chinapaw, M.J.M., Bakker, B., HiraSing, R.A., Terwee, C.B., Altenburg, T.M. 2013. Bioelectrical impedance analysis to estimate body composition in children and adolescents: a systematic review and evidence appraisal of validity, responsiveness, reliability and measurement error. *Obes Rev*, 14(11): 895–905.
- Taylor, R.W., Grant, A.M., Williams, S.M., Goulding, A. 2010. Sex differences in regional body fat distribution from pre-to postpuberty. *Obesity*, 18(7): 1410–1416.
- WHO. 2017. *Assessing and managing children at primary health-care facilities to prevent overweight and obesity in the context of the double burden of malnutrition, Updates for The Integrated Management of Childhood Illness (IMCI)*. WHO, New York
- Weber, D.R., Leonard, M.B., Shults, J., Zemel, B.S. 2014. A comparison of fat and lean body mass index to BMI for the identification of metabolic syndrome in children and adolescents. *J Clin Endocrinol Metab*, 99(9): 3208–3216.
- WHO. 2008. Waist circumference and waist-hip ratio report of a WHO expert consultation. WHO, New York.