

REFERENCES

- IUCN, 2018, The IUCN Red List of Threatened Species, diakses pada 7 June 2019
<http://www.iucnredlist.org/details/166894/0>
- Appelbaum, S., Schemmel. 1983. Dermal Sense Organs and Their Significance in the Feeding Behaviour of the Common Sole *Solea Vulgaris*, *Marine Ecology Progress Series* 13, no. 1 (1983): 29-36.
- Budelmann, B.U., Bleckmann, H.. 1988. A lateral line analogue in cephalopods: Water waves generate microphonic potentials in the epidermal head lines of *Sepia* and *Loliguncula*, *Journal of Comparative Physiology A* 164 (1): 1–5.
- Deelder, C.L. - 1985. Exposé synoptique des données biologiques sur l'anguille, *Anguilla anguilla* (Linnaeus, 1758). *FAO Synop. Pêches*, (80) Rev.1 . 71 p ..
- Egginton, S., Johnston, A., 1982, Muscle Fibre Differentiation and Vascularisation in Juvenile European Eel *Anguilla Anguilla* L.), *Cell and Tissue Research. springer. Verlag.* 222:563-577
- Fischer, E. R., Hansen, B. T., Nair, V., Hoyt, F. H., Dorward, D. W., 2012, Scanning Electron Microscopy, *Current Protocols in Microbiology*.
- Ghattas, S.M., Yanai, T. 2010. Light Microscopical Study on the skin of European Eel (*Anguilla Anguilla*). *World Journal of Fish and Marine Science* 2(3): 152-161
- Herianti, I. 2005. *Rekayasa Lingkungan Untuk Memacu Perkembangan Ovarium Ikan Sidat (Anguilla bicolor)*. *Oseanologi dan Limnologi* No. 37: 25-41.
- Jacoby, D., Harrison, I.J., Gollock, M. 2014. *Anguilla bicolor* . The IUCN Red List of Threatened Species 2014: e.T166894A67015710.
- Leonard, J. B., Summers, R.G. 1976. The Ultrastructure of the Integument of the American Eel, *Anguilla rostrata*. *Cell Tiss. Res.* 171.1-30 (1976)
- Partridge, B. L., Tony J.P. The sensory basis of fish schools: Relative roles of lateral line and vision. *Journal of comparative physiology* 135 (1980): 315-325.
- Matsui, I. 1993. *Theory And Practice Of Eel Culture*. AA. Balkema/Rotterdam.
- Montgomery, J. C., Macdonald, F., Baker, C. F., Carton, A. G. 2002. Hydrodynamic Contributions to Multimodal Guidance of Prey Capture Behavior in Fish. *Brain, Behavior and Evolution*, 59(4), 190–198.
- Munz, H., Claas, B. 1981. The Functional Organization of Neuromasts In the Lateral-Line System of a Cichlio Fish. Fakultät für Biologie, Universität Bielefeld, 0-4800 Bielefeld 1, F.R. of Germany
- Nelson, J.S. 1994. *Fishes Of The World*, 3rd editions. John Wiley & Sons, Inc. New York.

- Northcutt, R.G., Holmes, P.H., Albert, J.S. 2000. Distribution and Innervation of Lateral Line organs in the Channel Catfish. *The Journal of Comparative Neurology* 421:571-522.
- Palstra, A.P., Planas.J.V. 2013. *Swimming Physiology of Fish*. pringer-Verlag Berlin Heidelberg
- Pankhurst, N.W., Lythgoe, J.N. 1982. Structure and Colour of Integument of the European eel *Anguilla Anguilla* (L.). *Journal of Fish Biology*. Vol: 21
- Peach, M. B. Rouse, G. W. 2000. The Morphology of the Pit Organs and Lateral Line Canal Neuromasts of *Mustelus Antarcticus* (Chondrichthyes: Triakidae). *Journal of the Marine Biological Association of the United Kingdom*. 80 (01): 155–162.
- Pichon, F., Ghysen, A. 2004. Evolution of posterior lateral line development in fish and amphibians. *Evolution and Development*, 6(3), 187–193.
- Popper, A.N., C. Plat. 1993. *Inner Ear and Lateral Line. The Physiology of Fishes*. CRC Press. Boca Raton.
- Priatna, Y. 2008. Uji Coba Penentuan Frekuensi Suara Dalam Pemikatan Ikan Mas (*Cyprinus carpio*). Fakultas Perikanan dan Ilmu Kelautan. Institut Pertanian Bogor. Bogor.
- Querat, B., Hardy, A., Fontaine, Y.A. 1991. Regulation of the type-II gonadotrophin α and β subunit mRNAs by α estradiol and testosterone in the European eel. *Journal of Molecular Endocrinology*, 7: 81-86.
- Senturk, G. E., Canilloglu, Y. E. 2014. Which Histochemical Staining Technique Should I Choose for Biological Specimens. *Microscopy: advances in scientific research and education* (A. Méndez-Vilas, Ed.)
- Sheehan, D.C., and Hrapchak, B.B.1987.*Theory And Practice Of Histotechnology*, 2nd Edition. Battelle Memorial Institute. Columbus. OH.
- Sinha, V., J. Jones. 1975. *The European Freshwater Eel*. Liverpool: Liverpool University Press.
- Sugeha, H.Y., Suharti, S.R. 2008. Discrimination and distribution of two tropical short-finned eels (*Anguilla bicolor* and *Anguilla bicolor pacifica*) in the Indonesian Water. *The Nagisa Westpac Congress*: 1-14.
- Tesch, F.W. 2003. *The Eel. Biology and Mangement of Anguilla Eels*. Chapman and Hall. London.
- Webb, J.F. 1989. Neuromast morphology and lateral line trunk canal ontogeny in two species of cichlids: An SEM study. *J. Morphol.*, 202: 53-68. doi:10.1002/jmor.1052020105
- Willemse, J.J. 1979. Guide to The Internal Morphology of European Eel (*Anguilla Anguilla*) (Pisces, Teleostei). *Elsevier Scientific Publishing Company*. Amstredam, Aquaculture, 17(1979) 91-103
- Williams, Terry. 1962. The staining of nervous elements by the Bodian method: The influence of factors preceding impregnation. *Quarterly Journal of Microscopical Science*, Vol. 103, part 2, pp. 155-62



UNIVERSITAS
GADJAH MADA

THE STRUCTURE OF NEUROMAST AND NERVOUS SYSTEM OF LATERAL LINE ON THE DEVELOPMENTAL STAGE OF *Anguilla*

bicolor (McClelland, 1844)

ARIEL HANANYA, Dr. Bambang Retnoaji, M.Sc

Universitas Gadjah Mada, 2020 | Diunduh dari <http://etd.repository.ugm.ac.id/>

Yamada, Y., Hama, K. 1972. *Fine structure of the lateral-line organ of the common eel, *Anguilla japonica**. Zeitschrift For Zellforschung Und Mikroskopische Anatomie, 124(4), 454–464.

Zacchei, A.M., Tovolaro.P. 1998. Lateral Line System During Life Cycle of *Anguilla Anguilla*. *Italian Journal of Zoology*, 55:1-4, 145-153