

DAFTAR PUSTAKA

- Altemimi A, Lakhssassi N, Baharlouei A, Watson DG, Lightfoot DA. Phytochemicals: Extraction, Isolation, and Identification of Bioactive Compounds from Plant Extracts. *Plants (Basel)*. 2017 Sep 22;6(4). pii: E42. doi:10.3390/plants6040042.
- Barara, M., Mendiratta, V., & Chander, R. (2012). Cryotherapy in treatment of keloids: evaluation of factors affecting treatment outcome. *Journal of cutaneous and aesthetic surgery*, 5(3), 185–189. doi:10.4103/0974-2077.101376
- Burrows NP, Lovell CR. Disorders of connective tissue. In : Burns T, Breathnach S, Cox N, editors. *Rook's Textbook of Dermatology*. 8th edition. Wiley-Blackwell, Ltd. 2010. p.45.54-6
- CCRC, *Ciplukan (Physalis angulata L.)*. Dikunjungi 9 November 2019, dari ccrc.farmasi.ugm.ac.id/?page_id=193
- Chen, J.-X. Xue, Hui-Jun Ye, Wen-Cai Fang, Bing-Hu Liu, Ya-Hong Yuan, ShaoHua Yu, Pei Wang, Yu-Qiang. (2009) Activity of physalins purified from *Physalis angulata* in in vitro and in vivo models of cutaneous leishmaniasis, *Biological & pharmaceutical bulletin*, 32(August), doi:10.1248/bpb.32.1385.,
- Chike-Obi, C., Cole, P., & Brissett, A. (2009). Keloids: Pathogenesis, Clinical Features, and Management. *Seminars in Plastic Surgery*, 23(03), 178-184. doi:10.1055/s-0029-1224797.
- Cho SB, Lee JH, Lee SH. (2010). Efficacy and safety of 1064 nm Q-switch Nd:YAG laser with low fluence for keloids and hypertrophic scars. *Journal European Academy of Dermatology and Venerreology*. 24 : 1070-4. DOI:10.1111/j.1468-3083.2010.03593.x
- Darma, A. P., Ashari, R. A., Nugroho, P. A., Monikawati, A., Fauzi, I. A., Hermawan, A., & Meiyanto, E. (2012). Aktivitas Sitotoksik Ekstrak Etanolik Herba Ciplukan (*Physalis Angulata* L.) pada Sel Kanker Leher Rahim HeLa melalui Modulasi Ekspresi Protein p53. *Farmasains : Jurnal Farmasi Dan Ilmu Kesehatan*, 1(2). doi:10.22219/far.v1i2.1163.
- [DEPKES] Departemen Kesehatan Republik Indonesia. 2000. *Parameter Standar Umum Ekstrak Tumbuhan Obat. Edisi I*. Direktorat Jenderal Pengawasan Obat dan Makanan. Direktorat Pengawasan Obat Tradisional. Jakarta.
- España A, Solano T, Quintanilla E. (2001) Bleomycin in the treatment of keloids and hypertrophic scars by multiple needle punctures. *Dermatol Surg*. Jan;27(1):23-7. PubMed PMID: 11231236.
- Hochman B, Locali RF, Matsuoka PK, Ferreira LM. (2008). Intralesional triamcinolone acetate for keloid treatment: a systematic review. *Aesthetic Plast Surg*. Jul;32(4):705-9. doi: 10.1007/s00266-008-9152-8

- Ranti, I., Wahyuningsih, M., & Wirohadidjojo, Y. W. (2018). The antifibrotic effect of isolate tagitinin C from *Tithonia diversifolia* (Hemsley) A. Gray on keloid fibroblast cell. *The Pan African medical journal*, 30, 264. doi:10.11604/pamj.2018.30.264.9994
- Fitri NL, Susetyarini RE, Waluyo Lud. 2016. The Effect Of Ciplukan (*Physalis angulata* L.) Fruit Extract On SGPT and SGOT Levels Against White Male Mice (*Mus musculus*) Hyperglycemia Induced By Alloxan As Biology Learning Resources . *Jurnal Pendidikan Biologi Indonesia*. 2(2):180–187.
- Gauglitz, G. (2013). Management of keloids and hypertrophic scars: Current and emerging options. *Clinical, Cosmetic and Investigational Dermatology*, 103. doi:10.2147/ccid.s35252.
- Gauglitz, G. G., Korting, H. C., Pavicic, T., Ruzicka, T., & Jeschke, M. G. (2010). Hypertrophic Scarring and Keloids: Pathomechanisms and Current and Emerging Treatment Strategies. *Molecular Medicine*, 17(1-2), 113-125. doi:10.2119/molmed.2009.00153.
- Harborne, J.B. 1984. *Phytochemical Methods: A Guide to Modern Technique of Plant Analysis*. (2nd edn). Chapman and Hall. London. 19. Pp.37–168
- Hunasgi, S., Koneru, A., Vanishree, M., & Shamala, R. (2013). Keloid: A case report and review of pathophysiology and differences between keloid and hypertrophic scars. *Journal of Oral and Maxillofacial Pathology*, 17(1), 116. doi:10.4103/0973-029x.110701.
- Imaniar, R., Hartati, W. M. S., & Widodo, W. Y. (2018). The antifibrotic effect of isolate tagitinin C from *Tithonia diversifolia* (Hemsley) A. Gray on keloid fibroblast cell. *Pan African Medical Journal*, 30. doi: 10.11604/pamj.2018.30.264.9994
- Juckett G, Adams HH. Management of keloids and hypertrophic scars. *Am Fam Physician* 2009; 80(3)
- Kalangi, Sonny J. R. (2013). Histofisiologi Kulit. *Jurnal Biomedik Volume 5, Nomor 3*. Manado:Universitas Sam Ratulangi
- Kendall, R., & Feghali-Bostwick, C. (2014). Fibroblasts in fibrosis: novel roles and mediators. *Frontiers In Pharmacology*, 5. doi: 10.3389/fphar.2014.00123.
- Kelly, A. Paul. “Update on the Management of Keloids.” *Seminars in Cutaneous Medicine and Surgery* 28,no. 2 (2009): 71–76. <https://doi.org/10.1016/j.sder.2009.04.002>.
- Khan SA. Keloid an update. *Physician Academy* 2008, 2(6): 25-9 . Diakses 24 Desember 2019, <<http://www.physicians-academy.com/Upload/753676f2-e6b7-4b94-b36f-7b88236d1ef9.pdf>>
- Kim, S.-M., Choi, J.-S., Lee, J.-H., Kim, Y.-J., & Jun, Y.-J. (2014). Prevention of Postsurgical Scars: Comparison of Efficacy and Convenience between Silicone Gel Sheet and Topical Silicone Gel. *Journal of Korean Medical Science*, 29(Suppl 3). doi: 10.3346/jkms.2014.29.s3.s249

- Kischer, C. W., & Hendrix, M. (1983). Fibronectin (FN) in hypertrophic scars and keloids. *Cell and Tissue Research*, 231(1). doi:10.1007/bf00215771.
- Kolarsick, Paul & Kolarsick, Maria & Goodwin, Carolyn. (2011). Anatomy and Physiology of the Skin. *Journal of the Dermatology Nurses' Association*. 3. 203-213. 10.1097/JDN.0b013e3182274a98.
- Kranendonk S, Obagi S. (2011) An algorithmic approach to hypertrophic scars and keloids: Maximizing nonsurgical options. *Cosmetic Dermatology*24(1).
- Lee, S.-S., Yosipovitch, G., Chan, Y.-H., & Goh, C.-L. (2004). Pruritus, pain, and small nerve fiber function in keloids: A controlled study. *Journal of the American Academy of Dermatology*, 51(6), 1002–1006. doi: 10.1016/j.jaad.2004.07.054
- Luliana, S., Susanti, R., & Agustina, E. (2017). Antiinflammatory Activity Test of Aqueous Extracts Herb of Ciplukan (*Physalis angulata* L.) in Caragenan Inducted Wistar Rat (*Rattus norvegicus* L.). *Majalah Obat Tradisional*, 22(3), 199. doi: 10.22146/mot.31556
- Mescher, A. L. (2010). *Junquiera's Basic Histology Text & Atlas 12th ed.* New York. The McGraw-Hill Companies, Inc
- Mouhari-Toure, A., Saka, B., Kombaté, K., Akakpo, S., Egbohou, P., Tchangaï-Walla, K., & Pitche, P. (2012). Is There an Association between Keloids and Blood Groups? *ISRN Dermatology*, 2012, 1–4. doi: 10.5402/2012/750908
- Muhammad, F., Yuliani, F.S., Wahyuningsih, M.S.H., 2017, Aktivitas Antifibrotik Ekstrak Klorofom Nerium indicum dalam Menghambat Proliferasi Fibroblas Keloid dengan MTT Assay, *Prosiding Seminar Nasional "Peran Herbal untuk mencegah Proses Degenerasi*, Auditorium Fakultas Kedokteran UGM, 22 April 2017, ISBN: 978- 602-50277-0-3: p.46-50
- Naeni FF, Najafian J, Ahmadpour K. (2006). Bleomycin tattooing as a promising therapeutic modality in large keloids and hypertrophic scars. *Dermatol Surg*; 32: 1023-30.
- O'Boyle, C. P., Shayan-Arani, H., & Hamada, M. W. (2017). Intralesional cryotherapy for hypertrophic scars and keloids: *A review. Scars, Burns & Healing*, 3, 205951311770216. doi:10.1177/2059513117702162
- Ogawa, R. (2017). Keloid and Hypertrophic Scars Are the Result of Chronic Inflammation in the Reticular Dermis. *International Journal Of Molecular Sciences*, 18(3), 606. doi: 10.3390/ijms18030606
- Ogundajo, A. L., Akpome, A. S., Tijani, N. A., & Ogunwan, I. A. (2015). Chemical Constituents of the Leaf Essential Oil of *Physalis angulata* L. *Asian Journal of Applied Sciences* , 03(04).
- Perdanakusuma, DS. (2012). Penanganan Parut Hipertrofik dan Keloid. Dalam: *Penanganan Parut Hipertrofik dan Keloid*. Airlangga University Press, Fakultas Kedokteran Unair, pp. 1-131. ISBN 978-602-6606-40-2

- Prachyapruit W, Vashrangsi N. A new therapeutic modality intralesional 5-fluorouracil in the treatment of keloids and hypertrophic scar. *Institute of Dermatology* 2010; (10)2: 157-63. DOI 10.1007/s00266-006-0253-y
- Prayong, P., S. Barusrux, and N. Weerapreeyakul. "Cytotoxic Activity Screening of Some Indigenous Thai Plants." *Fitoterapia* 79, no. 7-8 (2008): 598–601. <https://doi.org/10.1016/j.fitote.2008.06.007>.
- Profyris, C., Tziotzios, C., & Sterling, J. (2012). Cutaneous scarring: Pathophysiology, molecular mechanisms, and scar reduction therapeutics. *Journal of the American Academy of Dermatology*, 66(1), 13–24. doi: 10.1016/j.jaad.2011.08.035
- Purnama, H., Sriwidodo, S., Mita SR. (2017). Proses Penyembuhan dan Perawatan Luka : Review Sistematis. *Jurnal Farmaka Universitas Padjadjaran*, 15(02) doi:10.24198/jf.v15i2.13366
- Putra, IB., Jusuf, NK. Studi Retrospektif Keloid di Poliklinik Kulit dan Kelamin RSUP H. Adam Malik Medan. *12th Asia-Pacific Environmental and Occupational Dermatology Symposium*; 2013 Oct 23-26; Yogyakarta: Indonesia
- Putra IB. Pengaruh ekstrak kelopak bunga rosella (*Hibiscus sabdariffa l*) terhadap penghambatan proliferasi sel fibroblast keloid manusia. *Disertasi. Medan : Program Doktor S3 Ilmu Kedokteran Fakultas Kedokteran Universitas Sumatra Utara*; 2012
- Ramakrishnan, K. M., Thomas, K. P., & Sundararajan, C. R. (1974). Study Of 1,000 Patients With Keloids In South India. *Plastic and Reconstructive Surgery*, 53(3), 276–280. doi: 10.1097/00006534-197403000-00004
- Rengifo, E., Vargas-arana, G. dan Científica, U. (2013) *Physalis angulata L. (Bolsa Mullaca)*.: A review of its traditional uses, *Chemistry and Pharmacology*, (September), Diakses 1 Oktober 2019, <https://www.researchgate.net/publication/260284500_Physalis_angulata_L_Bolsa_Mullaca_A_Review_of_its_Traditional_Uses_Chemistry_and_Pharmacology>.
- Robles, D. T., & Berg, D. (2007). Abnormal wound healing: Keloids. *Clinics in Dermatology*, 25(1), 26-32. doi:10.1016/j.clindermatol.2006.09.009
- Robles DT, Moore E, Draznin M. (2007) Keloids: Pathophysiology and management. *Dermatology Online Journal*; 13(3): 9-13.
- Rosita, S.M.D., Rostiana, O., Pribadi, dan Hernani, (2007), "Penggalian IPTEK Etnomedisin di Gunung Gede Pangrango". *Bul. Littro*. 18 (1): 13-28, Diakses: 22 Oktober 2017, <<http://ejurnal.litbang.pertanian.go.id/index.php/bultro/article/view/1925>>.
- Rusciani L, Paradisi A, Alfano C, Chiummariello S, Rusciani A. Cryotherapy in the treatment of keloids. *J Drugs Dermatol*. 2006 Jul-Aug;5(7):591-5.
- Seifert, O., & Mrowietz, U. (2009). Keloid scarring: bench and bedside. *Archives of Dermatological Research*, 301(4), 259–272. doi: 10.1007/s00403-009-0952-8

- Shridharani, MS., Magarakis, M.(2010) Emerging role of antineoplastic agents in the treatment of keloids and hypertrophic scars. *Annals of Plast Surg*: 64: 355-61. DOI: 10.1097/SAP.0b013e3181afaab0.
- Sutrisna, EM., Indwianistuti, Haryadi 2012. The Ethanol Extract of *Physalis angulata* Linn Inhibits COX-2 Activity in MCF-7 Cell In Vitro . *Surakarta: Jurnal kedokteran Universitas Muhammadiyah Surakarta* Vol. 3 No.1.
- Ukwubile, C.A., & Oise, I.E. (2016). Analgesic and Anti-inflammatory Activity of *Physalis angulata* Linn. (Solanaceae) Leaf Methanolic Extract in Swiss Albino Mice. *International Biology Biomedical Journal*. 2(14).
- Urioste SS, Arndt KA, Dover JS. (1999). Keloid and hypertrophic scars: Review and treatment strategies. *Seminars in Cutaneous Medicine and Surgery* 18(2): 159-71
- USDA, NRCS. 2020. *The PLANTS Database* (<http://plants.usda.gov>, 21 January 2020). National Plant Data Team, Greensboro, NC 27401-4901 USA
- Wahyuningsih, MSH., Wirohadidjojo, YW., Hidayat R., Sadid A. Antifibrotic Effect of Standardized Ethanol Extract of *Tithonia diversifolia* (Hemsley) A. Gray on Keloid Fibroblasts. *International Journal of Pharmacognosy and Phytochemical Research*. 2015; 7(4); 642-647. ISSN: 0975-4873
- Wahyuningsih, M. S. H., Yuliani, F. S., Rahmawati, D. Y., & Pratiwi, A. N. (2018). Antifibrotic Effect of Ethanolic Extract of *Nerium indicum* Mill. Standardized 5 α -Oleandrin on Keloid Fibroblasts Cells. *Majalah Obat Tradisional*, 23(1), 70. doi: 10.22146/mot.35116
- Wahyuningsih, M. S. H., Nugrahaningsih, D. A. A., & Budiyanto, A. (2019). Ethanolic Extract Of *Tithonia diversifolia* (Hemsley) A. Gray Inhibits Migration Activity And Decrease The Transforming Growth Factor-Beta1, Vegf Expression On Keloid Fibroblasts. *Asian Journal of Pharmaceutical and Clinical Research*, 12(1), 342. doi: 10.22159/ajpcr.2018.v12i1.29850
- Wolfram, D., Tzankov, A., Pülzl, P., & Piza-Katzer, H. (2009). Hypertrophic Scars and Keloids—A Review of Their Pathophysiology, Risk Factors, and Therapeutic Management. *Dermatologic Surgery*, 35(2), 171–181. doi: 10.1111/j.1524-4725.2008.34406.x
- Xi-Qiao, W., Ying-Kai, L., Chun, Q., & Shu-Liang, L. (2009). A Review of the Effectiveness of Antimitotic Drug Injections for Hypertrophic Scars and Keloids. *Annals of Plastic Surgery*, 63(6), 688–692. doi: 10.1097/sap.0b013e3181978753
- Vitasari, O. N. (2012). Uji Aktivitas Antibakteri Ekstrak Etanol Daun Ciplukan (*Physalis angulata* L.) terhadap *Staphylococcus aureus* dan *Pseudomonas aeruginosa*. Skripsi. Fakultas Matematika dan Ilmu Pengetahuan Alam, Universitas Sebelas Maret, Surakarta.
- Zhang, Q., Lin, L. & Ye, W. Techniques for extraction and isolation of natural products: a comprehensive review. *Chin Med* 13, 20 (2018) doi:10.1186/s13020-018-0177-x



UNIVERSITAS
GADJAH MADA

Aktivitas Ekstrak Metanol Ciplukan (*Physalis angulata* L.) terhadap Proliferasi Fibroblas Keloid dengan Metode MTT

FADHLUR ROHMAN NOOR IKBAR, Prof. Dr. Mae Sri Hartati W, Apt, M.Si;Dr. dr. Rul Afiyah Syarif, M.Kes;Dr. Med. (c)
Universitas Gadjah Mada, 2020 | Diunduh dari <http://etd.repository.ugm.ac.id/>